

UNIVERSIDAD DE CANTABRIA



E.T.S. INGENIEROS DE CAMINOS, CANALES Y PUERTOS.

DPTO. DE CIENCIAS Y TÉCNICAS DEL AGUA Y DEL MEDIO AMBIENTE.

AREA DE ECOLOGÍA

TESIS DOCTORAL

CONTRIBUCIÓN AL CONOCIMIENTO DEL ERIZO DE MAR *Paracentrotus lividus* (LAMARCK, 1816) EN EL MAR CANTÁBRICO: CICLO GONADAL Y DINÁMICA DE POBLACIONES

Presentada por: **JOSÉ MANUEL GONZÁLEZ IRUSTA**

Dirigida por: **JUAN CARLOS CANTERAS JORDANA**
GERARDO GARCÍA-CASTRILLO RIESGO

Santander, julio 2009

CAPÍTULO 6

BIBLIOGRAFÍA



Foto de Javier Ansorena

Esta tesis doctoral nunca habría podido realizarse sin todas las personas que a continuación se citan:

CAPÍTULO 6: BIBLIOGRAFÍA

1. **Aksland M (1994)**. A general cohort analysis method. *Biometrics* 50(4): 917-932.
2. **Allain JY (1972)** Structure des populations de *Paracentrotus lividus* (Lamarck), (Echinodermata, Echinoidea) soumises à la pêche sur les côtes du Nord de Bretagne. *Revue des Travaux de l'Institut des Pêches Maritimes* 39(2): 171-212.
3. **Allain JY (1975)**. Structure des populations de *Paracentrotus lividus* soumises à la pêche sur les côtes Nord de Bretagne. *Rev Trav. Inst. Pêches Marit*, 39(2): 171-212.
4. **Allain JY (1978)**. Age et croissance de *Paracentrotus lividus* (Lmk.) et de *Psammechinus miliaris* (Gmelin) des côtes Nord de Bretagne (Echinoidea). *Cah. Biol. Mar.* 19: 11-21.
5. **Andrew NL, Agatsuma Y, Ballesteros E, Bazhin AG, Creaser EP, Barnes DKA, Botsford LW, Bradbury A, Campbell AC, Dixon JD (2002)**. Status and management of world sea urchin fisheries. *Oceanography and Marine Biology Annual Review* 40: 343-425.
6. **Appeldoorn RS (1987)**. Modification of a seasonally oscillating growth function for use with mark-recapture data. *ICES J Mar Sci* 43(3): 194-198.
7. **Arafa S, Sadok S, El Abed A (2006)**. Variation in nitrogenous compounds and gonad index in fed and starved sea urchins (*Paracentrotus lividus*) during live storage. *Aquaculture*, 257(1-4): 525-533.

8. **Azzolina JF y Willsie A (1987)**. Abundance des juveniles de *Paracentrotus lividus* au sein de l'herbier a posidonie oceanica. In : Boudouresque, GIS Posidonie, IFREMER (eds.). *Colloque International sur Paracentrotus lividus et les oursins comestibles*. Marseille: 159-167.
9. **Azzolina JF (1988)**. Contribution a l'etude de la dynamique des populations de l'oursin comestible *Paracentrotus lividus* (Lmck). Croissance, recrutement, mortalite, migrations.
10. **Balch T y Scheibling RE (2000)**. Temporal and spatial variability in settlement and recruitment of echinoderms in kelp beds and barrens in Nova Scotia. *Mar. Ecol. Prog. Ser.* 205: 139-154.
11. **Ballesteros, E. y García-Rubies, A. (1987)**. La peche aux oursins en Espagne et plus particulièrement en Catalogne. In : Boudouresque, GIS Posidonie, IFREMER (eds.). *Colloque International sur Paracentrotus lividus et les oursins comestibles*. Marseille: 325-328.
12. **Barbaglio A, Sugni M, Di Benedetto C, Bonasoro F, Schnell S, Lavado R, Porte C, Candia Carnevali DM (2007)**. Gametogenesis correlated with steroid levels during the gonadal cycle of the sea urchin *Paracentrotus lividus* (Echinodermata: Echinoidea). *Comparative Biochemistry and Physiology, Part A* 147(2): 466-474.
13. **Barnes DKA y Crook AC (2001)**. Implications of temporal and spatial variability in *Paracentrotus lividus* populations to the associated commercial coastal fishery. *Hydrobiologia* 465: 95-102.
14. **Barnes DKA, Verling E, Crook AC, Davidson I, O'Mahoney M (2002)**. Local population disappearance follows (20 yr after) cycle collapse in a pivotal ecological species. *Marine Ecology Progress Series* 226: 311-313.

-
15. **Bayed A, Quiniou F, Benrha A, Guillou M (2005).** The *Paracentrotus lividus* populations from the northern Moroccan Atlantic coast: Growth, reproduction and health condition. *Journal of the Marine Biological Association of the United Kingdom* 85(4): 999-1007.

 16. **Bay-Schmith E y Pearse JS (1987).** Effect of fixed daylengths on the photoperiodic regulation of gametogenesis in the sea urchin *Strongylocentrotus purpuratus*. *International Journal of Invertebrate Reproduction and Development* 11(3): 287-294.

 17. **Beck MW (1995).** Size-specific shelter limitation in stone crabs: A test of the demographic bottleneck hypothesis. *Ecology*: 968-980.

 18. **Benedetti-Cecchi L y Cinelli F (1995).** Habitat heterogeneity, sea urchin grazing and the distribution of algae in littoral rock pools on the west coast of Italy (western Mediterranean). *Mar Ecol. Prog. Ser.* 126(1): 203-212.

 19. **Bhattacharya CG (1967).** A simple method of resolution of a distribution into Gaussian components. *Biometrics* 23(1): 115-135.

 20. **Bode A, Casas B, Fernández E, Marañón E, Serret P, Varela M (1996).** Phytoplankton biomass and production in shelf waters off NW Spain: Spatial and seasonal variability in relation to upwelling. *Hydrobiologia* 341(3): 225-234.

 21. **Bonnet A (1925).** Documents pour servir a l'étude des variations chez les echinides. *Bull Inst Oceanogr* 462: 1-28.

 22. **Botas JA, Fernández E, Bode A, Anadón R (1990).** A persistent upwelling off the central Cantabrian coast (Bay of Biscay). *Estuarine, Coastal and Shelf Science*. 30(2): 185-199.

- 23. Böttger SA, Devin MG, Walker CW (2006).** Suspension of annual gametogenesis in North American green sea urchins (*Strongylocentrotus droebachiensis*) experiencing invariant photoperiod. Applications for land-based aquaculture. *Aquaculture* 261(4): 1422-1431.
- 24. Boudouresque CF, Verlaque M, Azzolina JF, Meinesz A, Nedelec H, Rico V (1989).** Evolution des populations de *Paracentrotus lividus* et d' *Arbacia lixula* (Echinoidea) le long d' un transect permanent a Galeria (Corse). *Trav. Sci. Parc. Nat. Reg. Nat. Corse* 22: 65-82.
- 25. Boudouresque C y Verlaque M. (2001).** Chapter 13: Ecology of *Paracentrotus lividus*. In: *Edible sea urchins: Biology and Ecology* John M. Lawrence (ed), pp 177-216.
- 26. Brewin PE, Lamare MD, Keogh JA, Mladenov PV (2000).** Reproductive variability over a four-year period in the sea urchin *Evechinus chloroticus* (Echinoidea: Echinodermata) from differing habitats in New Zealand. *Mar. Biol.* 137(3): 543-557
- 27. Briscoe, C.S. y Sebens, K.P. (1988).** Omnivory in *Strongylocentrotus droebachiensis* (Mueller) (Echinodermata: Echinoidea): Predation on subtidal mussels. *J. Exp. Mar. Biol. Ecol.* 115(1): 1-24
- 28. Brockington S y Clarke A (2001).** The relative influence of temperature and food on the metabolism of a marine invertebrate. *J. Exp. Mar. Biol. Ecol.* 258(1): 87-99
- 29. Brody S. (1945).** *Bioenergetics and growth.* Reinhold, New York.

-
- 30. Buckle L.F., Guisado CH, Tarifeno E, Zuleta A, Cordova L, Serrano C (1978).** Biological studies on the sea urchin *Loxechinus albus* (Molina) (Echinodermata; Echinoidea) IV. Maturation cycle and seasonal biochemical changes in the gonad. *Cien. Mar.* 5(1): 1-18
- 31. Bulleri F, Benedetti-Cecchi L, Cinelli F (1999).** Grazing by the sea urchins *Arbacia lixula* L. and *Paracentrotus lividus* Lam. in the northwest Mediterranean. *J. Exp. Mar. Biol. Ecol.* 241(1): 81-95.
- 32. Byrne M (1990).** Annual reproductive cycles of the commercial sea urchin *Paracentrotus lividus* from an exposed intertidal and a sheltered subtidal habitat on the west coast of Ireland. *Marine Biology* 104(2): 275-289.
- 33. Caltagirone A, Francour P, Fernandez C. (1992).** Formulation of an artificial diet for the rearing of the urchin *Paracentrotus lividus*: I. Comparison of different binding agents. In: *Echinoderm Research*. Scalera-Liaci, L. y Canicatti, C. (ed) Balkema, Rotterdam, pp 115.
- 34. Cardona L, Sales M, López D (2007).** Changes in fish abundance do not cascade to sea urchins and erect algae in one of the most oligotrophic parts of the mediterranean. *Estuar. Coast Shelf Sci.* 72(1-2): 273-282.
- 35. Catoira JL (1991)** Harvesting sea urchins in Galicia Spain during the 1990-1991 season. *Echinoderm Research*, 1991: 199-200.
- 36. Catoira JL (1995).** Spatial and temporal evolution of the gonad index of the sea urchin *Paracentrotus lividus* (Lamarck) in Galicia, Spain. *Echinoderm Research* 1995 : 295-298.
- 37. Catoira JL (2004).** History and current state of sea urchin *Paracentrotus lividus* (Lamarck, 1816) fisheries in Galicia, NW Spain. *Sea Urchins: Fisheries and Ecology* : 64-73.

- 38. Cellario C y Fenaux L (1990).** *Paracentrotus lividus* (Lamarck) in culture (larval and benthic phases): Parameters of growth observed during two years following metamorphosis. *Aquaculture*, 84(2): 173-188.
- 39. Christie H, Fredriksen S, Rinde E (1998)** Regrowth of kelp and colonization of epiphyte and fauna community after kelp trawling at the coast of Norway. *Hydrobiologia* 375-376: 49-58.
- 40. Clemente S, Hernández JC, Toledo K, Brito A (2007).** Predation upon *Diadema affz. antillarum* in barren grounds in the Canary Islands. *Scientia Marina* 71(4).
- 41. Conand, C. y Sloan, N.A. (1989).** World fisheries for echinoderms. In: *Marine invertebrate fisheries: Their assessment and management*. John Wiley & Sons, New York ed. pp 647-663
- 42. Cook EJ y Kelly MS (2007) a.** Effect of variation in the protein value of the red macroalga *Palmaria palmata* on the feeding, growth and gonad composition of the sea urchins *Psammechinus miliaris* and *Paracentrotus lividus* (Echinodermata). *Aquaculture* 270(1-4): 207-217.
- 43. Cook EJ y Kelly MS (2007) b.** Enhanced production of the sea urchin *Paracentrotus lividus* in integrated open-water cultivation with Atlantic salmon *Salmo salar*. *Aquaculture*, 273(4): 573-585.
- 44. Crapp GB y Willis ME (1975).** Age determination in the sea urchin *Paracentrotus lividus* (Lamarck), with notes on the reproductive cycle. *Journal of Experimental Marine Biology and Ecology* 20(2): 157-178.

- 45. Dafni, J. & Tobol, R.** Population structure patterns of a common red sea echinoid (*Tripneustes gratilla elatensis*). Israel J. Zool. 34: 191-204.
- 46. Dance C (1987).** Patterns of activity of the sea urchin *Paracentrotus lividus* in the Bay of Port-Cros (France, Mediterranean). Mar. Ecol.8(2): 131-142.
- 47. Daza Cordero JL, Del Castillo y Rey, F., Márquez Pascual I. (2002).** La pesquería del erizo y la anémona de mar en el litoral de Cádiz y Málaga. Ed. Consejería de Agricultura y Pesca. Junta de Andalucía.
- 48. Delmas P y Régis MB. (1992)** Etude des populations de *Paracentrotus lividus* (Lam.) (Echinodermata: Echinoidea) soumises à une pollution complexe en provence nord-occidentale densités, structures, processus de détoxification (zn, cu, pb, cd, fe). Tesis doctoral. Universidad Aix-Marseille 3.
- 49. Diez I, Secilla A, Santolaria A, Gorostiaga JM (2000).** The north coast of Spain. Seas at the Millennium - an Environmental Evaluation - Volume 1 : 135-150.
- 50. Domenici P, González-Calderón D, Ferrari RS (2003).** Locomotor performance in the sea urchin *Paracentrotus lividus*. J. Mar. Biol. Assoc UK 83(2): 285-292.
- 51. Dominique F (1973)** Contribution a l'étude du cycle annuel de reproduction de deux espèces d'échinoides (echinodermata) des cotes de Bretagne. Thesis, Université Libre de Bruxelles.
- 52. Dumont C, Himmelman JH, Russell MP (2004).** Size-specific movement of green sea urchins *Strongylocentrotus droebachiensis* on urchin barrens in eastern Canada. Mar. Ecol. Prog. Ser. 276(1): 93-101.

- 53. Ebert, T (1967).** Negative growth and longevity in the purple sea urchin *Strongylocentrotus purpuratus* (Stimpson). *Science* Vol 157: 557-558.
- 54. Ebert TA (1973).** Estimating growth and mortality rates from size data. *Oecologia* 11: 281-298.
- 55. Ebert TA (1980).** Estimating parameters in a flexible growth equation, the Richards function. *Can. J. Fish. Aquat. Sci.* 37(4): 687-692.
- 56. Ebert TA (1982).** Longevity, life history, and relative body wall size in sea urchins. *Ecol. Monographs* 52(4): 353-394.
- 57. Ebert TA, Dixon JD, Schroeter SC, Kalvass PE, Richmond NT, Bradbury WA, Woodby DA (1999).** Growth and mortality of red sea urchins *Strongylocentrotus franciscanus* across a latitudinal gradient. *Marine Ecology Progress Series* 190: 189-209.
- 58. Fabens AJ (1965).** Properties and fitting of the von Bertalanffy growth curve. *Growth* 29: 265-289.
- 59. Fenaux L (1968).** Maturation des gonades et cycle saisonnier des larves chez *A. lixula*, *P. lividus* et *P. microtuberculatus* (echinides) à Villafranche-Sur-Mer. *Vie Milieu* 19: 1-52.
- 60. Fenaux L, Etienne M, Quelart G (1987).** Suivi écologique d'un peuplement de *Paracentrotus lividus* (Lamarck) dans la baie de Villefranche Sur Mer. : 187–197.
- 61. Fenaux L y Pedrotti ML (1988)** Metamorphosis of echinoid larvae in midwater. *Marine Ecology* 9(2): 93-107.

-
- 62. Fernandez C y Caltagirone A (1994)** Growth rate of adult sea urchins, *Paracentrotus lividus* in a lagoon environment: The effect of different diet types. Echinoderms through time. Balkema, Rotterdam: 655-660.
- 63. Fernandez C y Boudouresque CF (1997).** Phenotypic plasticity of *Paracentrotus lividus* (Echinodermata: Echinoidea) in a lagoonal environment. Marine Ecology Progress Series 152(1-3): 145-154.
- 64. Fernandez C y Boudouresque CF (2000).** Nutrition of the sea urchin *Paracentrotus lividus* (Echinodermata: Echinoidea) fed different artificial food. Marine Ecology Progress Series 204: 131-141.
- 65. Fernandez C, Caltagirone A, Johnson M (2001)** Demographic structure suggests migration of the sea urchin *Paracentrotus lividus* in a coastal lagoon. J Mar. Biol. Assoc U K 81(2): 361-362.
- 66. Fernandez C, Pasqualini V, Boudouresque CF, Johnson M, Ferrat L, Caltagirone A, Mouillot D (2006).** Effect of an exceptional rainfall event on the sea urchin (*Paracentrotus lividus*) stock and seagrass distribution in a Mediterranean coastal lagoon. Estuar Coast Shelf Sci 68(1-2): 259-270.
- 67. Ford E (1933).** An accounting of the herring investigations conducted at Plymouth during the years from 1924-1933. J Mar. Biol. Assoc UK 19: 305-384.
- 68. Fuji A (1960).** Studies on the biology of the sea urchin. III. reproductive cycle of two sea urchins, *Strongylocentrotus nudus* and *S. intermedius* in southern Hokkaido. Bull. Fac. Fish. Hokkaido Univ. 11: 49-57.
- 69. Fuji A (1967).** Ecological studies on the growth and food consumption of Japanese common littoral sea urchin, *Strongylocentrotus intermedius* (A. Agassiz). Mem. Fac. Fish Hokkaido Univ. 15(2): 1-160.

- 70. Fujisawa H y Shigei M (1990).** Correlation of embryonic temperature sensitivity of sea urchins with spawning season. *J. Exp. Mar. Biol. Ecol.*136(2): 123-139.
- 71. Gabín-Sánchez C y Lorenzo De Dios F (1993).** El erizo de mar un recurso con futuro. Aula del mar, Fundación Caixa Galicia.
- 72. Gage JD y Tyler PA (1985).** Growth and recruitment of the deep-sea urchin *Echinus affinis*. *Mar. Biol.* 90(1): 41-53.
- 73. Gage JD y Tyler, P.A. & Nichols, D. (1986).** Reproduction and growth of *Echinus acutus* var. *norvegicus* Düben & Koren and *E. elegans* Düben & Koren on the continental slope off Scotland. *J. Exp. Mar. Biol. Ecol.*101: 61-83.
- 74. Gage JD (1987).** Growth of the deep-sea irregular sea urchins *Echinostigma phiale* and *Hemiaster expergitus* in the rockall trough (NE Atlantic Ocean). *Mar. Biol.* 96(1): 19-30.
- 75. Gage JD (1991)** Skeletal growth zones as age-markers in the sea urchin *Psammechinus miliaris*. *Mar. Biol.* 110(2): 217-228.
- 76. Gago L, Range P, Luis O (2003).** Growth, reproductive biology and habitat selection of the sea urchin *Paracentrotus lividus* in the coastal waters of Cascais, Portugal.
- 77. Gallucci, V.F. y Quinn, T.J. (1979).** Reparameterizing, fitting, and testing a simple growth model. *Trans Am Fish Soc*, 108: 14-25.
- 78. García-Soto C (2006).** Blooms de fitoplancton (SeaWiFS) en el golfo de Vizcaya y canal de la mancha a lo largo de la ruta del ferry Pride of Bilbao. *Revista De Teledetección* : 10-14

-
- 79. Gayanilo FC, Sparre P, Pauly D. (2005).** FAO-ICLARM stock assessment tools II: User's guide. World fish Center.
- 80. Gianguzza P, Chiantore M, Bonaviri C, Cattaneo-Vietti R, Vielmini I, Riggio S (2006).** The effects of recreational *Paracentrotus lividus* fishing on distribution patterns of sea urchins at Utica island MPA (western Mediterranean, Italy). Fisheries Research 81(1): 37-44.
- 81. Girard D, Hernández JC, Toledo K, Clemente S, Br (2006)** Aproximación a la biología reproductiva del equinoideo *Paracentrotus lividus* (Lamarck 1816) en el litoral de Tenerife. Libro de Resúmenes XIV SIEBM.
- 82. Gonor JJ (1972).** Gonad growth in the sea urchin, *Strongylocentrotus purpuratus* (Stimpson) (Echinodermata: Echinoidea), and the assumptions of gonad index methods. J. Exp. Mar. Biol. Ecol.10: 89-103.
- 83. Gonor JJ (1973).** Reproductive cycles in Oregon populations of the echinoid, *Strongylocentrotus purpuratus* (Stimpson) I. Annual gonad growth and ovarian gametogenic cycles. Journal of Experimental Marine Biology and Ecology 12(1): 45-64.
- 84. Gosselin P y Jangoux M (1995).** Induction of metamorphosis in *Paracentrotus lividus* larvae (Echinodermata, Echinoidea). Oceanol Acta 19: 292-296.
- 85. Grosjean P, Spirlet C, Jangoux M (1996).** Experimental study of growth in the echinoid *Paracentrotus lividus* (Lamarck, 1816) (Echinodermata). Journal of Experimental Marine Biology and Ecology 201(1-2): 173-184.
- 86. Grosjean P (2001).** Growth model of the reared sea urchin *Paracentrotus lividus*. Ph. D. Thesis.

- 87. Grosjean P, Spirlet C, Jangoux M (2003).** A functional growth model with intraspecific competition applied to a sea urchin, *Paracentrotus lividus*. Canadian Journal of Fisheries and Aquatic Sciences 60(3): 237-246.
- 88. Guettaf M y San Martin GA (1995).** Study of gonad index variability of the edible sea-urchin *Paracentrotus lividus* (Echinodermata: Echinidae) in the north-western Mediterranean. Vie Milieu 45(2): 129-137.
- 89. Guettaf M (1997).** Contribution à l'étude de la variabilité du cycle reproductif (indice gonadique et histologie des gonades) chez *Paracentrotus lividus* (Echinodermata: Echinoidea) en Méditerranée sud occidentale (Algérie). These, Universite de la Mediterranee Aix-Marseille III.
- 90. Guettaf M, San Martin GA, Francour P (2000).** Interpopulation variability of the reproductive cycle of *Paracentrotus lividus* (Echinodermata: Echinoidea) in the south-western Mediterranean. Journal of the Marine Biological Association of the United Kingdom 80(5): 899-907.
- 91. Guidetti P, Frascchetti S, Terlizzi A, Boero F (2003)** Distribution patterns of sea urchins and barrens in shallow Mediterranean rocky reefs impacted by the illegal fishery of the rock-boring mollusc *Lithophaga lithophaga*. Mar. Biol. 143(6): 1135-1142.
- 92. Guidetti P, Terlizzi A, Boero F (2004).** Effects of the edible sea urchin, *Paracentrotus lividus*, fishery along the Apulian rocky coast (SE Italy, Mediterranean sea). Fisheries Research 66(2-3): 287-297.
- 93. Guidetti P y Dulčić J (2007).** Relationships among predatory fish, sea urchins and barrens in Mediterranean rocky reefs across a latitudinal gradient. Mar. Environ. Res. 63(2): 168-184.

-
- 94. Guillou M y Michel C (1993).** Reproduction and growth of *Sphaerechinus granularis* (Echinodermata: Echinoidea) in southern Brittany. Journal of Marine Biological Association of the United Kingdom 73(1): 179-192.
- 95. Guillou M y Michel C (1994).** The influence of environmental factors on the growth of *Sphaerechinus granularis* Lamarck) (Echinodermata: Echinoidea). J. Exp. Mar. Biol. Ecol.178(1): 97-111.
- 96. Guillou M y Lumingas LJJL (1998).** The reproductive cycle of the ‘blunt’ sea urchin. Aquacult. Int. 6(2): 147-160.
- 97. Gulland JA y Holt SJ (1959).** Estimation of growth parameters for data at unequal time intervals. ICES J Mar Sci 25(1): 47.
- 98. Hagen NT (1996)** Echinoculture: From fishery enhancement to closed cycle cultivation. World Aquac. 27(4): 6-19.
- 99. Harrington LH, Walker CW, Lesser MP (2007).** Stereological analysis of nutritive phagocytes and gametogenic cells during the annual reproductive cycle of the green sea urchin, *Strongylocentrotus droebachiensis*. Invertebr. Biol. 126(2): 202-209.
- 100. Harrold C, Lisin S, Light KH, Tudor S (1991).** Isolating settlement from recruitment of sea urchins. J. Exp. Mar. Biol. Ecol.147(1): 81-94.
- 101. Haya de la Sierra, D. (1990).** Biología y Ecología de *Paracentrotus lividus* en la zona intermareal. Tesis doctoral, Universidad de Oviedo.
- 102. Hereu B, Zabala M, Linares C, Sala, E (2004).** Temporal and spatial variability in settlement of the sea urchin *Paracentrotus lividus* in the NW Mediterranean. Mar. Biol. 144(5): 1011-1018.

103. **Hereu B, Zabala M, Linares C, Sala E (2005).** The effects of predator abundance and habitat structural complexity on survival of juvenile sea urchins. *Mar. Biol.* 146(2): 293-299.
104. **Highsmith RC (1982).** Induced settlement and metamorphosis of sand dollar (*Dendraster ecentricus*) larvae in predator-free sites: Adult sand dollar beds. *Ecology*: 329-337.
105. **Himmelman JH. (1981).** Synchronization of spawning in marine invertebrates by phytoplankton. *Advances in invertebrates reproduction. Vol II.* Elsevier North-Holland, New York, NY, pp 3-19.
106. **Hixon MA y Beets JP (1993).** Predation, prey refuges, and the structure of coral-reef fish assemblages. *Ecol.Monogr*: 77-101.
107. **Jacquin AG, Donval A, Guillou J, Leyzour S, Deslandes E, Guillou M (2006).** The reproductive response of the sea urchins *Paracentrotus lividus* (G.) and *Psammechinus miliaris* (L.) to a hyperproteinated macrophytic diet. *J. Exp. Mar. Biol. Ecol.* 339(1): 43-54.
108. **James PJ, Heath P, Unwin MJ (2007).** The effects of season, temperature and initial gonad condition on roe enhancement of the sea urchin *Evechinus chloroticus*. *Aquaculture* 270(1-4): 115-131.
109. **Jensen M (1981).** The Strongylocentrotidae, a morphologic and systematic study. *Sarsia* 57: 113-148.
110. **Jensen M (1969).** Age determination of echinoids. *Sarsia* 37: 41-44

-
111. **Jolicoeur P (1985).** Flexible 3-parameter curve for limited or unlimited somatic growth. *Growth* 49: 271-281.
112. **Keats DW, Steele DH, South GR (1984).** Depth-dependent reproductive output of the green sea urchin, *Strongylocentrotus droebachiensis* (O.F. Mueller), in relation to the nature and availability of food. *J. Exp. Mar. Biol. Ecol.* 80: 77-91.
113. **Keesing JK, Cartwright CM, Hall KC (1993).** Measuring settlement intensity of echinoderms on coral reefs. *Mar. Biol.* 117(3): 399-408.
114. **Keesing JK y Hall KC (1998).** Review of harvests and status of world sea urchin fisheries points to opportunities for aquaculture. *J Shellfish Res.* 17: 1597-1608.
115. **Kelly MS (2000).** The reproductive cycle of the sea urchin *Psammechinus miliaris* (Echinodermata: Echinoidea) in a Scottish sea loch. *Journal of the Marine Biological Association of the UK* 80(05): 909-919.
116. **Kelly MS (2001).** Environmental parameters controlling gametogenesis in the echinoid *Psammechinus miliaris*. *J. Exp. Mar. Biol. Ecol.* 266(1): 67-80.
117. **Kempf M (1969).** Recherches d'écologie comparée sur *Paracentrotus lividus* et *Arbacia lixula*. *Rec. Trav. St. Mar. Endoume* 25(39): 45-115.
118. **King CK, Hoegh-Guldberg O, Byrne M (1994).** Reproduction cycle of *Centrostephanus rogersii* (Echinoidea) with recommendations for the establishment of a sea urchin fishery in New South-Wales.

- 119. Lamare MD y Mladenov PV (2000).** Modelling somatic growth in the sea urchin *Evechinus chloroticus* (Echinoidea: Echinometridae). *Journal of Experimental Marine Biology and Ecology* 243(1): 17-43.
- 120. Lamare MD y Barker MF (2001)** Settlement and recruitment of the New Zealand sea urchin *Evechinus chloroticus*. *Mar. Ecol. Prog. Ser.* 218: 153-166.
- 121. Lawrence, J.M. & Lane, J.M. (1982).** The utilization of nutrients by postmetamorphic echinoderms. In: M. Jangoux & J.M. Lawrence (ed) *Echinoderm nutrition*. Balkema, Rotterdam, pp 331-371.
- 122. Lawrence JM. (2001).** The edible sea urchins. In: Lawrence JM (ed) *Edible sea urchins: Biology and ecology*. pp 1
- 123. Le Direach L, Boudouresque CF, Antolic B, Kocatas A, Panayotidis P, Pancicci A, Semroud R, Span A, Zaquali J, Zavodnik D (1987).** Rapport sur l'exploitation des oursins en Méditerranée. In : Boudouresque, GIS Posidonie, IFREMER (eds.). *Colloque International sur Paracentrotus lividus et les oursins comestibles*. Marseille: 199-222.
- 124. Le Gall (1987).** La peche des oursins en Bretagne. In : Boudouresque, GIS Posidonie, IFREMER (eds.). *Colloque International sur Paracentrotus lividus et les oursins comestibles*. Marseille: 311-324.
- 125. Le Gall P y Bucaille D (1990).** Influence de la temperature sur la croissance de deux oursins comestibles, *Paracentrotus lividus* et *Psammechinus miliaris*. In: *Echinoderms research*, De Ridder C, Dubois P, Lahaye MC, Jangoux M (eds) pp 183-188.

-
126. **Lessios HA (1988).** Mass mortality of *Diadema antillarum* in the Caribbean: What have we learned? *Annu Rev Ecol.Syst* 19: 371-393.
127. **López S, Turon X, Montero E, Palacín C, Duarte CM, Tarjuelo I (1998).** Larval abundance, recruitment and early mortality in *Paracentrotus lividus* (Echinoidea). Interannual variability and plankton-benthos coupling. *Marine Ecology Progress Series* 172: 239-251.
128. **Lozano J, Galera J, Lopez S, Turon X, Palacin C, Morera G (1995)** Biological cycles and recruitment of *Paracentrotus lividus* (Echinodermata: Echinoidea) in two contrasting habitats. *Marine Ecology Progress Series* 122(1-3): 179-192.
129. **McClanahan TR y Muthiga NA (1989).** Patterns of predation on a sea urchin, *Echinometra mathaei* (De Blainville), on Kenyan coral reefs. *J. Exp. Mar. Biol. Ecol.* 126(1): 77-94.
130. **Metaxas A y Scheibling RE (1993).** Community structure and organization of tidepools. *Mar. Ecol. Prog. Ser.* 98(1-2): 187-198.
131. **Miguez Rodríguez LJ, Catoira JL, Gallego Castejón L (1990).** Nuevas aportaciones sobre el rendimiento gonadal y las relaciones biométricas de *Paracentrotus lividus* (Lamarck) en la ría de la Coruña. *Bentos VI.* : 275-281.
132. **Miller RJ, Adams AJ, Ebersole JP, Ruiz E (2007)** Evidence for positive density-dependent effects in recovering *Diadema antillarum* populations. *Journal of Experimental Marine Biology and Ecology* 349(2): 215-222.
133. **Montero-Torreiro MF y Garcia-Martinez P (2003).** Seasonal changes in the biochemical composition of body components of the sea urchin,

- Paracentrotus lividus*, in Lorbe (Galicia, north-western Spain). Journal of the Marine Biological Association of the UK 83(03): 575-581.
134. **Mottet MG (1976)**. The fishery biology of sea urchin in the family stronglylocentrotidae. Wash. Dept. Fisch. Tech. Rep. 20: 1-66.
135. **Munro JL (1982)**. Estimation of natural mortality rates from selectivity and catch length-frecuency data. ICLARM Fishbyte 1(1): 5-6.
136. **Nataf G (1954)**. Sur la croissance de *Paracentrotus lividus* et des *Psammechinus miliaris* (Gmelin) Bull. Mus. Hist. Nat. Paris, 2(26): 244-251.
137. **Naya I (2007)**. Meta-análisis estadístico de los patrones de crecimiento del erizo *Paracentrotus lividus* (Lamarck, 1816). DEA, Universidad de la Coruña.
138. **Nichols D (1981)**. The Cornish sea-urchin fishery. Cornish Studies 9: 5-16.
139. **Niell F y Pastor R (1973)**. Relaciones tróficas de *Paracentrotus lividus* (Lmk.) en la zona litoral. Invest.Pesq 37(1): 1-7.
140. **Ourens R (2007)**. Patrones geográficos, poblacionales y estacionales en los parámetros reproductivos del erizo *Paracentrotus lividus*. DEA, Universidad de la Coruña.
141. **País A, Chessa LA, Serra S, Ruiu A, Meloni G, Donno Y (2007)**. The impact of commercial and recreational harvesting for *Paracentrotus lividus* on shallow rocky reef sea urchin communities in north-western Sardinia, Italy. Estuarine, Coastal and Shelf Science 73(3-4): 589-597.

-
142. **Palacín, C., Giribet, G., Turon, X. (1997).** Patch recolonization through migration by the sea urchin *Paracentrotus lividus* (Lamarck) in communities with high algal cover and low sea urchins densities. *Cah. Biol. Mar.* 38: 267-271.
143. **Palacín C, Turon X, Ballesteros M, Giribet G, López S (1998) a.** Stock evaluation of three littoral echinoid species on the Catalan coast (North-Western Mediterranean). *Mar. Ecol.* 19(3): 163-177.
144. **Palacín C, Giribet G, Carner S, Dantart L, Turon X (1998) b.** Low densities of sea urchins influence the structure of algal assemblages in the Western Mediterranean. *J. Sea Res.* 39(3-4): 281-290.
145. **Pastor R (1971)** Distribución del erizo de mar, *Paracentrotus lividus* (Lmk), en la ría de Vigo. *Publ. Tecn. Dir. Gen. Pesca Marit, Spain* 9: 255-270.
146. **Pauly, D. y Gaschutz, G., (1979).** A simple method for fitting oscillating length growth data, with a program for pocket calculators. *ICES CM* 6: 24-26.
147. **Pearce CM, Daggett TL, Robinson SMC (2004).** Effect of urchin size and diet on gonad yield and quality in the green sea urchin (*Strongylocentrotus droebachiensis*). *Aquaculture* 233(1-4): 337-367.
148. **Pearse JS (1969).** Reproductive periodicities of indo-pacific invertebrates in the Gulf of Suez. II. The echinoid *Echinometra mathaei* (de Blainville). *Bull Mar Sci* 19(3): 580-612.
149. **Pearse JS (1981).** Synchronization of gametogenesis in the sea urchins *Strongylocentrotus purpuratus* and *S. franciscanus*. En: *Advanced in Invertebrate Reproduction*, pp 53-68.

- 150. Pearse JS y Cameron RA (1991)** Echinodermata: Echinoidea. *Reproduction of Marine Invertebrates* 6: 513–662.
- 151. Pedrotti ML y Fenaux L (1992).** Dispersal of echinoderm larvae in a geographical area marked by upwelling (Ligurian Sea, NW Mediterranean). *Marine Ecology Progress Series* 86: 217-217.
- 152. Pedrotti ML (1993).** Spatial and temporal distribution and recruitment of echinoderm larvae in the Ligurian Sea. *J Mar. Biol. Assoc. UK* 73(3): 515-530.
- 153. Preece MA y Baines MJ (1978).** A new family of mathematical models describing the human growth curve. *Ann. Hum. Biol.* 5: 1-24.
- 154. Régis MB (1978).** Croissance de deux échinoides du golfe de Marseille (*Paracentrotus lividus* (Lmk) et *Arbacia lixula* L.). *Aspects écologiques de la microstructure du squelette et de l' evolution des indices physiologiques*. Tesis doctoral, Universidad de Aix-Marseille III.
- 155. Régis MB (1979).** Analyse des fluctuations des indices physiologiques chez deux échinoides (*Paracentrotus lividus* (Lmk) et *Arbacia lixula* L.) du golfe de Marseille. *Tethys* 9(2): 167-181.
- 156. Richards FJ (1959)** A flexible growth function for empirical use. *J. Exp. Bot* 10(29): 290-300.
- 157. Ricker WE (1979).** Growth rates and models. *Fish physiology*. Academic Press, New York.

-
- 158. Roberts CM y Ormond RFG (1987).** Habitat complexity and coral reef fish diversity and abundance on red sea fringing reefs. *Marine Ecology Progress Series*. 41(1): 1-8.
- 159. Roden, C. M.; Hensey, M.; Ludgate R. (1987).** Aquaculture in Kilkieran and Bertraghbuoy Bays (Co. Galway); an atlas on environmental factors. *Aquacult. Tech. Bull N°12*. National Board for Science and Technology, Dublin.
- 160. Rowley RJ (1989).** Settlement and recruitment of sea urchins (*Strongylocentrotus* sp.) in a sea-urchin barren ground and a kelp bed: Are populations regulated by settlement or post-settlement processes? *Mar. Biol.* 100(4): 485-494.
- 161. Russel MP (1987).** Life history traits and resource allocation in the purple sea urchin *Strongylocentrotus purpuratus* (Stimpson). *J. Exp. Mar. Biol. Ecol.* 108: 199-216.
- 162. Russell MP y Meredith RW (2000).** Natural growth lines in echinoid ossicles are not reliable indicators of age: A test using *Strongylocentrotus droebachiensis*. *Invertebr Biol*: 410-420.
- 163. Sala E y Zabala M (1996).** Fish predation and the structure of the sea urchin *Paracentrotus lividus* populations in the NW Mediterranean. *Marine Ecology Progress Series* 140(1): 71-81.
- 164. Sala E (1997).** Fish predators and scavengers of the sea urchin *Paracentrotus lividus* in protected areas of the north-west Mediterranean Sea. *Mar. Biol.* 129(3): 531-539.

- 165. Sala E, Ribes M, Hereu B, Zabala M, Alvà V, Coma R, Garrabou J (1998).** Temporal variability in abundance of the sea urchins *Paracentrotus lividus* and *Arbacia lixula* in the Northwestern Mediterranean: Comparison between a marine reserve and an unprotected area. *Marine Ecology Progress Series* 168: 135-145.
- 166. Sánchez F y Olosó I (2004).** Effects of fisheries on the Cantabrian Sea shelf ecosystem. *Ecol. Model* 172(2-4): 151-174.
- 167. Sánchez-España AI, Martínez-Pita I, García FJ (2004).** Gonadal growth and reproduction in the commercial sea urchin *Paracentrotus lividus* (Lamarck, 1816) (Echinodermata: Echinoidea) from southern Spain. *Hydrobiologia* 519(1-3): 61-72.
- 168. Santos R y Flammang P (2007)** Intra and interspecific variation of attachment strength in sea urchins. *Marine Ecology Progress Series* 332: 129-142.
- 169. Savy, S. (1987).** Les prédateurs de *Paracentrotus lividus* (Echinodermata). In : Boudouresque, GIS Posidonie, IFREMER (eds.). *Colloque International sur Paracentrotus lividus et les oursins comestibles*. Marseille: 413-423.
- 170. Schnack-Schiel SB e Isla E (2005).** The role of zooplankton in the pelagic-benthic coupling of the southern ocean. *Scientia Marina* 69 (2). 39-55.
- 171. SCOR-UNESCO (1966).** Determination of photosynthetic pigments in sea-water. Report of working group 17. *Monographs on Oceanographic Methodology*, no. 1

-
- 172. Sellem F y Rezig M (1995).** Variabilité du cycle sexual de l'oursin *Paracentrotus lividus* Lamarck, 1816 (Echinodermata-Echinoidea) dans le golfe de Tunis (Méditerranée). *Oceanologica Acta* 23(5): 607-613.
- 173. Sellem F y Guillou M (2007).** Reproductive biology of *Paracentrotus lividus* (Echinodermata: Echinoidea) in two contrasting habitats of northern Tunisia (south-east Mediterranean). *Journal of the Marine Biological Association of the United Kingdom* 87(3): 763-767.
- 174. Sellem F, Langar H, Pesando D (2000).** Âge et croissance de l'oursin *Paracentrotus lividus* Lamarck, 1816 (Echinodermata-Echinoidea) dans le Golfe de Tunis (Méditerranée). *Oceanologica Acta* 23(5): 607-613.
- 175. Semroud R y Kada H (1987).** Contribution a l'etude de l'oursin *Paracentrotus lividus* (Lamarck) dans la region d'alger (Algerie): Indice de repletion et indice gonadique. *Colloque International Sur Paracentrotus Lividus et Les Oursins Comestibles*: 117-124.
- 176. Shears NT y Babcock RC (2002).** Marine reserves demonstrate top-down control of community structure on temperate reefs. *Oecologia* 132(1): 131-142.
- 177. Shirayama Y y Thornton H (2005).** Effect of increased atmospheric CO₂ on shallow water marine benthos. *J.Geophys.Res* 110.
- 178. Shpigel M, McBride SC, Marciano S, Lupatsch I (2004).** The effect of photoperiod and temperature on the reproduction of European sea urchin *Paracentrotus lividus*. *Aquaculture*, 232(1-4): 343-355.

- 179. Shpigel M, McBride SC, Marciano S, Ron S, Ben-Amotz A (2005).** Improving gonad colour and somatic index in the European sea urchin *Paracentrotus lividus*. *Aquaculture*, 245(1-4): 101-109.
- 180. Siikavuopio SI, Christiansen JS, Dale T (2006).** Effects of temperature and season on gonad growth and feed intake in the green sea urchin (*Strongylocentrotus droebachiensis*). *Aquaculture* 255(1-4): 389-394.
- 181. Sime AAT y Cranmer GJ (1985).** Age and growth of North Sea echinoids. *Journal of the Marine Biological Association of the United Kingdom*. 65(3): 583-588.
- 182. Sloan NA (1985).** Echinoderm fisheries of the world: A review. *Proceedings of the Fifth International Echinoderm conference*. Pp 109-124
- 183. Smith JR y Strehlow DR (1983)** Algal-induced spawning in the marine mussel *Mytilus californianus*. *Int.J.Invertebr.Reprod.* 6(2): 129-133.
- 184. Soriano M y Pauly D (1989).** A method for estimating the parameters of a seasonally oscillating growth curve from growth increment data. *International Center for Living Aquatic Resources Management, Fishbyte* 7: 18-21.
- 185. Spirlet C, Grosjean P, Jangoux M (2001)** Cultivation of *Paracentrotus lividus* (Echinodermata: Echinoidea) on extruded feeds: Digestive efficiency, somatic and gonadal growth. *Aquaculture Nutrition* 7(2): 91-99.
- 186. Spirlet C, Grosjean P, Jangoux M (2000).** Optimization of gonad growth by manipulation of temperature and photoperiod in cultivated sea urchins, *Paracentrotus lividus* (Lamarck) (Echinodermata). *Aquaculture* 185(1-2): 85-99.

-
- 187. Spirlet C, Grosjean P, Jangoux M (1998).** Reproductive cycle of the echinoid *Paracentrotus lividus*: Analysis by means of the maturity index. *Invertebrate Reproduction and Development* 34(1): 69-81.
- 188. Starr M, Himmelman JH, Therriault JC (1990).** Direct coupling of marine invertebrate spawning with phytoplankton blooms. *Science* 247(4946): 1071-1074.
- 189. Starr M, Himmelman JH, Therriault JC (1993).** Environmental control of green sea urchin, *Strongylocentrotus droebachiensis*, spawning in the St. Lawrence estuary. *Can J Fish Aquat Sci* 50(5): 894-901.
- 190. Tanaka M (1982).** A new growth curve which expresses infinite increase. *Publ Amakusa Mar. Biol. Lab* 6: 167-177.
- 191. Tanaka M (1988).** Eco-physiological meaning of parameters of ALOG growth curve. *Publ Amakusa Mar. Biol. Lab* 9: 103-106.
- 192. Tegner MJ y Dayton PK (1981).** Population structure, recruitment and mortality of two sea urchins (*Strongylocentrotus franciscanus* and *S. purpuratus*) in a kelp forest. *Mar Ecol.Prog. Ser.* 5: 255-268.
- 193. Tegner MJ y Levin LA (1983).** Spiny lobsters and sea urchins: Analysis of a predator-prey interaction. *J. Exp. Mar. Biol. Ecol.* 73(2): 125-150.
- 194. Thompson RJ (1982).** The relationship between food ration and reproductive effort in the green sea urchin *Strongylocentrotus droebachiensis*. *Oecologia* 56: 50-57.

195. **Tomas F, Romero J, Turon X (2004).** Settlement and recruitment of the sea urchin *Paracentrotus lividus* in two contrasting habitats in the Mediterranean. Marine Ecology Progress Series 282: 173-184.
196. **Tomas F, Álvarez-Cascos D, Turon X, Romero J (2006).** Differential element assimilation by sea urchins *Paracentrotus lividus* in seagrass beds: Implications for trophic interactions. Marine Ecology Progress Series 306: 125-131.
197. **Tomassone R, Dervin C, Masson JP (1993).** Biométrie: Modélisation de phénomènes biologiques. Masson, Paris.
198. **Tortonese E. (1965).** Echinodermata fauna d'Italia. Calderini,
199. **Turon X, Giribet G, Lopez S, Palacin C (1995).** Growth and population structure of *Paracentrotus lividus* (echinodermata: Echinoidea) in two contrasting habitats. Marine Ecology Progress Series 122(1-3): 193-204.
200. **Tuya F, Cisneros-Aguirre J, Ortega-Borges L, Haroun RJ (2007)** Bathymetric segregation of sea urchins on reefs of the Canarian archipelago: Role of flow-induced forces. Estuar. Coast Shelf Sci. 73(3-4): 481-488.
201. **Urgorri V, Reboreda P, Troncoso JS (1994).** Dispersión, demografía y producción gonadal de una población de *Paracentrotus lividus* (Lamarck, 1816). Memoria final de resultados.
202. **Vadas RL,Sr., Smith BD, Beal B, Dowling T (2002).** Sympatric growth morphs and size bimodality in the green sea urchin (*Strongylocentrotus droebachiensis*). Ecol.Monogr 72(1): 113-132.

203. **Vaughan, D.S. y Kanciruk, P. (1982).** An empirical comparison of estimation procedures for the Von Bertalanffy growth equation. *J Cons Int Explor Mer* 40: 211-219.
204. **Von Bertalanffy L (1938).** A quantitative theory of organic growth (inquiries of growth laws II). *Hum. Biol.* 10(2): 181-213.
205. **Von Bertalanffy L (1957).** Quantitative laws in metabolism and growth. *Quart Rev Biol* 32(3): 217-231.
206. **Vukovic A (1982).** Florofaunistic changes in the infralitoral zone after the sea urchin *Paracentrotus lividus* (L.) population explosion. *Acta Adriat* 23(1-2): 237-241.
207. **Walford LA (1946).** A new graphic method of describing the growth of animals. *Biol Bull* 90(2): 141-147.
208. **Walker C y Lesser M (1998).** Manipulation of food and photoperiod promotes out-of-season gametogenesis in the green sea urchin, *Strongylocentrotus droebachiensis*: Implications for aquaculture. *Mar. Biol.* 132(4): 663-676.
209. **Watanabe JM y Harrold C (1991).** Destructive grazing by sea urchins *Strongylocentrotus* sp. in a central California kelp forest: Potential roles of recruitment, depth, and predation. *Mar.Ecol.Prog. Ser.* 71(2): 125-141.
210. **Weibull W (1951).** A statistical distribution function of wide applicability. *J. Appl. Mech.* 18: 293-296.

211. **Williams CT y Harris LG (1998).** Growth of juvenile green sea urchins on natural and artificial diets. Proceedings of the Ninth International Echinoderm Conference.
212. **Yamaguchi M (1975).** Estimating growth parameters from growth rate data. *Oecologia*, 230: 321-332.
213. **Zamora S y Stotz W (1992)** Ciclo reproductivo de *Loxechinus albus* (Molina, 1782)(Echinodermata: Echinoidea) en punta lagunillas, IV región, Coquimbo, Chile. *Rev Chile Hist Nat* 65: 121-133.
214. **Zavodnik D (1987).** Synopsis on the sea urchin *Paracentrotus lividus* (Lamarck, 1816) in the Adriatic sea. Colloque International sur *Paracentrotus Lividus* et les Oursins Comestibles: 221-240.

WEBS

1. **Myers P, Espinosa R, Parr CS, Jones T, Hammond GS, Dewey TA (2008)** The animal diversity web.
2. **Smith AB (2004).** The echinoid directory. World Wide Web electronic publication.
3. www.fao.org
4. www.pescadegalicia.com
5. www.bim.ie
6. www.tematico.asturias.es/dgpesca/
7. www.teledeteccionoceanografica-ieo.net
8. www.fomento.es