

8.- BIBLIOGRAFIA

Agency for Toxic Substances and Disease Registry (ATSDR, 1992). Toxicological Profile for Vanadium. Departament of Health and Human Services, Atlanta.

Agency for Toxic Substances and Disease Registry (ATSDR, 1993). Toxicological Profile for Arsenic, Beryllium, Cadmium, Chromium, Lead, and Toluene. Departament of Health and Human Services, Atlanta

Agency for Toxic Substances and Disease Registry (ATSDR, 1994). Toxicological Profile for Pentachlorophenol. Departament of Health and Human Services, Atlanta

Agency for Toxic Substances and Disease Registry (ATSDR, 1996). Toxicological Profile for Nickel, Benzene, Hexaclorbenzene, Polychlorinated Biphenyls (PCBs) and Xylene. Departament of Health and Human Services, Atlanta.

Agency for Toxic Substances and Disease Registry (ATSDR, 1999). Toxicological profile for Mercury and Chlorinated Dibenzo-p-Dioxins (CDDs). Departament of Health and Human Services, Atlanta.

Agency for Toxic Substances and Disease Registry (ATSDR, 2000) Toxicological profile for manganese. Department of Health and Human Services, Public Health Service, Atlanta.

Agramunt MC, Domingo JL, Schuhmacher M, Gómez M, Llobet JM, Müller L (2002) Biological monitoring of organic substances in workers at a hazardous waste incinerator of Catalonia, Spain. Organohalogen Compd 55: 263-266.

Aguamarket (2002) Hexaclorbenceno. Diccionario del Agua.
(<http://aguamarket.com/diccionario/terminos>).

Alcock RE, Gemmill R, Jones KC (1999) Improvements to the UK PCDD/F and PCB atmospheric emission inventory following an emission measurement programme. Chemosphere 38: 759-770.

Alfvén T, Elinder CG, Carlsson MD, Grubb A, Hellström L, Persson B, Pettersson C, Spang G, Schütz, Jarup L (2000) Low-level cadmium exposure and osteoporosis. *J Bone Mineral Research* 15: 1579-1586.

Alimonti A, Petrucci F, Krachler M, Bocca B, Caroli S (2000) Reference values for chromium, nickel and vanadium in urine of youngsters from the urban area of Rome. *J Environ Monit* 2: 351-354.

Aller AJ (1990) The clinical significance of beryllium. *Trace Elem Electrolytes Health Dis* 4: 1-6.

American Conference of Goverment Industrial Hygienists (ACGIH, Worldwide). Guide to occupational Exposure Values, 1999.

AMICLOR (1997) Dioxinas. Organización de Usuarios y Trabajadores de la Química del Cloro. (<http://www.amiclor.org/opciones/dioxinas.shtml>)

Angerer J, Heinzel B, Reimann DO, Knorz W, Lehnert G (1992) Internal exposure to organic substances in a municipal waste incinerator. *Int Arch Occup Environ Health* 64: 265-273.

Aoki Y (2001) Polychlorinated biphenyls, polychlorinated dibenzo-p-dioxins, and polychlorinated dibenzofurans as endocrine disrupters- What we have learned from Yusho disease. *Environ Res* 86: 2-11.

Apostoli P, Lucchini R, Alessio L (2000) Are current biomarkers suitable for the assessment of manganese exposure in individual workers? *American J Ind Med* 37: 283-290.

Apostoli P, Maranelli G, Duca PG, Bavazzano P, Bortoli A, Cruciani A, Elia G, Minoia C, Piccinini R, Sabbioni E, Sciarra G, Soave C (1997) Reference values of urinary chromium in Italy. *Int Arch Occup Environ Health* 70: 173-179.

Arfi C, Seta N, Fraisse D, Revel A, Escande JP, Momas I (2001) Dioxins in adipose tissue of non-occupationally exposed persons in France: correlation with individual food exposure. *Chemosphere* 44: 1347-1352.

Aschner M, Aschner JL (1990) Mercury neurotoxicity: mechanisms of blood-brain barrier transport. *Neurosci Biobehav Rev* 14: 169-176.

Astor S (1979) Über die unwelttoxikologische Bedeutung des pentachlorphenols. Kiel. Schriftenreihe der Untersuchungsstelle für Umwelttoxikologie des Landes Schleswig-Holstein. Núm. 6.

Avkhimenko M (2000) Medical and ecological consequences of environmental contamination with PCBs, in: Polychlorinated Biphenyls-Supertoxicants of XXI Century, VINYITY Pres, Moscow: 14-30.

Avnstorp C (1992) Cement eczema. An epidemiological intervention study. *Acta Derm Venereol Suppl Stockh* 179: 1-22.

Baldwin DR, Marshall W (1999) Heavy metal poisoning and its laboratory investigation. *Ann Clin Biochem* 36: 267-300.

Ballschmiter K, Braunmiller I, Niemcyk R, Swerev M (1988) Reaction pathways for the formation of polychlorodibenzodioxins (PCDD) and -dibenzofurans (PCDF) in combustion processes. II. Chlorobenzenes and chlorophenols as precursors in the formation of polychloro-dibenzodioxins and -dibenzofurans in flame chemistry. *Chemosphere* 17: 995-1005.

Barman SC, Bhargava SK (1997) Accumulation of heavy metals in soil and plants in industrially polluted fields. *Ecological Environ Imp Ass* : 289-314.

Barregard L, Sallsten G, Jarvholm B (1990) Mortality and cancer incidence in chloralkali workers exposed to inorganic mercury. *Br J Ind Med* 47: 99-104.

Bartell SM, Ponce RA, Sanga RN, Faustman EM (2000) Human variability in mercury toxicokinetics and steady state biomarker ratios. Environ Res 84: 127-132.

Baughman R, Meselson M (1973) Environ Health Perspect 5: 27-35. En: Kumar KS, Kannan K, Paramasivar O i cols (2001) Polychlorinated biphenyls in human tissues, meat, fish, and wildlife samples from India. Environ Sci Technol 35: 3448-3455.

Benes B, Spevackova V, Smid J, Cejchanova M, Cerna M, Subrt P, Marecek J (2000) The concentration levels of Cd, Pb, Hg, Cu, Zn and Se in blood of the population in the Czech Republic. Cent Eur J Public Health 8: 117-119.

Birnbaum LS (1994) The mechanism of dioxin toxicity: The relationship to risk assessment. Environ Health Perspect 102: 157-167.

Blanusa M(1996) Assessment of total human exposure to lead and cadmium. Environ Manag Health 7: 23-28.

Bordado J, Ferreira H, Gomes J (1999) Dioxinas e dibenzofuranos no meio ambiente. Departamento de Engenharia Química. Instituto Superior Técnico. Av. Rovisco Pais, 1906 Lisboa Codex.

Börjesson J, Gerhardsson L, Schütz A, Perfekt R, Mattson S, Skerfvin S (2001) Kidney cadmium as compared to other markers of cadmium exposure in workers at a secondary metal smelter. Am J Ind Med 39: 19-28.

BUA-Beratergremium für Umweltrelevante Altstoffe der Gesellschaft Deutscher Chemiker (1988) Benzol. Basilea, Heidelberg & Zurich Nº 24.

Burse VW, Groce DF, Caudill SP, Korver MP, Philips DL, McClure PC, Lapeza CR, Head SL, Miller DT, Buckley DJ (1994) Determination of polychlorinated biphenyl levels in the serum of residents and in the homogenates of seafood from the New Bedford, Massachusetts, area: a comparison of exposure sources through pattern recognition techniques. Sci Total Environ 144: 153-177.

- Cano-Pavon JM, García de Torres A, Sánchez-Rojas F, Cañada-Rudner P (1999) Analytical methods for mercury speciation in environmental and biological samples-an overview. *Int J Environ Anal Chem* 75: 93-106.
- Chang MB, Wu HS, Huang CK (2000) Evaluation on speciation and removal efficiencies of mercury from municipal solid waste incinerators in Taiwan. *Sci Total Environ* 246: 165-173.
- Chevrier J, Dewailly E, Ayotte P, Mauriege P, Despres JP, Treblay A (2000) Body weight loss increases plasma and adipose tissue concentrations of potentially toxic pollutants in obese individuals. *Int J Obes Metab Disord* 24: 1272-1278.
- Chiarenazelli JR, Alexabder C, Isley A, Serudato R, Pagano J, Ramirez W (2001) Polychlorinated biphenyls in no accumulating, century-old sediments: sources, signatures, and mechanism of introduction. *Environ Sci Technol* 35: 2903-2908.
- Christensen JM, Kristiansen J, Nielsen NH, Menne T, Byralsen J (1999) Nickel concentrations in serum and urine of patients with nickel eczema. *Toxicol Lett* 108: 185-189.
- Clarke AN, Megehee MM, Lowe DL, Clarke JH (1994) A review of polychlorinated dibenzofurans and polychlorinated dibenzodioxines in sediments in the United States and international waterways. *Hazard Waste Hazard Materials* 11: 253-275.
- Clayton CA, Pellizari ED, Whitmore RW, Perrit RL, Quackenboss JJ (1999) National human exposure assessment survey (NHEXAS): distribution and associations of lead, arsenic and volatile organic compounds in EPA Region 5. *J Exp Anal Environ Epidemiol* 9: 381-392.
- Cledes J, Allain P (1992) Nephropatie saturnine chronique. Epidemiologie et dianostic. *Presse Med* 21: 759-762.
- Cole DC, Kearney J, Ryan JJ, Gilman AP (1997) Plasma levels and profiles of dioxin and dioxin-like compounds in Ontario Great lakes anglers. *Chemosphere* 34: 1401-1409.

Coleman PJ, Lee RGM, Alcock RE, Jones KC (1997) Observations on PAH, PCB, and PCD/F trends in UK urban air, 1991-1995. Environ Sci Technol 31: 2120-2124.

Collins JF, Brown JP, Painter PR, Jamall IS, Zeise LA, Alexeeff GV, Wade MJ, Siegel DM, Wong JJ (1992) On the carcinogenicity of cadmium by the oral route. Regul Toxicol Pharmacol 16: 57-72.

Concha G, Vogler G, Lezcano D (1998) Exposure to inorganic arsenic metabolites during early human development. Toxicol Sci 44: 185-190.

Consejero General de Colegios Oficiales de Farmacéuticos (2002) Informe Dioxinas (www.cof.es/consejo/dioxina.htm).

Corbella J (2000) Esquemes de toxicologia industrial (I). Introducció-Metalls. Publicacions del Seminari Pere Mata de la Universitat de Barcelona, núm. 24.

Cornelis R, Heinzel B, Herber RFM, Christensen JM, Poulsen OM, Sabbioni E, Templeton DM, Tomesen Y, Vahter M, Vesterberg O (1996) Simple collection guidelines for trace elements in blood and urine. J Trace Elem Med Biol 10: 103-127.

Cornelis R, Sabbioni E, Van der Venne MT (1994) Trace element reference values in tissues from inhabitants of the European Community. VII. Review of trace elements in blood, serum, and urine of the Belgian population and critical evaluation of their possible use as reference values. Sci Total Environ 158: 191-226.

Counter SA, Buchanan LH, Ortega F, Laurell G (2002) Elevated blood mercury and neuro-otological observations in children of the Ecuadorian gold mines. J Toxicol Environ Health 65: 149-163.

Crine JP (1988) Hazards, Decontamination and Replacement of PCB. A Comprehensive Guide. Crine (ed) New York & London.

Daniel V, Huber W, Bauer K, Suesal C, Conradt C, Opelz G (2001) Associations of blood levels of PCBs, HCHs, and HCB with numbers of lymphocyte subpopulations, in vitro lymphocyte response, plasma cytokine levels, and immunoglobulin autoantibodies. Environ Health Persp 109: 173-178.

Deohdar SD, Barna BP (1991) Immune mechanisms in beryllium lung disease. Cleveland Clin J Med 58: 157-160.

De Rosa Ch T, Pohl HR, Willians M, Ademoyer AA, Selene CH, Chou J, Jones DE (1998) Public health implications of environmental exposures. Environ Health Perspect 106: 369-378.

De Vito MJ, Birnbaum LS (1994) Toxicology of dioxins and related chemicals. A: Dioxins and Health. Schecter (ed), Plenum Press, NY.

Deml E, Mangelsdorf I, Greim H (1996) Chlorinated dibenzodioxins and dibenzofurans (PCDD/F) in blood and human milk of non occupationally exposed persons living in the vicinity of a municipal waste incinerator. Chemosphere 33: 1941-1950.

Dempsey CR, Oppelt ET (1993) Incineration of hazardous waste: a critical review update. Air Waste 43: 25-73.

DFG Senatskommission zur Prüfung gesundheitsschädlicher Arbeitsstoffe (eds) (1995) MAK – und BAT-Werte-Liste. VCH, Weinheim.

DOCE. Diari Oficial de les Comunitats Europees (1994) Posició Comú N° 26/94 relativa a la incineració de residus perillósos, N°C232: 35-50.

Domingo JL, Gómez M, Sánchez DL (1995) Toxicology of vanadium compounds in diabetic rats: the action of chelating agents on vanadium accumulation. Mol Cell Biochem 153: 233-240.

Domingo JL, Schuhmacher M, Granero S, Llobet JM (1999) PCDDs and PCDFs in food samples from Catalonia, Spain. An assessment of dietary intake. Chemosphere 38: 3517-3528.

Domingo JL, Schuhmacher M, Agramunt MC, Müller L, Neugebauer F (2001) Levels of metals and organic substances in blood and urine of workers at a new hazardous waste incinerator. Int Arch Occup Environ Health 74: 263-269.

Droz PO (1989) Biological Monitoring I: Sources of variability in human response to chemical exposure. Appl Ind Hyg 4: F20-G24.

Ducroffe G, Claeys F, Bruaux P (1990) Lowering time trend of blood lead levels in Belgium since 1978. Environ Res 51: 25-34.

Dunstan RH, Donohoe M, Taylor W, Roberts TK, Murdoch RN, Watkins JA, McGregor NR (1995) A preliminary investigation of chlorinated hydrocarbons abd chronic fatigue syndrome. Med J Aust 163: 294-297.

Ebinghaus R, Jennings SG, Schroeder WH, Berg T (1999) International field intercomparison measurements of atmospheric mercury species at Mace Head, Ireland. Atmos Environ 33: 3063-3073.

Eduljee G (1999) Secondary exposure to dioxins through exposure to PCP and its derivates. Sci Total Environ 232: 193-214.

Elghany NA, Schumacher MC, Slattery ML, West DW, Lee JS (1990) Occupation, cadmium exposure, and prostate cancer. Epidemiology 1: 107-115.

Eljarrat E, Monjonell A, Caixach J, Rivera J (2002) Toxic potency of polychlorinated dibenzo-p-dioxins, polychlorinated dibenzofurans, and polychlorinated biphenyls in food samples from Catalonia (Spain). J Agr Food Chem 50: 1161-1167.

EPA (1997) Mercury study report to congress. US Environmental Protection Agency, Washington, DC.

European Centre for Ecotoxicology and Toxicology of Chemicals (1996) Risk assessment for carcinogens. Brussels. Monograph 24: 45.

Fabiánová E, Hettychová L, Koppová K i cols (2000) Health risk assessment for inhalation exposure to arsenic. Cent Eur J Publ Health 8: 28-32.

Flesch-Janys D, Becher H, Gurn P, Jung D, Konietzko J, Manz A, Päpke O (1996) Elimination of polychlorinated dibenzo-p-dioxins and dibenzofurans in occupationally exposed persons. J Toxicol Environ Health 47: 363-378.

Fiedler H (1998) Thermal formation of PCDD/PCDF: a survey. Environ Eng Sci 15: 49-58.

Fiedler H (1999) Dioxin and furan inventories. National and Regional Emissions of PCDD/PCDF. UNED chemicals, Geneva, Switzerland.

Fiedler H, Hutzinger O (1990) Dioxins: sources of environmental load and human exposure. Toxicol Environ Chem 29: 157-234.

Finkelstein Y, Markowitz ME, Rosen JF (1998) Low-lead lead-induced neurotoxicity in children: an update on central nervous system effects. Brain Res Rev 27: 168-176.

Fitzgerald DF, Clarkson TW (1991) Mercury and monomethylmercury: present and future concerns. Environ Health Perspect 96: 159-166.

Fleming CR (1988) Trace element metabolism in adult patients requiring total parenteral nutrition. Am J Clin Nutr 49: 573-579.

Fries GF (1995) Transport to organic environmental contaminants to animal products. Rev Environ Contam Toxicol 141: 71-109.

Furman A, Laleli M (1999) Analysis of lead body burden in Turkey. Sci Total Environ 234: 37-42.

Gabrio T, Piechotowski I, Wallenhorst T, Klett M, Cott L, Friebel P, Link B, Schwenk M (2000) PCB-blood levels in teachers, working in PCB-contaminated schools. Chemosphere 40: 1055-1062.

García Ariño C (1996) Síntesis diagnóstica de las enfermedades profesionales producidas por metales, productos químicos y agentes vivos. Ed. MAPFRE.

Gerhardsson L, Nordberg GF (1993) Lung cancer in smelter workers- interactions of metals as indicated by tissue levels. Scand J Work Environ Health 19: 90-94.

Gilman H, Dietrich J (1957) J Amer Chem Soc 79: 1439. En: Soriano V. Dioxinas (<http://www.opolanco.es/Apat/Boletín> 14/dioxinas.htm)

Gleadle A, Gem M, Harrision N (2000) Monitoring dietary exposure to dioxins and PCBs in the UK. Organohalogen Compd 47: 292-295.

Glynn AW, Wolk A, Aune M, Atuma S, Zettermark S, Maehle-Schimdt M, Darnerud PO, Becker W, Vessby B, Adami HO (2000) Serum concentrations of organochlorines in men: a search for markers of exposure. Sci Total Environ 263: 197-208.

Göen Th, Gündel J, Schaller KH, Angerer J (1995) The elimination of 1-hydroxypyrene in the urine of the general population and workers with different occupational exposures to PAH. Sci Total Environ 163: 195-201.

González CA, Kogevinas M, Huici A, Gadea E, Ladona M, Bosch A, Bleda MJ (1998) Blood levels of polychlorinated dibenzodioxins, polychlorinated dibenzofurans and polychlorinated biphenyls in the general population of a Spanish Mediterranean city. Chemosphere 3: 419-426.

Granero S, Llobet JM, Schuhmacher M, Corbella J, Domingo JL (1998) Biological monitoring of environmental pollution and human exposure to metals in Tarragona, Spain. I. Levels in hair of school children. Trace Elem Electrol 15: 39-43.

Haberman AL, Pratt M, Storrs FJ (1993) Contact dermatitis from beryllium in dental alloys. *Contact Dermatitis* 28: 157-162.

Hamilton E, Sabbioni E, Van der Venne MT (1994) Trace element reference values in tissues from inhabitants of the European Community. VI. Review of elements in blood, plasma, and urine, and a critical evaluation of reference values for the United Kingdom population. *Sci Total Environ* 158: 165-190.

Hansen AM, Christensen JM, Sherson D (1995) Estimation of reference values for urinary 1-hydroxypyrene and α -naphthol in Danish workers. *Sci Total Environ* 163: 211-219.

Hansson M, Barregard L, Sällsten G, Svensson BG, Rappe C (1997) Polychlorinated dibenzo-p-dioxin and dibenzofuran levels and patterns in polyvinylchloride and chloroalcaly industry workers. *Int Arch Occup Environ Health* 70: 51-56.

Hansson M, Grimstad T, Rappe C (1995) Occupational exposure to polychlorinated dibenzo-p-dioxins and dibenzofurans in a magnesium production plant. *Occup Environ Med* 52: 823-826.

Harada M, Nakanishi J, Yasoda E, Pinheiro -mcn, Oikawa T, Guimaraes GA, Cardoso BS, Kizaki T, Ohno H (2001) Mercury pollution in the Tapajos River basin, Amazon mercury level of head hair and health effects. *Environ Inter* 27: 285-290.

Harrad SJ, Jones KC (1992) A source inventory and budget for chlorinated dioxins and furans in the United Kingdom environment. *Sci Total Environ* 126: 89-107.

Harvey RG (1991) Polycyclic aromatic hydrocarbons chemistry and carcinogeniticy. New York: Cambridge Press.

Hasselriis F, Licata A (1996) Analysis of heavy metal emission data from municipal waste combustion. *J Hazard Mater* 47: 77-102.

Hays SM, Aylward LL (2001) Temporal trends in body-burden suggest that dioxin exposures in the general population have declined significantly. *Organic Compounds Human Exposure*.

Heederik D, Hooiveld M, Bueno-de-Mesquita HB (1998) Modelling of 2,3,7,8-tetrachlorodibenzo-*p*-dioxins levels in a cohort of workers with exposure to phenoxy herbicides and chlorophenols. *Chemosphere* 37: 1743-1754.

Heiselman DE, Cannon LA (1990) Benzene and the aromatic hydrocarbons. En: Adddad LM, Winchester JF, eds. *Clinical management of poisoning and drugs overdose*. Filadelfia: W.B. Saunders 1990; 1222-1230.

Herce-Pagliai C, González G, Camean AM, Repetto M (1999) Presence and distribution of arsenical species in beers. *Food Addit Contam* 16: 267-271.

Hernandez M, Schuhmacher M, Fernandez JD, Domingo JL, Llobet JM (1996) Urinary cadmium levels during pregnancy and postpartum. A longitudinal study. *Biol Trace Elem Res* 53: 205-212.

Hill RH, Head SL, Baker S, Gregg M, Shealy DB, Bailey SL, Williams CC, Sampson EJ, Needham LL (1995) Pesticide residues in urine of adults living in the United States: reference range concentrations. *Environ Res* 71: 99-108.

Hoffmann K, Becker K, Friedrich C, Helm D, Krause C, Seifert B (2000) The German Environmental Survey 1990/1992 (GerEs II): cadmium in blood, urine and hair of adults and children. *J Exp An Environ Epidemiol* 10: 126-135.

Horng CJ, Tsai JL, Lin SR (1999) Determination of urinary arsenic, mercury, and selenium in steel production workers. *Biol Trace Elem Res* 70: 29-40.

Houston DK, Johnson MA (1999) Lead as a risk factor for hypertension in women. *Nutr Rev* 57: 277-279.

Hu H (2000) Exposure to metals. *Occup Environ Med* 4: 983-996.

Hu SW, Shy CM (2001) Health effects of waste incineration: a review of epidemiologic studies. *J Air Waste Manage Assoc* 51: 1100-1209.

Huang CC, Lu CS, Chu NS, Hochberg F, Lilienfield D, Olanow W, Calne DB (1993) Progression after chronic manganese exposure. *Neurology* 43: 1479-1483.

IARC (1983) Polycyclic aromatic compounds: Part 1. Chemical, environmental data. In: IARC monographs on the evaluation of carcinogenic risks of chemicals to humans. Vol. 32.

IARC (1984) Monographs on the evaluation of carcinogenic risks of chemicals to humans. Polynuclear aromatic compounds. Vol.4, part 3. Lyon, France.

IARC (1990) Monographs on evaluation of carcinogenic risks to humans: chromium, nickel and welding. The Internation Agency for Cancer Research. Lyon, France. Vol. 49.

ICDA (1997) Criteria Document for hexavalent chromium. Internal Chromium Development Association. Paris, France.

Imbeault P, Tremblay A, Simoneau JA, Joannis DR (2002) Weight loss-induced rise in plasma pollutant is associated with reduced skeletal muscle oxidative capacity. *Am J Physiol Endocrinol Metab* 282: E574-579.

Iwata K, Saito H, Moriyama M, Nakano A (1992) Follow up study of renal tubular dysfunction and mortality in residents of an area polluted with cadmium. *Br J Ind Med* 49: 736-737.

Jakob A, Stucki S, Kuhn P (1995) Evaporation of heavy metals during the heat treatment of municipal solid waste fly ash. *Environ Sci Technol* 29: 2429-2436.

Janssen CR, Schamphelaere KD, Heijerick D, Muyssen B, Lock K, Bossuyt B, Vangheluwe M, Van Sprang P (2000) Uncertainties in the environmental risk assessment of metals. *Human Ecol Risk Assess* 6: 1003-1018.

Jarup L, Hellström L, Alfvén T, Carisson MD, Grubb A, Persson B, Pettersson C, Spang G, Schütz A, Elinder CG (2000) Low level exposure to cadmium and early kidney damage: the OSCAR study. *Occup Environ Med* 10: 668-672

Jiménez B, Hernández LM, Eljarrat E, Rivera J, González MJ (1996) Levels of PCDDs, PCDFs and non-ortho PCBs in serum samples of non-exposed individuals in Madrid (Spain). *Chemosphere* 33: 2403-2410.

Jones TD (1995) Toxicological potency of 2,3,7,8-tetrachlorodibenzo-p-dioxin relative to 100 other compounds: a relative potency analysis of in vitro and in vivo test data. *Arch Environ Contam Toxicol* 29: 77-85.

Jongeneelen FJ (2001) Benchmark guideline for urinary 1-hydroxypyrene as biomarker of occupational exposure to polycyclic aromatic hydrocarbons. *Ann Occup Hyg* 45: 3-13.

Kahan E, Derazne E, Rosenboim J, Ashkenazi R, Ribak J (1992) Adverse health effects in workers exposed to cadmium. *Am J Ind Med* 21: 527-537.

Kaise T, Watanabe S, Itoh K (1985) The acute toxicity of arsenobetaine. *Chemosphere* 14: 1327-1332.

Kalen SN, Goldman RH (2002) Mercury exposure: current concepts, controversies, and a clinic's experience. *J Occup Environ Med* 44: 143-154.

Kano K, Horikawa M, Utsunomiya T, Tati M, Satoh K, Yamaguchi S (1993) Lung cancer mortality among a cohort of male chromate pigment workers in Japan. *Int J Epidemiol* 22: 16-22.

Karimi A, Moniri F, Nasihatkon A, Zarepoor MJ, Alborzi A (2002) Mercury exposure among residents of a building block in Shirz, Iran. *Environ Research Section A* 88: 41-43.

Kasuba V, Rozgaj R, Sentija K (2000) Cytogenetic changes in subjects occupationally exposed to benzene. Chemosphere 40: 307-310.

Kawada T, Tohyama C, Suzuki S (1990) Significance of the excretion of urinary indicator proteins for a low level of occupational exposure to cadmium. Int Arch Occup Environ Health 62: 95-100.

Kent H (1999) Vanadium for diabetes. CMAJ 160: 17.

King R (1988) Mercury poisoning and its control. (ed) United Trade Press Limited, Londres.

Kitamura K, Kikichi Y, Watanabe S, Waechter i cols (2000) Health effects of chronic exposure to polychlorinated dibenzo-p-dioxins (PCDD), dibenzofurans (PCDF) and coplanar PCB (Co-PCB) of municipal waste incinerator workers. J Epidemiol 10: 262-270.

Komaromy-Hiller G, Ashm KO, Costa R, Howerton K (2000) Comparison of representative ranges based on US patient population and literature reference intervals for urinary trace elements. Clin Chim Acta 296: 71-90.

Korrick SA, Hunter DJ, Rotnizky A, Hu H, Speizer FE (1999) Lead and hypertension in a sample of middle-aged women. Am J Public Health 89: 330-335.

Krachler M, Habersack-Walner S, Eber B i cols. (1996) Influence of a low-fat diet and activity program on the trace element status of patients with coronary heart disease. Metal Ions in Biology and Medicine. Ed. Ph Collery 4: 560-562.

Kreiss K, Newman LS, Mroz M (1991) Blood testing for chronic beryllium disease. J Occup Med 33: 1188-1189.

Kristiansen J, Christensen JM, Iversen BS, Sabbioni E (1997) Toxic trace element reference levels in blood and urine: influence of gender and lifestyle factors. Sci Total Environ 204: 147-160.

Kurkjian R (2000) Metal contamination in the Republic of Armenia. Environ Manag 25: 477-483.

Kuschner M (1981) The carcinogenicity of beryllium. Environ Health Perspect 40: 101-105.

Kutz FW, Barnes DG, Bottimore DP (1990) The international toxicity equivalency factor (I-TEF) method of risk assessment for complex mixtures of dioxins and related compounds. Chemosphere 20: 751-757.

La Dou J (1986) Health issues in the microelectronics. State of the Art Reviews: Occupational Medicine 1: 1-11.

Langard S (1990) One hundred years of chomium and cancer: a review of epidemiological evidence and selected case reports. Am J Ind Med 17: 189-215.

Langworth S, Almkvist O, Wikstrom BO (1992) Effects of occupational exposure to mercury vapour on the central nervous system. Br J Ind Med 49: 545-555.

Lehnert G, Angerer J, Göen T, Schaller KH (1994) Referenzwerte für persistente Gefahrstoffe am Beispiel der Konzentration an polychorierten biphenylen in humanserum. Arbeitsmed Sozialmed Umweltmed 29: 454-458.

Levy L, Shuker LK, Rowbotham AL (2001) An evaluation of total personal exposure to chromium in the UK environment and associated possible adverse health effects. Environ Geo Health 23: 181-186.

Li H, Krieger RI, Li QX (2000) Improved HPLC method for analysis of 1 hydroxypyrene in human urine specimens of cigarette smokers. Sci Total Environ 257: 147-153.

Liem AKD, Furst P, Rappe C (2000) Exposure of populations to dioxins and related compounds. Food Addit Contam 17: 241-259.

Lin TW, Huang SD (2001) Direct and simultaneous determination of copper, chromium, and manganese in urine with a multielement graphite furnace atomic absorption spectrometer. *Anal Chem* 73: 4319-4325.

Lisk DL (1988) Environmental implications of incineration of municipal solid waste and disposal. *Sci Total Environ* 74: 39-66.

López-Artigues M, Soria ML, Cameán A, Repetto M (1993) Cadmium in the diet of the local population of Seville (Spain). *Bull Environ Contam Toxicol* 50: 417-424.

Lorber M, Cleverly D, Schaum J (1994) Development and validation of an air-to-beef food chain model for dioxine-like compounds. *Sci Total Environ* 156: 39-65.

Llobet JM, Granero S, Schuhmacher M, Corbella J, Domingo JL (1998a) Biological monitoring of environmental pollution and human exposure to metals in Tarragona, Spain IV. Estimation of the dietary intake. *Trace Elem Elec* 15: 136-141.

Llobet JM, Granero S, Torres A, Schuhmacher M, Domingo JL (1998b) Biological monitoring of environmental pollution and human exposure to metals in Tarragona, Spain III. Blood levels. *Trace Elem Electr* 15: 76-80.

Lustenhouwer JWA, Olie K, Hutzinger O (1980) Chlorinated dibenzo-*p*-dioxins and related compounds in incinerator effluents. *Chemosphere* 9: 501-522.

Malisch R, Bruns-Weller E, Knoll A, Fürst P, Mayer R, Wiesmüller T (2000) Results of an "emergency quality control study" as confirmation of a PCDD/PCDF-contamination of milk and butter samples. *Chemosphere* 40: 1033-1040.

Mastrantonio G (2000) Compuestos organoclorados como contaminantes persistentes: el caso de las dioxinas y los bifenilos policlorados. Progeso SA - Grupo Consultores Independiente. Nota técnica nº 2: 1-2.

McLauchlan GA (1991) Acute mercury poisoning. *Anaesthesia* 46: 110-112.

McLachlan MS, Thomas H, Reissinger M, Hutzinger O (1990) PCDD/F in an agricultural food chain. Part 1: PCDD/F mass balance of a lactating cow. Chemosphere 20: 1013-1020.

Meharg AA, Osborn D (1995) Dioxins released from chemical accidents. Nature 375: 353-354.

Mercado Calderón F (1993) Evaluación del 1-hidroxipireno como bioindicador de la exposición industrial a los hidrocarburos aromáticos policíclicos. Salud Pública de México 35: 6.

Mergler D, Baldwin M, Bélanger S, Larribe F, Beuter A, Bowler R, Panisset M, Edwards R, De Geoffroy A, Sassine MP, Hudnell K (1999) Manganese neurotoxicity, a continuum of dysfunction: results from a community based study. Neurotoxicology 20: 327-342.

Messerer P, Zober A, Becher H (1998) Blood lipid concentrations of dioxins and furans in a sample of BASF employees included in the IARC registry of workers exposed to phenoxy acid herbicides and/or chlorophenols. Environ Health Persp 106: 733-735.

Narasimhan TR, Kim HL, Safe SH (1991) Effects of hydroxylated polychlorinated biphenyls on mouse liver mitochondrial oxidate phosphorylation. J Biochem Toxicol 6: 229-236.

Nogawa K, Ishizaki A, Kobayashi E (1979) A comparison between health effects of cadmium and cadmium concentration in urine among inhabitants of the Itai-Itai disease endemic district. Environmental Research 18: 397-409.

Nriagu JO (1988) A silent epidemic of environmental metal poisoning?. Environ Pollut 50: 139-161.

Nriagu JO, Pacyna JM (1988) Quantitative assessment on worldwide contamination of air, water and soils by trace metals. Nature 333: 134-139.

NRC (2001) Arsenic in drinking water. Subcommittee on arsenic in drinking water, National Research Council. National Academy of Sciences. National Academy Press, Washington DC.

OMS (1977) IARC Monographs on the evaluation of the cancerogenic risk of chemicals to man. Some fumigants. The Herbicides 2,4-d and 2,4,5-t. Chlorinated Dibenzodioxins and miscellaneous industrial chemicals. IARC núm. 15. Lyon.

Orlinskii D, Pripitina I, Popova A, Shalanda A, Tsongas T, Hinman G, Butcher W (2001) Influence of environmental contamination with PCBs on human health. Environ Geoch Health 23: 317-332.

Outridge PM, Scheuhammer AM (1993) Bioaccumulation and toxicology of chromium: implications for wild life. Rev Environ Contam Toxicol 130: 31-77.

Pandey PK, Patel KS, Lenicek J (1999) Polycyclic aromatic hydrocarbons: need for assessment of health risks in India? Environ Mon Ass 59: 287-319.

Päpke O, Ball M, Lis A (1993) Potencial occupational exposure of municipal waste incinerator workers with PCDD/PCDF. Chemosphere 27: 203-209.

Päpke O, Ball M, Lis A (1994) PCDD/PCDF in humans. An update of background data. Chemosphere 29: 2355-2360.

Päpke O, Ball M, Lis A, Wuthe J (1996) PCDD/PCDF in humans. Follow-up of background data for Germany, 1994. Chemosphere 32: 575-582.

Paschal DC, Burt V, Caudill SP, Gunter EW, Pirkle JL, Sampson EJ, Miller DT, Jackson RJ (2000) Exposure of US population age 6 years and older to cadmium: 1988-1994. Arch Environ Contam Toxicol 38: 377-383.

Patterson DJJ, Fingerhut MA, Roberts DW, Needham LL, Haring C, Sweeney M, Marlow DA, Andrews JS, Halperin WE (1989) Levels of polychlorinated dibenzo-p-dioxins (PCDDs) and dibenzofurans (PCDFs) in workers exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin. Am J Ind Med 16: 135-146.

Paustendbach D, Connor K (2000) International concern over persistent organic pollutans (POPs) in foods: risk assessment and public policy. Organohalogen Compd 47: 284-287.

Payton M, Riggs KM, Spiro A, Weiss ST, Hu H (1998) Relations of bone and blood lead to cognitive function: The VA Normative Aging Study. Neurot Terat 20 (1) : 19-27.

Pilliére F, Roussenli X (1992) La surveillance biologique. De la théorie à la pratique. Cahier de notes documentaires 506 : 626-630.

Pirkle JL, Brody DJ, Gunter EW, Kramer RA, Paschal DC, Flegal KM, Matte TD (1994) The decline in blood lead levels in the United States. The National Health and Nutrition Examination Surveys (NHANES). JAMA 272: 284-291.

Pohl H, Holler J (1995) Halogenated aromatic hydrocarbons and toxicoty equivalency factors (TEFs) from the public health assessment perspective. Chemosphere 31: 2547-2559.

Rappe C (1992) Sources of PCDDs and PCDFs. Introduction, reactions, levels, patterns, profiles and trends. Chemosphere 25: 41-44.

Robards K, Worsford P (1991) Cadmium: toxicology and analysis. A review. Analyst 116: 549-568.

Rodushkin I, Ödman F, Appelblad PK (1999) Multielement determination and lead isotope ratio measurement in alcoholic beverages by high-resolution inductively couple plasma spectrometry. J Food Comp Anal 12: 243-257.

Rosenberg C, Kontsas H, Tornaeus J, Mutanen P, Vainio H, Patterson DG, Needham LL (1995) PCDD/PCDF levels in the blood of workers at a pulp and paper mill. Chemosphere 31: 3933-3944.

Rowbotham AL, Levy LS, Shuker LK (2000) Chromium in the environment: an evaluation of exposure of the UK general population and possible adverse health effects. J Toxicol Environ Health 3: 145-178.

Ryan JJ, Dewailly E, Gilman A (1997) Dioxin-like compounds in fishing people from the lower north shore of the St. Lawrence River, Québec, Canada. Arch Environ Health 52: 309-316.

Saavedra P, Arrieta FJ, Jorge A, Mateos CK i cols. (1996) Relationship between chromium levels in plasma and urine and some metabolic alterations associated with diabetes mellitus related to cardiovascular accidents. Metal Ions in Biology and Medicine, Eds. Ph Collery 4: 589-591.

Salway AG, Eggleston HS, Goodwin JW, Murrells TP (1997) UK emissions of air pollutants 1970-1995 (AEAT-1746/Issue 1), Abingdon , UK, AEA Technology.

Sanz P, Nogué S (1990) Intoxicación por metales de origen alimentario. Med Clin 94: 215-217.

Sanz P, Nogué S (1991) Intoxicación por metales de origen laboral. Med Clin 97: 96-98.

Schaller KH, Weltle D, Angerer J (2000) Analytical validity of the determination of mercury in whole blood and urine – Results of the German external assurance programme for toxicological analysis in biological materials. Fresenius J Anal Chem 366: 449-452.

Scherer G, Frank S, Riedel K, Meger-Kossien I, Renner T (2000) Biomonitoring of exposure to polycyclic aromatic hydrocarbons of non occupationally exposed persons. Cancer Epidemiology, Biomarkers & Prevention 9: 373-380.

Schechter AJ, Malkinn R, Päpke O, Ball M, Brandt-Rauf PW (1991) Dioxin levels in blood of municipal incinerator workers. *Med Sci Res* 19: 331-332.

Schrey P, Wittsiepe J, Budde U, Heinzel B, Idel H, Willhelm M (2000) Dietary intake of lead, cadmium, copper and zinc by children from the German North sea island Amrum. *Int J Hyg Environ Health* 203: 1-9.

Schuhmacher M, Batista J, Bosque MA, Domingo JL, Corbella J (1994) Mercury concentrations in marine species from the coastal area of Tarragona Province, Spain. Dietary intake of mercury through fish and seafood consumption. *Sci Total Environ* 156: 269-273.

Schuhmacher M, Bellés M, Rico A, Domingo J, Corbella JL (1996) The impact of reduction of lead in gasoline on the blood and hair lead levels in the population of Tarragona Province, Spain, 1990-1995. *Sci Total Environ* 184: 203-209.

Schuhmacher M, Bosque MA, Domingo JL, Corbella J (1994) Urinary cadmium levels in an unexposed population: relationship to age, sex, and smoking habits. *Trace Elelctr* 11: 38-41.

Schuhmacher M, Domingo JL, Agramunt MC, Gomez M, Llobet JM, Müller L (2000) Internal exposure to organic substances in workers of a new hazardous waste incinerator in Catalonia, Spain. Baseline Levels. *Organohalogen Compounds* 48: 40-43.

Schuhmacher M, Domingo JL, Agramunt MC, Bocio A, Müller L (2002) Biological monitoring if metals and organic substances in hazardous-waste incineration workers. *Int Arch Occup Environ Health* (En premsa).

Schuhmacher M, Domingo JL, Llobet JM, Corbella J (1992) Lead concentration and d-aminolevulinic acid dehydratase activity in the blood of the general population of Tarragona Province, Spain. *Sci Total Environ* 116: 253-259.

Schuhmacher M, Domingo JL, Llobet JM, Lindström G, Wingfors H (1999) Dioxin and dibenzofuran concentrations in blood of a general population from Tarragona Spain. Chemosphere 3: 1123-1133.

Schwartz BS, Stewart WF, Bolla KI, Simon OD, Bandeen-Roche K, Gordon PB, Links JM, Todd AC (2000). Past adult lead exposure is associated with longitudinal decline in cognitive function. Neurology 55: 1144-1150.

Seegal RF, Schantz SL (1994) Neurochemical and behavioral sequelae of exposure to dioxins and PCBs, in: A. Schecter (ed). Dioxins and Health, Plenum Publishing, New York: 409-447.

Seifert B, Becker K, Helm D, Krause C, Schulz C, Seiwert M (2000) The German Environmental Survey 1990/1992 (GerEs II): reference concentrations of selected environmental pollutants in blood, urine, hair, house dust, drinking water and indoor air. Expo Anal Environ Epidemiol 10: 552-565.

Sendelbach LE, Tryka AE, Witschi H (1989) Progressive lung injury over a one-year period after a single inhalation exposure to beryllium sulphate. Am Rev Respir Dis 139: 1003-1009.

Seta N, Arfi C, Momas I (2000) Exposition de la population générale aux dioxines, contaminants ubiquitaires. Rev Epidemiol et Santé Publ 48: 281-293.

Shi Z (1988) Acute nickel carbonil poisoning: a report of 179 cases. Br J Ind Med 43: 422-424.

Slob W (1993) Modeling long-term exposure of the whole population to chemicals in food. Risk Anal 13: 525-530.

Smith CJ, Livingston SD, Doolittle DJ (1997) An international literature survey of “IARC group I carcinogens” reported in mainstream cigarette smoke. Food Chem Toxicol 35: 1107-1130.

Snyder R (2000) Overview of the toxicology of benzene. *J. Tox. Environ Health* 61: 339-346.

Soengas JA, San Miguel A, Gonzalez M, Lopez R (1997) Metales en el organismo: patologias y métodos de determinación. *JANO* 53: 1973-1984.

Staessen J, Amery A, Bernard A, Bruaux P, Buchet JP, Bulpitt CJ, Claeys F, De Plaen P, Ducoffre G, Fagard R, Lauwerys RR, Lijnen P, Nock L, Saint Remy A, Roels H, Rondia D, Sartor F, Thijs L (1991) Blood pressure, the prevalence of cardiovascular diseases, and exposure to cadmium: a population study. *Am J Epidemiol* 134: 257-267.

Startin JR (1994) Dioxins in food. A: Dioxins and Health, Schecter A (ed), Plenum Press, NY.

Stephens RD, Petreas MX, Hayward DG (1995) Biotransfer and bioaccumulation of dioxins and furans from soil: chickens as a model for foraging animals. *Sci Total Environ* 175: 253-273.

Strandberg B, Hites RA (2001) Concentration of organochlorine pesticides in wine corks. *Chemosphere* 44: 729-735.

Sun AY, Yang WL, Kim HD (1993) Free radical and lipid peroxidation in manganese-induced neuronal cell injury. *Ann NY Acad Sci* 679: 358-363.

Sunderman FW (1975) Toxicology and metabolism of nickel compounds. Progress report. Department of Commerce, National Technical Information Service, Springfield, USA.

Sunderman FW (2001) Nasal toxicity, carcinogenicity, and olfactory uptake of metals. *Annals Clinical Lab Sci* 31: 3-24.

Suzuki T, Hongo T, Matsuo N, Imai H, Nakazawa M, Abe T, Yamamura Y, Yoshida M, Aoyama H (1992) An acute mercuric mercury poisoning: chemical speciation of hair mercury shows a peak of inorganic mercury value. *Hum Exp Toxicol* 11: 53-57.

Sweeney A, Symanski E, Burau KD, Kim YJ, Humphrey HEB, Smith MA (2001) Changes in serum PBB and PCB levels over time among women of varying ages at exposure. Environ Res Sect 86: 128-139.

Telisman S, Jurasic J, Pizent A, Cwitkovic P (2001) Blood pressure in relation to biomarkers of lead, cadmium, copper, zinc, and selenium in men without occupational exposure to metals. Environ Res Sect 87: 57-68.

Telolahy P, Javeland B, Cluet J i cols (1993) Urinary excretion of porphyrins by smelter workers chronically exposed to arsenic dust. Toxicol Lett 66: 89-95.

Thomas VM, Socolow RH, Fanelli JJ, Spiro TG (1999) Effects of reducing lead in gasolines: an analysis of the international experience. Environ Sci Technol 33: 3942-3948.

Treble RG, Thompsom TS (1996) Normal values for pentachlorophenol in urine samples collected from a general population. J Anal Toxicol 20: 313-317.

Triebig G, Schaller KH (2000) Mercury, metallic mercury and inorganic mercury compounds. In biological exposure values for occupation toxicants and carcinogens. Critical data evaluation for BAT and BKA values. Deutsche Forschungsgemeinschaft (ed. H Greim and G Lehnert). Wiley-VCH: Weinheim.

Tsuritani I, Honda R, Ishizaki M, Yamada Y, Kido T, Nogawa K (1992) Impairment of vitamin D metabolism due to environmental cadmium exposure, and possible relevance to sex-related differences in vulnerability to the bone damage. J Toxicol Environ Health 37: 519-533.

Vahter M, Berglund M, Akesson A, Lidén C (2002) Metals and women's health. Environ Res Sect 88: 145-155.

van der Berg M, De Jongh J, Poiger H, Olson JR (1994) The toxicokinetics and metabolism of polychlorinated dibenzo-p-dioxins (PCDD/Fs) and dibenzofurans (PCDFs) and their relevance for toxicity. Crit Rev Toxicol 24:1-74.

van der Berg M, Van Virgelen A, Birnbaum L (2000) Consultation on assessment of the health risk of dioxins; re-evaluation of the tolerable daily intake (TDI): Executive Summary. Food Add Cont 17: 223-240.

van der Molen GW, Kooijman SA, Michalek JE, Slob W (1998) The estimation of elimination rates of persistent compounds: a re-analysis of 2,3,7,8-tetrachlorodibenzo-p-dioxin levels in Vietnam veterans. Chemosphere 37: 1833-1844.

van Leeuwen FXR, Feeley M, Schrenk D, Larsen JC Farland W, Younes M (2000) Dioxins: WHO's tolerable daily intake (TDI) revisited. Chemosphere 40: 1095-1101.

Vartiainen T, Lampi P, Tolonen K, Tuomisto J (1995) Polychlorodibenzo-p-dioxin and polychlorodibenzofuran concentrations in lake sediments and fish after a ground water pollution with chlorophenols. Chemosphere 30: 1439-1451.

Vieth B, Heinrich-Hirsch B, Mathar W (2000) Trends in dioxin intake and in human milk levels in Germany. Organohalogen Compd 47: 300-303.

Visvanathan C (1996) Hazardous waste disposal. Resour Conserv Recycl 16: 201-212.

Wasser W, Yusuf S, D'Agati V (1997) Over the counter chromium and renal failure. Ann Intern Med 127: 656.

Watanabe T, Nakatsuka H, Shimbo S (1996) Reduced cadmium and lead burden in Japan in the past 10 years. Int Arch Occup Environ Health 68: 305-314.

WHO (1981) Arsenic. Environmental Health Criteria 18. World Health Organization, Geneva.

WHO (1992). Cadmium. Environmental Health Criteria 134, Geneva.

WHO (1995) Environment Health Criteria 165: Inorganic lead. World Health Organization, Geneva.

WHO (1998) Assessment of the health risk of dioxins: re-evaluation of the tolerable daily intake (TDI). WHO European Centre for Environmental and Health International Programme on Chemical Safety.

WHO/ICPS (1989) Polychlorinated dibenzo-p-dioxins and dibenzofurans. environmental health criteria 88. World Health Organization, Geneva, Switzerland .

Wittsiepe J, Schrey P, Ewers U, Wilhelm M, Selenka F (2000) Decrease of PCDD/F Levels in human blood-trend analysis for the german population, 1991-1996. Environ Res 83: 46-53.

Wrbitzky R, Beyer B, Thoma H, Flatau B, Henning M, Weber A, Angerer J, Lehnert G (2001) Internal Exposure to Polychlorinated Dibenzo-p-Dioxins and Polychlorinated Dibenzofurans (PCDD/Fs) of Bavarian Chimney Sweeps. Arch Environ Contam Toxicol 40: 136-140.

Wrbitzky R, Göen T, Letzel S, Frank F (1995) Internal exposure of waste incineration workers to organic and inorganic substances. Int Arch Occup Environ Health 68: 13-21.

Wu WZ, Li W, Xu Y, Wang JW (2001) Long-term toxic impact of 2,3,7,8-tetrachlorodibenzo-p-dioxin on the reproduction, sexual differentiation, and development of different life stages of Gobiocypris rarus and Daphnia magna. Ecotoxi & Environ Safety 48: 293-300.

Ysart G, Miller P, Croasdale M, CrewwsH, Robb P, Baxter M, de L'Argy C, Harrison N (2000) 1997 UK total diet study--dietary exposures to aluminium, arsenic, cadmium, chromium, copper, lead, mercury, nickel, selenium, tin and zinc. Food Addit Contam 17: 775-786.

Zheng J, Kosmus W, Pichler-Semmelrock F, Köck M (1999) Arsenic speciation in human urine reference materials using high-performance liquid chromatography with inductively couple plasma mass spectrometric detection. J Trace Elements Med Biol 13: 150-156.

Zook DR, Rappe C (1994) Environmental sources, distributions, and fate of polychlorinated dibenzodioxins, dibenzofurans, and related organochlorines. In Dioxins and Health; Schecter A (ed). Plenum Press, New York; 79-113.