

Several enzymes implicated in the ATP metabolism such as an spinach chloroplast ATP synthase [60] and a pea nuclei apyrase [61] are also CK2 *in vitro* substrates. Other plant CK2 targets are a soybean lipoxygenase [62] and a proteasome subunit [63]. In pea nuclei, CK2 has been found in the lamina-matrix and it is able to phosphorylate a lamina-like protein [7].

In yeast and in mammalian cells it has been demonstrated the involvement of CK2 in cell cycle control. Using the synchronizable tobacco BY-2 cell line it has been reported that the Ck2 activity oscillates during the cell cycle, peaking at G1/S and M phases [15]. It is proposed that the CK2 $\beta$  regulatory subunits may play a central role in controlling the level of CK2 activity. Recently, it has been reported that CK2 is involved in the activation of the Salicylic acid-mediated pathway [64], which is one of the most studied pathways in the plant defense reactions. It is the first time in which activation of the CK2 by a plant hormone is reported.

## Acknowledgements

This work was supported by grant BIO2000-1562. M.R. was supported by grant 200 1 TDO C000 12 from Generalitat de Catalunya (Spain).

## References

- Seldin DC, Leder P: Casein kinase II alpha transgene-induced murine lymphoma: Relation to theileriosis in cattle. *Science* 267: 894–896, 1995
- Lüscher B, Kuenzel EA, Krebs EG, Eisenman RN: Myc oncoproteins are phosphorylated by casein kinase II. *EMBO J* 8: 1111–1119, 1989
- Hanna D, Rethinaswamy A, Glover CVC: Casein kinase II is required for cell cycle progression during G1 and G2/M in *Saccharomyces cerevisiae*. *J Biol Chem* 270: 25905–25914, 1995
- Yan T-FJ, Tao M: Purification and characterization of a wheat germ protein kinase. *J Biol Chem* 257: 7037–7043, 1982
- Dobrowolska G, Meggio F, Marchiori F, Pinna LA: Specific determinants of maize casein kinase II-B are related but distinct from those of rat liver casein kinase-2. *Biochem Biophys Acta* 1010: 274–277, 1989
- Klimczak LJ, Schindler U, Cashmore AR: DNA binding activity of the Arabidopsis G-box binding factor GBF1 is stimulated by phosphorylation by casein kinase II from broccoli. *Plant Cell* 4: 87–98, 1992
- Li H, Roux SJ: Casein kinase II protein kinase is bound to lamina-matrix and phosphorylates lamina-like protein in isolated pea nuclei. *Proc Natl Acad Sci USA* 89: 8434–8438, 1992
- Dobrowolska G, Boldyreff B, Issinger O-G: Cloning and sequencing of the casein kinase 2 a subunit from Zea mays. *Biochem Biophys Acta* 1129: 139–140, 1991
- Mizoguchi T, Yamaguchi-Shinozaki K, Hayashida N, Kamada H, Shinozaki K: Cloning and characterization of two cDNAs encoding casein kinase II catalytic subunits in *Arabidopsis thaliana*. *Plant Mol Biol* 21: 279–289, 1993
- Collinge MA, Walker JC: Isolation of an *Arabidopsis thaliana* casein kinase II  $\beta$  subunit by complementation in *Saccharomyces cerevisiae*. *Plant Mol Biol* 25: 649–658, 1994
- Sugano S, Andronis C, Green RM, Wang Z-Y, Tobin EM: Protein kinase CK2 interacts with and phosphorylates the Arabidopsis circadian clock-associated 1 protein. *Proc Natl Acad Sci USA* 95: 11020–11025, 1998
- Niefind K, Guerra B, Pinna LA, Issinger O-G, Schomburg D: Crystal structure of the catalytic subunit of protein kinase CK2 from Zea mays at 2.1 Å resolution. *EMBO J* 17: 2451–2462, 1998
- Riera M, Peracchia G, de Nadal E, Ariño J, Pagès M: Maize protein kinase CK2: Regulation and functionality of three  $\beta$  regulatory subunits. *Plant J* 25: 2001 (in press)
- Lee Y, Lloyd AM, Roux SJ: Antisense expression of the CK2 alpha-subunit gene in Arabidopsis. Effects on light-regulated gene expression and plant growth. *Plant Physiol* 119: 989–1000, 1999
- Espunya MC, Combettes B, Dot J, Chaubet-Gigot N, Martínez MC: Cell-cycle modulation of CK2 activity in tobacco BY-2 cells. *Plant J* 19: 655–666, 1999
- Litchfield DW, Lozemann FJ, Piening C, Sommercorn J, Takio K, Walsch KA, Krebs EG: Subunit structure of casein kinase II from bovine testis. *J Biol Chem* 265: 7638–7644, 1990
- Padmanabha R, Chen-Wu JLP, Hanna DE, Glover CVC: Isolation, sequencing, and disruption of the CKA2 gene: Casein kinase II is essential for viability in *Saccharomyces cerevisiae*. *Mol Cell Biol* 10: 4089–4099, 1990
- Peracchia G, Jensen AB, Culiáñez-Macià FA, Grosset J, Goday A, Issinger O-G, Pagès M: Characterization, subcellular localization and nuclear targeting of casein kinase II from Zea mays. *Plant Mol Biol* 40: 199–211, 1999
- Guerra B, Boldyreff B, Sarno S, Cesaro L, Issinger O-G, Pinna LA: CK2: A protein kinase in need of control. *Pharmacol Ther* 82: 303–313, 1999
- Niefind K, Putter M, Guerra B, Issinger O-G, Schomburg D: GTP plus water mimic ATP in the active site of protein kinase CK2. *Nat Struct Biol* 6: 1100–1103, 1999
- Battistutta R, Sarno S, De Moliner E, Marin O, Issinger O-G, Zanotti G, Pinna LA: The crystal structure of the complex of Zea mays alpha subunit with a fragment of human beta subunit provides the clue to the architecture of protein kinase CK2 holoenzyme. *Eur J Biochem* 267: 5184–5190, 2000
- Livak KJ: Detailed structure of the *Drosophila melanogaster* stellite genes and their transcripts. *Genetics* 124: 303–316, 1990
- Grankowski N, Boldyreff B, Issinger O-G: Isolation and characterization of recombinant human casein kinase II subunits  $\alpha$  and  $\beta$  from bacteria. *Eur J Biochem* 198: 25–30, 1991
- Meggio F, Boldyreff B, Marin O, Marchiori F, Perich JW, Issinger O-G, Pinna LA: Role of  $\beta$  subunit of casein kinase-2 on the stability and specificity of the recombinant reconstituted holoenzyme. *Eur J Biochem* 204: 293–297, 1992
- Bidwai AP, Reed JC, Glover CVC: Phosphorylation of calmodulin by the catalytic subunit of casein kinase II is inhibited by the regulatory subunit. *Arch Biochem Biophys* 300: 265–270, 1993
- Bidwai AP, Reed JC, Glover CVC: Casein kinase II of *Saccharomyces cerevisiae* contains two distinct regulatory subunits,  $\beta$  and  $\beta'$ . *Arch Biochem Biophys* 309: 348–355, 1994
- Saxena A, Padmanabha R, Glover CVC: Isolation and sequencing of cDNA clones encoding the alpha and beta subunits of drosophila casein kinase II. *Mol Cell Biol* 7: 3409–3417, 1987
- Bidwai AP, Zhao W, Glover CVC: A gene located at 56F1-2 in *Drosophila melanogaster* encodes a novel metazoan beta-like subunit of casein kinase II. *Mol Cell Biol Res Commun* 1: 21–28, 1999