

7. BIBLIOGRAFIA

- Aguilar A., Casas C. i Lema J.M. 1995. Degradation of volatile fatty acids by differently enriched methanogenic cultures: Kinetics and inhibition. *Water Res.* **29**(2), 505-509.

- Albaigés, J. 1980. Identification and geochemical significance of long chain acyclic isoprenoid hydrocarbons in crude oils. *Adv. Org. Geochem.* 1979. (Eds. A.G.Douglas i J.R. Maxwell) 19-28. Pergamon Press.

- Aldrich H.C., Robinson R.W. i Williams D.S. 1986. Ultrastructure of *Methanosarcina mazei*. *Syst. Appl. Microbiol.* **7**, 314-319.

- Aldrich H.C., Beimborn D.B. i Schönheit P. 1987. Creation of artifactual internal membranes during fixation of *Methanobacterium thermoautotrophicum*. *Can. J. Microbiol.* **33**, 844-849.

- Balch W.E. i Wolfe R.S. 1976. New approach to the cultivation of methanogenic bacteria: 2-mercaptoethanesulfonic acid (HS-CoM)-dependent growth of *Methanobacterium ruminantium* in a pressurized atmosphere. *Appl. Environ. Microb.* **32**,781-791.

- Baresi L., Mah R.A., Ward D.M. i Kaplan I.R. 1978. Methanogenesis from acetate: Enrichment studies. *Appl. Environ. Microb.* **36** (1), 186- 197.

- Bhattacharya S.K., Uberoi V. i Dronamraju M.M. 1996. Interaction between acetate fed sulfate reducers and methanogens. *Water Res.* **30**(10), 2239-2246.

- Burggraaf S., Stetter K.O., Rouviere P. i Woese C.R. 1991. *Methanopyrus kandleri* : An Archaeal Methanogen Unrelated to all Other Known Methanogens. *Syst. Appl. Microbiol.* **14**, 346-351.

- Cabirol N., Villemur R., Perrier J., Jacob F., Fouillet B., i Chambon P. 1998. Isolation of a methanogenic bacterium, *Methanosarcina* sp strain FR, for its ability to degrade high concentrations of perchloroethylene. Can. J. Microbiol. **44**, 1142-1147.
- Cabrera, M. 1999. Estudi dels biomarcadors de conques altament reductores. Tesi Doctoral. U.P.C. Manresa. 412 pp.
- Cavalier-Smith, T. 1992. Lectures on Cell Evolution, Phylogeny and Macrosystematics. CIRIT. 21-25 Abril 1992, Barcelona.
- Chappe B., Albrecht P i Michaelis W. 1982. Polar lipids of archaebacteria in sediments and petroleums. Science. **217**, 65-66.
- Chaudhuri B.K. i Wiesmann, U. 1996. Kinetic study of the anaerobic degradation of toluene by a mixed culture. Acta Biotechnol. **16**(1), 31-41.
- Choquet G.C., Patel G.B. i Sprott G.D. (1996). Heat sterilization of archaeal liposomes. Can. J. Microbiol. **42**, 183-186.
- Comita P.B. i Gagosian, R.B. 1983. Membrane lipid from deep-sea hydrothermal vent Methanogen: a new macrocyclic glycerol diether. Science **222**, 1329-1331.
- Comita P.B., Gagosian, R.B., Pang & Costello. 1984. Structural elucidation of a unique membrane lipid from a new, extremely thermophilic, deep-sea hydrothermal vent Archaeobacterium, *Methanococcus jannaschii*. J. Biol. Chem. **259**(24), 15234-15241.
- De las Heras i Cisa F.X. 1991. Geoquímica orgànica de conques lacustres fòssils. Institut d'estudis catalans. Arxius de la Secció de Ciències, XCII. 324pp.
- De las Heras i Cisa F.X. 1992. Geoquímica orgànica de sòls. Restauració de sòls. Butll. Soc. Cat. Cièn. **XIII**, 653-661.
- De Rosa M. 1996. Archaeal lipids: structural features and supramolecular organization. Thin Solid Films. **284-285**, 13-17.

- De Rosa M. i Gambacorta A. 1986. Lipid biogenesis in archaeobacteria. *Syst. Appl. Microbiol.* **7**(2-3), 278-285.
- De Rosa M., Bullock J.D., Gambacorta A., Grant W.D., Nicolaus B. i Ross H.N.M. 1982. An asymmetric archeobacterial diether lipid from alkaliphilic halophiles. *J. Gen. Microbiol.* **128**(Feb), 343-348.
- De Rosa M., Albrecht P., Chappe B., Gambacorta A. i Nicolaus B. 1983a. Isoprenoid ethers. Backbone of complex lipids of the archaeobacteria *Sulfolobus solfataricus*. *Biochim. Biophys. Acta.* **753**(2), 249-256.
- De Rosa M., Gambacorta A., Grant W.D., i Nicolaus B. 1983b. A C-25, C-25 diether core lipid from archaeobacterial haloalkaliphiles. *J. Gen. Microbiol.* **129**(Aug), 2333-2337.
- De Rosa M., Gambacorta A., Lanzotti V., Trincone A., Harris J.E. i Grant W.D. 1986. A range of ether core lipids from the methanogenic archaeobacterium *Methanosarcina barkeri*. *Biochim. Biophys. Acta.* **875**, 487-492.
- De Rosa M., Basso A., Gambacorta A., Holz I., Trincone A. i Zillig W. 1987. Lipids of *Thermococcus-Celer*, a sulfur-reducing Archaeobacterium- Structure and Biosynthesis. *Syst. Appl. Microbiol.* **9**(1-2), 1-5.
- De Rosa M., Gambacorta A., Grant W.D., Lanzotti V. i Nicolaus B. 1988. Polar lipids and glycine betaine from haloalkaliphilic archaeobacteria. *J. Gen. Microbiol.* **134** part 1, 205-211.
- De Rosa M., Morana A. i Riccio A. 1994. Lipids of the Archaea: a new tool for bioelectronics. *Biosens. Bioelectron.* **9**, 669-675.
- Demizu K., Ohtsubo S., Kohno S., Miura I., Nishihara M. i Koga Y. 1992. Quantitative determination of Methanogenic cells based on analysis of ether-linked glycerolipids by High-Performance Liquid Chromatography. *J. Ferment. Bioeng.* **73** (2), 135-

139.

- Dolfing J. i Mulder J.W. 1985. Comparison of methane production rate and coenzyme F₄₂₀ content of methanogenic consortia in anaerobic granular sludge. *Appl. Environ. Microb.* **49**(5), 1142-1145.
- Donato M.M., Seleiro E.A. i Da Costa M.S. 1991. Polar lipid and fatty acid composition of strains of *Thermus ruber*. *Syst. Appl. Microbiol.* **14**, 235-239.
- Drzyzga O. i Blotevogel K.H. 1997. Microbial degradation of diphenylamine under anoxic conditions. *Curr. Microbiol.* **35**, 343-347.
- Dyreborg S., Arvin E. i Broholm K. 1997. Biodegradation of NSO-compounds under different redox-conditions. *J. Contaminant Hydrology.* **25**, 177-197.
- Edwards, E.A. i Grbic-Galic. 1994. Anaerobic degradation of toluene and o-xylene by a methanogenic consortium. *Appl. Environ. Microb.* **60**(1), 313-322.
- Elder D.J.E. i Kelly D.J. 1994. The bacterial degradation of benzoic acid and benzenoid compounds under anaerobic conditions: Unifying trends and new perspectives. *FEMS Microbiol. Rev.* **13**, 441-468.
- Ekiel I., Smith I.C.P. i Sprott G.D. 1983. Biosynthetic pathways in *Methanospirillum hungatei* as determined by ¹³C nuclear magnetic resonance. *J. Bacteriol.* **156**, 316-326.
- Eismann F., Becker F., Kusch P. i Stottmeister U. 1996. Alternative electron acceptors in microbial coal-conversion wastewater treatment. *Appl. Microbiol. Biot.* **46**, 604-609.
- Evans R.W., Kates M. i Kuswaha S.C. 1980. The lipids of *Halobacterium marismortui*, an extremely halophilic bacterium in the Dead-Sea. *Biochim. Biophys. Acta.* **619**(3), 533-544.

- Ferrante G., Ekiel I. i Sprott G.D. 1986. Structural characterization of the lipids of *Methanococcus voltae*, including a novel N-acetylglucosamine 1-phosphate diether. J. Biol. Chem. **261**, 17062-17066.
- Ferrante G., Ekiel I. i Sprott G.D. 1987. Structures of diether lipids of *Methanospirillum hungatei* containing novel head groups N,N-dimethylamino- and N,N,N-trimethylaminopentane-tetrol. Biochim. Biophys. Acta. **921**(2), 281-291.
- Ferrante G., Ekiel I., Patel G.B. i Sprott G.D. 1988 a. Structure of the major polar lipids isolated from the acetoclastic methanogen, *Methanotherx concilii* GP6. Biochim. Biophys. Acta. **963**(2), 162-172.
- Ferrante G., Ekiel I., Patel G.B. i Sprott G.D. 1988 b. A novel core Lipid isolated from the Aceticlastic Methanogen *Methanotherx Concilii* GP6. Biochim. Biophys. Acta. **963**, 173-182.
- Ferry J.G. 1993. Fermentation of acetate. A Methanogenesis (ed. J. G. Ferry) Chapman & Hall. Londres. pp. 304-334.
- Finster K., Tanimoto Y. i Bak F. 1992. Fermentation of methanethiol and dimethylsulfide by a newly isolated methanogenic bacterium. Arch. Microbiol. **157**, 425-430.
- Fuhrmann J.A., McCallum K. i Davis A.A. 1992. Novel major archaeobacterial group from marine plankton. Nature. **356**, 148-149.
- Gambacorta A., Trincone A., Nicolaus B., Lama L. i De Rosa M. 1994. Unique Features of lipids of Archaea. Syst. Appl. Microbiol. **16**, 518-527.
- Gliozzi A., Basak T.K, Bruno S., De Rosa M. i Gambacorta A. 1986 a. Organization and dynamics of bipolar lipids from *Sulfolobus solfataricus* in bulk phases and in monolayer membranes. Syst. Appl. Microbiol. **7**(2-3), 266-271.

- Gliozzi A., Paoli G., Pisani D., Gliozzi F., De Rosa M. i Gambacorta A. 1986 b. Phase transitions of bipolar lipids of thermophilic archaeobacteria. *Biochim. Biophys. Acta.* **861**(3), 420-428.
- Grossi V., Hirschler A., Raphael D., Rontani J.F., de Leeuw J.W. i Bertrand J.C. 1998. Biotransformation pathways of phytol in recent anoxic sediments. *Org. Geochem.* **29**(4), 845-861.
- Gulik A., Luzzatti V., De Rosa M. i Gambacorta A. 1986. Structure and polymorphism of tetraether lipids from *Sulfolobus solfataricus*. I: Crystallographic analysis. *Syst. Appl. Microbiol.* **7**(2-3), 258-261.
- Gupta A., Flora J.R.V., Gupta M., Sayles G.D. i Suidan M.T. 1994a. Methanogenesis and sulfate reduction in chemostats-I. Kinetic studies and experiments. *Water Res.* **28**(4), 781-793.
- Gupta A., Flora J.R.V., Sayles G.D. i Suidan M.T. 1994b. Methanogenesis and sulfatereduction in chemostats-II. Model development and verification. *Water Res.* **28**(4), 795-803.
- Hanna K., Bengisgarber C., Kates M., Kogut M. i Kushner D.J. 1984. The effect of salt concentration on the phospholipid and fatty-acid composition of the moderate halophile *Vibrio-costicola*. *Can. J. Microbiol.* **30**(5), 669-675.
- Harada H., Uemura S. i Momonoi K.. 1994. Interactions between sulfate-reducing bacteria and methane-producing bacteria in UASB reactors fed with low strength wastes containing different levels of sulfate. *Water Res.* **28**(2), 355-367.
- Hermann M., Noll K.M., i Wolfe R.S. 1986. Improved agar bottle plate for isolation of methanogens or other anaerobes in a defined gas atmosphere. *Appl. Environ. Microb.* **51**, 1124-1126.
- Holzer G., Oró J. i Tornabene T.G. 1979. Gas chromatographic-mass Spectrometric

analysis of neutral lipids from methanogenic and thermoacidophilic bacteria. J. Chromatogr. **186**(Dec), 795-809.

- Holzer G.U., Kelly P.J. i Jones W.J. 1988. Analysis of lipids from a hydrothermal vent methanogen and associated vent sediment by supercritical fluid chromatography. J. Microbiol. Meth. **8**(3), 161-173.

- Hirata A., Lee H.S., Tsuneda S. i Takai T. 1997. Treatment of photographic processing wastewater using anaerobic-aerobic biofilm reactor. Water Sci.Technol. **36**(12), 91-99.

- Hoefs M.J.L., Shouten S., de Leeuw J. W., King L.L., Wakeham S.G. i Sinninghe Damsté J.S. 1997. Ether lipids of planktonic archaea in the marine water column. Appl. Environ. Microb. **63**(8), 3090-3095.

- James A.G., Watson-Craik I.A. i Senior E. 1998. The effects of organic acids on the methanogenic degradation of the landfill leachate molecules butyrate and valerate. Water Res. **3**, 792-800.

- Jetten M.S.M., Stams A.J.M. i Zehnder A.J.B. 1992. Methanogenesis from acetate: a comparison of the acetate metabolism in *Methanotherix soehngeni* and *Methanosarcina* spp. FEMS Microbiol. Rev. **88**(1992), 181-198.

- Jones W.J., Whitman W.B., Fields R.D. i Wolfe R.S. 1983. Growth and plating efficiency of Methanococci on agar media. Appl. Environ. Microbiol. **46**(1), 220-226.

- Jones W.J., Nagle D.P. i Whitman W.B. 1987. Methanogens and the diversity of Archaeobacteria. Microbiol. Rev. **51**(1), 135-137.

- Jones W.J. i Holzer G.U. 1991. The polar and neutral lipid composition of *Methanosphaera stadtmanae*. Syst. Appl. Microbiol. **14**(2), 130-134.

- Kakinuma K., Obata Y., Matsuzawa T., Uzawa T. i Oshima T. 1990. The

stereochemical fate of glycerol during the biosynthesis of membrane lipids in thermoacidophilic Archaeobacteria *Sulfolobus acidocaldarius*. J. Chem. Soc. Chem. Commun. 925-927.

- Kataoka N., Tokiwa Y. i Takeda K. 1991. Improved technique for identification and enumeration of Methanogenic Bacterial Colonies in Roll Tubes by Epifluorescence Microscopy. Appl. Environ. Microb. **57**(12), 3671-3673.

- Kates M. 1997. Diether and tetraether phospholipids and glycolipids as molecular markers for Archaeobacteria (Archaea). A Molecular markers in environmental geochemistry. cap 13. pp 35-48. American Chemical Society.

- Kates M., Yengoyan L.S. i Sastry P.S. 1965. A diether analog of phosphatidyl glycerophosphate in *Halobacterium cutirubrum*. Biochim. Biophys. Acta **98**, 252-268.

- Kazumi J., Caldwell M.E., Suflita J.M., Lovley D.R. i Young L.Y. 1997. Anaerobic Degradation of Benzene in Diverse Anoxic Environments. Environ. Sci. Technol. **31**(3), 813-818.

- Kim C.G., Clarke W.P. i Lockington D. 1999. Feasibility test of biological degradation of heterocyclic sulfur compounds in anaerobic state. J. Environ. Sci. Health. **A34**(4), 899-918.

- King L.L., Pease T.K. i Wakeham S.G. 1998. *Archaea* in Black Sea water column particulate matter and sediments- evidence from ether lipid derivatives. Org. Geochem. **28**(11), 677-688.

- Koene-Cottaar F.H.M. i Schraa G. 1998. Anaerobic reduction of ethene to ethane in an enrichment culture. FEMS Microbiol. Ecol. **25**, 251-256.

- Koga Y., Ohga M., Nishihara M. i Morii H. 1987. Distribution of a diphytanyl ether analog of phosphatidylserine and an ethanolamine-containing tetraether lipid in methanogenic bacteria. Syst. Appl. Microbiol. **9**, 176-182.

- Koga Y., Nishihara M., Morii H. i Akagawa-Matsushita M. 1993. Polar lipids of Methanogenic Bacteria. *Microbiol. Rev.* **57**, 168-182.
- König H. 1988. Archaeobacterial cell envelopes. *Can. J. Microbiol.* **34**, 395-406.
- König H. i Stetter K.O. 1989. Archaeobacteria A Staley J.T., Bryant M.P., Pfenning N. i Holt J.G. *Bergey's Manual of Systematic Bacteriology.* (Eds Williams i Wilkin) pp. 2171-2253.
- Krajewski-Bertrand M.A., Hayer M., Wolff G., Milon A., Albrecht A.-M., Heissler D., Nakatani Y. i Ourisson G. 1990. Tricyclohexaprenol and an octaprenediol, two of the primitive amphiphilic lipids do improve phospholipidic membranes. *Tetrahedron* **46**, 3143-3154.
- Kreisl P. i Kandler O. 1986. Chemical structure of the cell wall polymer of *Methanosarcina*. *Syst. Appl. Microbiol.* **7**, 293-299.
- Krylova N.I., Janssen P.H. i Conrad R. 1997. Turnover of propionate in methanogenic paddy soil. *FEMS Microbiol. Ecol.* **23**, 107-117.
- Kushwaha S.C., Juez G., Kates M., Kushner D.J. i Rodríguez-Valera F. 1982a. Polar lipids of an Extremely Halophilic bacterial strain (R-4) isolated from salt ponds in Spain. *Biochim. Biophys. Acta.* **711**(1), 19-25.
- Kushwaha S.C., Kates M., Sprott G.D. i Smith I.C.P. 1981a. Novel polar lipids from the methanogen *Methanospirillum hungatei* Gp1. *Biochim. Biophys. Acta.* **664**, 156-173.
- Kushwaha S.J., Juezperez G., Kates M., Kushner D.J. i Rodríguez-Valera F. 1982b. Survey of lipids of a new group of extremely halophilic bacteria from salt ponds in Spain. *Can. J. Microbiol.* **28**(12), 1365-1372.

- Langworthy T.A. 1977. *Biochim. Biophys. Acta.* **487**, 37-50.
- Langworthy T.A. i Pond J.L. 1986. Archaeobacterial ether lipids and chemotaxonomy. *Syst. Appl. Microbiol.* **7**(2-3), 253-257.
- Lanzotti V., De Rosa M., Trincone A., Basso A.L., Gambacorta A., i Zillig W. 1987. Complex lipids from *Desulfurococcus mobilis*, a sulfur reducing archaeobacterium. *Biochim. Biophys. Acta.* **922**(2), 95-102.
- Lanzotti V., Nicolaus B., Trincone A., De Rosa M., Grant W.D. i Gambacorta A. 1989a. A complex lipid with a cyclic phosphate from the archaeobacterium *Nanotrooccus occultus*. *Biochim. Biophys. Acta.* **1001**(1), 31-34.
- Lanzotti V., Nicolaus B., Trincone A., De Rosa M., Grant W.D. i Gambacorta A. 1989b. An isoprenoid ether analogue of phosphatidic acid from a halophilic archaeobacterium. *Biochim. Biophys. Acta.* **1002**(3), 398-400.
- Lanzotti V., Trincone A., Nicolaus B. Zillig W., De Rosa M. i Gambacorta A. 1989c. Complex lipids of *Pyrococcus* and AN1, thermophilic members of archaeobacteria belonging to *Thermococcales*. *Biochim. Biophys. Acta.* **1009**, 44-48.
- Lazrak T., Wolf G., Albrecht A.M., Nakatani Y., Ourisson G. i Kates M. 1988. Bacterioruberins reinforce reconstituted Halobacterium lipid membranes *Biochim. Biophys. Acta.* **939**(1), 160-162.
- Lechevalier H. i Lechevalier M.P. 1988. Chemotaxonomic use of lipids-an overview. *Microbial Lipids*. vol. 1 (eds. Radledge i Wilkinson), 873-902. Academic Press.
- Lelkes P.I., De Rosa M., Gambacorta A., Gliozzi A., Goldenberg D. i Miller I.R. 1983. Vesicles from mixtures of bipolar archaeobacterial lipids with egg phosphatidylcholine.

Biochim. Biophys. Acta. **732**(3), 714-718.

- Li Y-Y., Lam S. i Fang H.H.P. 1996. Interactions between methanogenic, sulfate-reducing and syntrophic acetogenic bacteria in the anaerobic degradation of benzoate. *Water Res.* **7**, 1555-1562.
- Lin C-Y., Chang C-H i Chang F-Y. 1999. Sensitivity of anaerobic sludge biogranule to sulfur compounds. *Biotechnol. Lett.* **21**, 421-423.
- Lomans B.P., Maas R., Luderer R., OP den Camp H.J.B., Pol A., Van Der Drift C. i Vogels G.D. 1999. Isolation and characterization of *Methanomethylovorans hollandica* gen.nov.sp.nov., Isolated from freshwater sediment, a methylotrophic methanogen able to grow on dimethyl sulfide and methanethiol. *Appl. Environ. Microb.* **Aug 1999**, 3641-3650.
- Londry K.L. i Suflita J.M. 1998. Toxicity effects of organosulfur compounds on anaerobic microbial metabolism. *Environ. Toxicol. Chem.* **17**(7), 1199-1206.
- MacDonald D. i Goldfine H. 1991. Effects of solvents and alcohols on the polar lipid composition of *Clostridium butyricum* under conditions of controlled lipid chain composition. *Appl. Environ. Microb.* **57**(12), 3517-3521.
- Madsen T., Rasmussen H.B. i Nilsson L. 1995. Anaerobic biodegradation potentials in digested sludge, a freshwater swamp and a marine sediment. *Chemosphere.* **31**(10), 4243-4258.
- Matthies C. i Schink B. 1993. Anaerobic degradation of long chain dicarboxylic acids by methanogenic enrichment cultures. *FEMS Microbiol. Lett.* **111**, 177-182.
- Mawson A.J., Earle R.L. i Larsen V.F. 1992. Inhibition of a methanogenic population by thioglycollate. *Lett. Appl. Microbiol.* **15**(1992), 100-102.

- McCartney D.M. i Oleszkiewicz J.A. 1993. Competition between methanogens and sulfate reducers: effect of COD: sulfate ratio and acclimation. *Water Environ. Res.* **65**(5), 655-664.
- Moldoveanu N. i Kates M. 1988. Biosynthetic Studies of the polar lipids of *Halobacterium cutirubrum*. Formation of isoprenyl ether intermediates. *Biochim. Biophys. Acta.* **960**(2), 164-182.
- Moldoveanu N., Kates M., Montero C.G. i Ventosa A. 1990. Polar lipids of non-alkaliphilic Halococci. *Biochim. Biophys. Acta.* **1046**(2), 127-135.
- Mollet J. (1991). Balanç d'hidrocarburs aromàtics policíclics a la Depuradora d'Aigües Residuals de Manresa. EUPM. Projecte final de carrera.
- Montero C.G., Ventosa A., Rodriguez-Valera F., Kates M., Moldoveanu N. i Ruiz-Berraquero F. 1989. *Halococcus saccharolyticus* sp.nov., a new species of extremely halophilic non alkaliphilic cocci. *Syst. Appl. Microbiol.* **12**(2), 167-171.
- Morii H. i Koga Y. 1986. Absolute stereochemical configuration of a diphytanyl ether analog of phosphatidylserine of *Methanobrevibacter arboriphilus*. *Biochim. Biophys. Acta.* **879**(1), 103-105.
- Morii H. i Koga Y. 1993. Tetraether type polar lipids increase after logarithmic growth phase of *Methanobacterium thermoautotrophicum* in compensation for the decrease of diether lipids. *FEMS Microbiol. Lett.* **109**(1993), 283-288.
- Morii H. i Koga Y. 1994. Asymmetrical topology of diether-and tetraether-type Polar Lipids in Membranes of *Methanobacterium thermoautotrophicum* Cells. *J. Biol. Chem.* **269**(14), 10492-10497.
- Müller V., Blaut M., i Gottschalk G. 1993. Bioenergetics of Methanogenesis. A Methanogenesis (ed. J. G.Ferry) Chapman & Hall. Londres.pp 360-406.

- Mycke B., Narjes F. i Michaelis W. 1988. Bacteriohopanetetrol from chemical degradation of an oil shale kerogen. *Nature*. **326**, 179-181.
- Nakamoto M., Ueki A. i Ueki K. 1996. Physiological properties of a sulfate-reducing bacterium isolated from municipal sewage sludge and its possible role as a syntrophic acidogen in the ecosystem. *J. Gen. Appl. Microbiol.* **42**, 109-120.
- Nakamura T., Maeda H. Takahashi Y. i Hama Y. 1990. Inevitable generation of primary alcohols during reduction of oxidized lipids with sodium borohydride. *Lipids*. **25**(9), 578-580.
- Nichols P.D., Mancuso C.A. i White D.C. 1987. Measurement of methanotroph and methanogen signature phospholipids for use in assesment of biomass and community structure in model systems. *Org. Geochem.* **11**(6), 451-461.
- Nichols P.D., Shaw P.M., Mancuso C.A. i Franzmann P.D. 1993. Analysis of archaeal phospholipid-derived di- and tetraether lipids by high temperature capillary gas chromatography. *J. Microbiol. Meth.* **18**, 1-9.
- Nicolaus B., Lanzotti V., Trincone A., De Rosa M. i Gambacorta A. 1989. Glycine betaine and polar lipid composition in halophilic archaeobacteria in reponse to growth in different salt concentrations. *FEMS Microbiol. Lett.* **59**, 157-160.
- Nishihara M. i Koga Y. 1987 a. Extraction amd composition of polar lipids from the archaeobacterium *Methanobacterium thermoautotrophicum*: effective extraction of tetraether lipids by an acidified solvent. *J. Biochem.- Tokio* **101**(4), 997-1005.
- Nishihara M., Morii H. i Koga Y. 1987b. Structure Determination of a quartet of novel tetraether lipids from *Methanobacterium thermoautotrophicum*. *J. Biochem.* **101**(4), 1007-1015.

- Nishihara M. i Koga Y. 1991. Hydroxyarchaetidylserine and Hydroxyarchaetidylmyoinositol in *Methanosarcina barkeri*. Polar lipids with a new ether core portion. Biochim. Biophys. Acta. **1082**(2), 211-217.
- Nishihara M. i Koga Y. 1995. Two new phospholipids, hydroxyarchaetidylglycerol and hydroxyarchaetidylethanolamine, from the archaea *Methanosarcina barkeri*. Biochim. Biophys. Acta. **1254**, 155-160.
- Oremland R.S., Kiene P.K., Mathrani I., Whiticar M.J. i Boone D.R. 1989. Description of an estuarine methylotrophic methanogen which grows on dimethyl sulfide. Appl. Environ. Microb. **55**(4), 994-1002.
- Oremland R.S. i Culbertson C.W. 1992. Importance of methane-oxidizing bacteria in the methane budget as revealed by the use of a specific inhibitor. Nature. **356**, 421-426.
- Paltauf F. 1994. Ether lipids in biomembranes. Chemistry and physics of lipids. **74**, 101-139.
- Pease T.K., Van Vleet E.S. i Barre J.S. 1992. Diphytanylglycerol ether distributions in sediments of the Orca Basin. Geochim. Cosmochim. Acta. **56**, 3469-3479.
- Peters K.E. i Moldowan J.M. 1993. The biomarker guide. Prentice-Hall. New Jersey. 363 pp.
- Philp R.P. Fossil fuel biomarkers. Applications and spectra. 1985. Methods in geochemistry and geophysics, 23. Elsevier. Amsterdam. 294 pp.
- Pond J.L., Langworthy T.A. i Holzer G. 1986. Long chain diols: A new class of membrane lipids from a thermophilic bacterium. 1986. Science. **231**, 1134-1136.
- Pugh E.L. i Kates M. 1994. Acylation of proteins of the archaebacteria *Halobacterium cutirubrum* and *Methanobacterium thermoautotrophicum*. Biochim. Biophys. Acta. **1196**, 38-44.

- Putzer K.P., Buchholz L.A., Lidstrom M.E. i Remsen C.C. 1991. Separation of methanotrophic bacteria by using percoll and its application to isolation of mixed and pure cultures. *Appl. Environ. Microb.* **57**(12), 3656-3659.
- Quinn P.J., Brain A.P.R., Steward L.C. i Kates M. 1986. The structure of membrane lipids of the extreme halophile, *Halobacterium cutirubrum*, in aqueous systems studied by freeze-fracture. *Biochim. Biophys. Acta.* **863**(2), 213-223.
- Razo-Flores E., Donlon B., Field J. i Lettinga G. 1996. Biodegradability of N-substituted aromatics and alkylphenols under methanogenic conditions using granular sludge. *Water Sci. Technol.* **33**(3), 47-57.
- Razo-Flores E., Luijten M., Donlon B., Lettinga G. i Field J. 1997. Biodegradation of selected azo dyes under methanogenic conditions. *Water Sci. Technol.* **36** (6-7), 65-72.
- Rorije E., Peijnenburg W.J.G.M. i Klopman G. 1998. Structural requirements for anaerobic biodegradation of organic chemicals: A fragment model analysis. *Environ. Toxicol. Chem.* **17**(10), 1943-1950.
- Ross H.N.M., Collins M.D., Tindall B.J., i Grant W.D. 1981. A rapid procedure for the detection of archaeobacterial lipids in halophilic bacteria. *J. Gen. Microbiol.* **123**, 75-80.
- Rothfuss F. i Conrad R. 1993. Thermodynamics of methanogenic intermediary metabolism in littoral sediment of Lake Constance. *FEMS Microbiol. Ecol.* **12**, 265-276.
- Rowland S.J. 1990. Production of acyclic isoprenoid hydrocarbons by laboratory maturation of methanogenic bacteria. *Org. Geochem* **15**(1), 9-16.
- Russell N.J. 1988. Structural and functional roles of lipids. *Microbial Lipids*. vol 1. (eds. Radledge i Wilkinson) Academic Press. pp. 317-365.
- Salanitro J.P. i Diaz L.A. 1995. Anaerobic biodegradability testing of surfactants.

Chemosphere. **30**(5), 813-830.

- Schink B. 1985. Degradation of unsaturated hydrocarbons by methanogenic enrichment cultures. FEMS Microbiol. Ecol. **31**. 69-77.

- Schink B. 1988. A Zehnder. Biology of anaerobic microorganisms. Wiley-Interscience. Chap 14. Principles and limits of anaerobic degradation: environmental and technological aspects. 771-846.

- Schmidt J.E. i Ahring B.K. 1994. Extracellular polymers in granular sludge from different upflow anaerobic sludge blanket (UASB) reactors. Appl. Microbiol. Biot. **42**, 457-462.

- Schouten S., Van der Maarel M.J.E.C., Huber R. i Sinninghe Damsté J.S. 1997. 2,6,10,15,19-pentamethylcosenes in *Methanobus bombayensis*, a marine methanogenic archaeon, and in *Methanosarcina mazei*. Org. Geochem. **26** (5/6), 409-414.

- Schouten S., Hoefs M.J.L., Koopmans M.P., Bosch H-J. i Sinninghe Damsté J.S. 1998. Structural characterization, occurrence and fate of archaeal ether-bound acyclic and cyclic biphytanes and corresponding diols in sediments. Org. Geochem. **29**(5-7), 1305-1319.

- Sinninghe Damsté J.S., de las Heras F.X.C., Van Bergen P.F. i de Leeuw J.W., 1993. Characterization of tertiary catalan lacustrine oil shales: Discovery of extremely organic sulphur-rich Type I kerogens. Geochim. Cosmochim. Acta. **57**, 389-415.

- Sinninghe Damsté J.S., Schouten S., van Vliet N. H., Huber R. i Geenevasen J. A. J. 1997. A polyunsaturated irregular acyclic C₂₅ isoprenoid in a methanogenic archaeon. Tetrahedron Lett. **38**(39), 6881-6884.

- Skerratt J.H., Nichols P.D., Mancuso C.A., James S.R., Dobson S.J., McMeekin T.A. i Burton H. 1991. The phospholipid ester-linked fatty acid composition of members of the family *Halomonadaceae* and genus *Flavobacterium*: A chemotaxonomic guide. Syst. Appl. Microbiol. **14**, 8-13.

- Smallbone B.W. i Kates M. 1981. Structural identification of minor glycolipids in *Halobacterium cutirubrum*. *Biochim. Biophys. Acta.* **665**(3), 551-558.
- Smet E., Lens P., i Van Lagenhove H.V. 1998. Treatment of waste gases contaminated with odorous sulfur compounds. *Critical Reviews in Environ. Sci. Technol.* **28**(1), 89-117.
- Smith P.F. 1988. Archaeobacteria and other specialized bacteria *Microbial Lipids*. vol 1.(Radledge i Wilkinson eds.) Academic Press. pp. 489-553.
- Smith G.C. i Floodgate G.D.1992. A chemical method for estimating methanogenic biomass. *Continental Shelf Research.* **12**(10), 1187-1196.
- Sprott G.D. i Mckellar R.C. 1980. Composition and properties of the cell wall of *Methanospirillum hungatii*. *Can. J. Microbiol.* **26**(2), 115-120.
- Sprott G.D., Ekiel I. i Dicaire C.J. 1990. Novel, acid-labile, hydroxydiether lipid cores in Methanogenic bacteria. *J. Biol. Chem.* **265**(23), 13735-13740.
- Sprott G.D., Dicaire C.J. i Patel G.B.1994 a. The ether lipids of *Methanosarcina mazei* and other *Methanosarcina* species, compared by fast atom bombardment mass spectrometry. *Can. J. Microbiol.* **40**(1994), 837-843.
- Sprott G.D., Ferrante G. i Ekiel I.1994b. Tetraether lipids of *Methanospirillum hungatei* with head groups consisting of phospho-N,N-dimethylaminopentane-tetrol, phospho-N,N,N-trimethylaminopentane-tetrol, and carbohydrates. *Biochim. Biophys. Acta.* **1214**(1994), 234-242.
- Sprott G. D., Agnew B. J. i Patel G. B. 1997. Structural features of ether lipids in the archaeobacterial thermophiles *Pyrococcus furiosus*, *Methanopyrus kandleri*, *Methanothermus fervidus*, and *Sulfolobus acidocaldarius*. *Can. J. Microbiol.* **43**, 467-476.
- Stams A.J.M. 1994. Metabolic interactions between anaerobic bacteria in methanogenic environments. *Antonie van Leeuwenhoek.* **66**, 271-294.

- Teixidor P. 1996. Biogeoquímica d'alquillglicerols i altres lípids en sistemes deposicionals hipersalins. Tesi Doctoral. Universitat de Barcelona. 278pp.
- Teixidor P., García-Veigas J., Pueyo J. J. i Grimalt J.O. 1993. Lipid composition in Messinian halites from Lorca Basin (SE Iberian Peninsula). Org. Geochem. (Ed. K. Oygard), 554-557. Falch Hurtigtrykk, Oslo (Noruega).
- Tindall B.J. 1990. A comparative study of the lipid composition of *Halobacterium saccharovorum* from various sources. Syst. Appl. Microbiol. **13**, 128-130.
- Tindall B. J., Tomlinson G.A. i Hochstein L.I. 1987. Polar lipid composition of a new halobacterium. Syst. Appl. Microbiol. **9**, 6-8.
- Tornabene T.G. i Langworthy T.A. 1978. Diphytanil and dibiphytanyl glycerol ether lipids of methanogenic Archaeobacteria. Science. **203**, 51-53.
- Torreblanca M., Juez G., Kamekura M., Kates M., Rodriguez-Valera F. i Ventosa A. 1986. Classification of non-alkaliphilic halobacteria based on numerical taxonomy and polar lipid-composition, and description of *Haloarcula* Gen-Nov and *Haloferax* Gen-Nov. Syst. Appl. Microbiol. **8**(1-2), 88-99.
- Trincone A., De Rosa M., Gambacorta A., Lanzotti V., Nicolaus B., Harris J.E., i Grant W.D. 1988. A simple chromatographic procedure for the detection of cyclized archaeobacterial glycerol-bisdiphytanyl-glycerol tetraether core lipids. J. Gen. Microbiol. **134**(12), 3159-3163.
- Trincone A., Lanzotti V., Nicolaus B., Zillig B., De Rosa M. i Gambacorta A. 1989. Comparative lipid composition of aerobically and anaerobically grown *Desulfurolobus ambivalens*, an autotrophic thermophilic archaeobacterium. J. Gen. Microbiol. **135**(10), 2751-2757.
- Trincone A., Nicolaus B., Lama L. De Rosa M., Gambacorta A. i Grant W.D. 1990.

The glycolipid of *Halobacterium sodomense*. J. Gen. Microbiol. **136**(11), 2327-2331.

- Trincone A., Nicolaus B., Palmieri G., De Rosa M., Huber R., Huber G., Stetter K.O. i Gambacorta A. 1992. Distribution of complex and core lipids within new hyperthermophilic members of the Archaea domain. Syst. Appl. Microbiol. **15**(10), 11-17.

- Ueki K., Ueki A., Takahashi K. i Iwatsu M. 1992. The role of sulfate reduction in methanogenic digestion of municipal sewage sludge. J. Gen. Appl. Microbiol. **38**, 195-207.

- Van Eekert M.H.A., Schroder T.H., Stams A.J.M., Schraa G. i Field J.A. 1998a. Degradation and fate of carbon tetrachloride in unadapted Methanogenic Granular Sludge. Appl. Environ. Microb. **64**(7), 2350-2356.

- Van Eekert M.H.A., Van Ras N.J.P., Mentink G.H., Rijnaarts H.H.M., Stams A.J.M., Field J.A. i Schraa G. 1998 b. Anaerobic transformation of β -hexachlorocyclohexane by methanogenic granular sludge and soil microflora. Environ. Sci. Technol. **32**, 3299-3304.

- Vasiliev V.B., Vavilin V.A., Ponomarev i Rytov A.V. 1994. Simulation model of methanogenic community. Revista de Biologia General. Moscú. (en rus). **55**(2), 211-237.

- Vavilin V.A., Vasiliev V.B., i Rytov A.V. 1995. The application of the simulation model "Methane" to the study of organic matter's anaerobic degradation. Revista de Biologia General. Moscú. (en rus). **56**(5), 588-602.

- Vella A.J. i Holzer G. 1990. Ether derived alkanes from sedimentary organic matter. Org. Geochem. **15**(2), 209-214.

- Veltman S., Schoenberg T. i Switzenbaum M. S. 1998. Alcohol and acid formation during the anaerobic decomposition of propylene glycol under methanogenic conditions. Biodegradation. **9**, 113-118.

- Veltman S., Schoenberg T. i Switzenbaum M.S. 1998. Alcohol and acid formation

during the anaerobic decomposition of propylene glycol under methanogenic conditions. *Biodegradation*. **9**, 113-118.

- Whitman W.B., Bowen T.L. i Boone D.R. 1992. The methanogenic bacteria. *A The prokaryotes*. (Eds. Balows *et al*). Springer-Verlag. 719-767. Heidelberg.

- Vila M. 1993. Detecció i caracterització de bacteris metanogènics en el tapís microbià de *Phormidium valderianum* (Salines de la Trinitat, Delta de l'Ebre). Tesi de Llicenciatura. CID-CSIC. 91 pp.

- Vink A., Schouten S., Sephton S. i Sinninghe Damsté J.S. 1998. A newly discovered norisoprenoid, 2,6,15,19-tetramethylcosane, in Cretaceous black shales. *Geochim. Cosmochim. Acta*. **62**(6), 965-970.

- Volkman J.K. i Maxwell J.R. 1986. Acyclic isoprenoids as biological markers. *A Biological Markers in the Sedimentary Records. Methods in Geochemistry and Geophysics* 24 (Ed. R.B Johns.) Elsevier, Amsterdam.

- Woese. 1987. Bacterial Evolution. *Microbiol. Rev.* **51**(2), 221-271.

- Woese C.R. 1992. Prokariote systematics: the evolution of a science. *A: The Prokariotes* (Balows *et al* eds.) Springer-Verlag. Heidelberg.

- Wolfe R.S. 1993. An historical overview of Methanogenesis. *A. Methanogenesis* (ed. James G. Ferry) Chapman & Hall. Londres. pp. 1-32.

- Wollenweber H-W. i Rietschel E.T. 1990. Analysis of lipopolysaccharide (lipid A) fatty acids. *J. Microbiol. Meth.* **11**, 195-211.

- Young L.Y. i Cerniglia C.E. eds. 1995. Microbial transformation and degradation of toxic organic chemicals. Wiley-liss. New York. 631 pp.

- Zehnder. A.J.B. ed. 1988. Biology of anaerobic microorganisms. Wiley-Interscience Publication. 846 pp.

- Zengler K., Richnow H.H., Rosselló-Mora R., Michaelis W. i Widdel F. 1999. Methane formation from long chain alkanes by anaerobic microorganisms. *Nature*. **401**, 266-269.

- Zinder S. H. 1993. Physiological ecology of methanogens. *A Methanogenesis* (ed. J.G Ferry). Chapman & Hall. Londres. pp 128-206.

- Zink G. i Lorber K.E. 1995. Mass spectral identification of metabolites formed by microbial degradation of polycyclic aromatic hydrocarbons. *Chemosphere*. **31**(9), 4077-4084.