

Universidad de Cantabria

Departamento de Física Moderna

CSIC - Universidad de Cantabria

Instituto de Física de Cantabria

**Detection of Point Sources in Maps of the
Cosmic Microwave Background Radiation
by means of Optimal Filters**

A dissertation submitted in partial of the requirements

for the degree of Doctor of Philosophy in Physics

by

Marcos López-Caniego Alcarria

2006

References

- [1] Baccigalupi, C., Bedini, L., Burigana, C., de Zotti, G., Farusi, A., Maino, D., Maris, M., Perrotta, F., Salerno, E., Toffolatti, L. & Tonazzini, A., 2000, "Neural networks and separation of Cosmic Microwave Background and astrophysical signals in sky maps", *MNRAS*, 318, 769.
- [2] Banday, A. J., Wolfendale, A. W., 1991, "Galactic dust emission and the cosmic microwave background", *MNRAS*, 252, 462.
- [3] Barcons, X., 1992, "Confusion noise and source clustering", *ApJ*, 396, 460.
- [4] Barreiro, R. B., Sanz, J. L., Martínez-González, Cayón, L., Silk, J., 1997, "Peaks in the Cosmic Microwave Background: flat versus open models", *ApJ*, 478.
- [5] Barreiro, R. B., Sanz, J. L., Herranz, D., Martínez-González, E., 2003, "Comparing filters for the detection of point sources", *MNRAS*, 342, 119.
- [6] Barreiro, R.B., Hobson, M.P., Banday, A.J., Lasenby, A.N., Stolyarov, V., Vielva, P., Górski, K.M., 2004, "Foreground separation using a flexible maximum-entropy algorithm: an application to COBE data", *MNRAS*, 351, 515.
- [7] Barreiro, R. B., 2006, "Data Analysis in Cosmology: Techniques for compact source extraction on CMB maps", Eds. Martinez, V. J., Martinez-Gonzalez, E., Pons-Borderia, M.J. and Saar, E., Springer-Verlag Lecture Notes in Physics, *astro-ph/0512538*, in press.
- [8] Becker R. H., White R. L., Helfand D. J., 1995, "The FIRST Survey: Faint Images of the Radio Sky at Twenty Centimeters", *ApJ*, 450, 559.
- [9] Bedini, L., Herranz, D., Salerno, E., Baccigalupi, C., Kuruoğlu, E.E., Tonazzini, A., 2004, "Separation of correlated astrophysical sources using multiple-lag covariance matrices", EURASIP JASP Special Issue "Applications of Signal Processing in Astrophysics and Cosmology".

- [10] Bennett, C. L., et al., 1996, "Four-Year COBE DMR Cosmic Microwave Background Observations: Maps and Basic Results", *ApJ*, 464, L1.
- [11] Bennett, C. L., et al., 2003, "First-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Preliminary Maps and Basic Results", *ApJS*, 148, 1.
- [12] Bijaoui, A., Bobichon, Y. and Vandame, B., 1997, "Multiscale image fusion in Astronomy", *Vistas in Astronomy*, 41, 365.
- [13] Blum, R.S., 2004, "Robust image fusion using a statistical signal processing approach", *Information Fusion*, in press.
- [14] Bond, J. R., Szalay, A. S., 1983, "The collisionless damping of density fluctuations in an expanding universe", *ApJ*, 274, 443.
- [15] Bond, J. R., Efstathiou G., 1987, "The statistics of cosmic background radiation fluctuations", *MNRAS*, 226, 655.
- [16] Bouchet, F. R., Gispert, R., Boulanger, F., Puget, J.-L., 1997, "Comparing Performances of CMB Experiments", Proceedings of the Moriond Astrophysics meeting, 481.
- [17] Bouchet, F.R., Gispert, R., 1999, "Foregrounds and CMB experiments I. Semi-analytical estimates of contamination", *New Astronomy*, 4, 6, 443.
- [18] Bouchet, F. R., Prunet, S., Sethi, S. K., 1999, "Multifrequency Wiener filtering of cosmic microwave background data with polarization", *MNRAS*, 302, 663.
- [19] Cayón, L., Sanz, J. L., Barreiro, R. B., Martínez-González, E., Vielva, P., Toffolatti, L., Silk, J., Diego, J. M., Argüeso, F., 2000, "Isotropic wavelets: a powerful tool to extract point sources from cosmic microwave background maps", *MNRAS*, 315, 757, 761.
- [20] Chiang, L.Y., Jørgensen, H.E., Naselsky I.P., Naselsky P.D., Novikov I.D., Christensen, P.R., 2002, "An adaptive filter for the construction of the Planck Compact Source Catalogue", *MNRAS*, 335, 1054.
- [21] Cleary K. A., et al., 2005, "Source subtraction for the extended Very Small Array and 33-GHz source count estimates", *MNRAS*, 360, 340.
- [22] Colafrancesco, S., Mazzotta, P., Rephaeli, Y., Vittorio, N. 1997, "Intracluster Comptonization of the Cosmic Microwave Background: Mean Spectral Distortion and Cluster Number Counts", *ApJ*, 479, 1.

- [23] Condon J.J., Cotton W.D., Greisen E.W., Yin Q.F., Perley R.A., Taylor G.B., c Broderick J.J., 1998, "The NRAO VLA Sky Survey", AJ, 115, 1693.
- [24] Cremonese, G., Marzari, F., Burigana, C., Maris, M., 2002, "Asteroid detection at millimetric wavelengths with the PLANCK survey", New Ast., 7, 483.
- [25] Cruz, M., Tucci, M., Martínez-González, E., Vielva, P., 2006, " The non-Gaussian Cold Spot in WMAP: significance, morphology and foreground contribution", MN-RAS, submitted (astro-ph/0601427).
- [26] Corey, B. E., Wilkinson, D. T., 1976, "A Measurement of the Cosmic Microwave Background Anisotropy at 19 GHz", BAAS, 8, 351.
- [27] Daubechies, I., 1988, "Orthonormal bases of compactly supported wavelets", Comm. Pure Appl. Math., 41, 909.
- [28] de Bernardis, P., et al., 2000, "A flat Universe from high-resolution maps of the cosmic microwave background radiation", Nature, 404, 955.
- [29] Delabrouille J., Cardoso J.-F., Patanchon G., 2003, MNRAS, "Multidetector multi-component spectral matching and applications for cosmic microwave background data analysis", 346, 1089.
- [30] de Zotti, G., Franceschini, A., Toffolatti, L., Mazzei, P., Danese, L., 1996, "Extragalactic Backgrounds: I. General Formalism", ApL&C, 35, 289.
- [31] de Zotti, G., Toffolatti, L., Argüeso, F., Davies, R. D., Mazzotta, P., Partridge, R. B., Smoot, G. F., Vittorio, N., 1999, "The Planck Surveyor mission: astrophysical prospects", AIPC, 476, 204.
- [32] de Zotti, G., Ricci, R., Mesa, D., Silva, L., Mazzotta, P., Toffolatti, L., González-Nuevo, J., 2005, "Predictions for high-frequency radio surveys of extragalactic sources", A&A, 431, 893.
- [33] De Zotti, G., Burigana, C., Negrello, M., Tinti, S., Ricci, R., Silva, L., González-Nuevo, J., Toffolatti, L., 2006, in "The many scales in the Universe", ed. J.C. Del Toro Iniesta et al., Springer, p. 45.
- [34] Dennison, B., Topasna, G. A., Simonetti, J. H., 1996, "An Imaging Survey of the Galactic H alpha Emission", AAS, 28, 889.
- [35] Desert, F.-X., Boulanger, F., Puget, J. L., 1990, "Interstellar dust models for extinction and emission", A&A, 237, 215.

- [36] Dicke, R. H., Peebles, P. J. E., Roll, P. G., Wilkinson, D. T., 1965, "Cosmic Black-Body Radiation.", *ApJ*, 142, 414.
- [37] Diego J. M., Vielva P., Martínez-González E., Silk J., Sanz J. L., 2002, "A Bayesian non-parametric method to detect clusters in Planck data", *MNRAS*, 336, 1351.
- [38] Dickinson, C., Davies, R. D., Davis, R. J., 2003, "Towards a free-free template for CMB foregrounds", *MNRAS*, 341, 369.
- [39] Dolag, K., Hansen, F.K., Roncarelli, M., Moscardini L., 2005, "The imprints of local superclusters on the Sunyaev-Zel'dovich signals and their detectability with Planck", *MNRAS*, 363, 1, 29.
- [40] Draine, B. T. & Lazarian, A., 1998, "Electric Dipole Radiation from Spinning Dust Grains", *ApJ*, 508, 157.
- [41] Draine B. T. & Lazarian A., 1998, "Diffuse Galactic Emission from Spinning Dust Grains", *ApJ*, 494, L19.
- [42] Dunlop, J. S., Peacock, J. A., 1990, "The Redshift Cut-Off in the Luminosity Function of Radio Galaxies and Quasars", *MNRAS*, 247, 19.
- [43] Eddington A. S., 1940, "The correction of statistics for accidental error", *MNRAS*, 100, 354.
- [44] Finkbeiner, D. P., Davis, M., Schlegel, D. J., 1999, "Extrapolation of Galactic Dust Emission at 100 Microns to Cosmic Microwave Background Radiation Frequencies Using FIRAS", *ApJ*, 524, 867.
- [45] Fixsen, D. J., Cheng, E. S., Gales, J. M., Mather, J. C., Shafer, R. A., Wright, E. L., 1996, "The Cosmic Microwave Background Spectrum from the Full COBE FIRAS Data Set", *ApJ*, 473, 576.
- [46] Forni O., Aghanim N., 2004, "Separating the kinetic Sunyaev-Zel'dovich effect from primary cosmic microwave background fluctuations", *A&A*, 420, 49.
- [47] Franceschini, A., Toffolatti, L., Danese, L., de Zotti, G., 1989, "Discrete source contributions to small-scale anisotropies of the microwave background", *ApJ*, 344, 35.
- [48] Gaustad J. E., McCullough P. R., Rosing W., Van Buren D., 2001, "A Robotic Wide-Angle H alpha Survey of the Southern Sky", *PASP*, 113, 1326.

- [49] Giardino, G., Banday, A. J., Górski, K. M., Bennett, K., Jonas, J. L., Tauber, J., 2002, "Towards a model of full-sky Galactic synchrotron intensity and linear polarisation: A re-analysis of the Parkes data", *A&A*, 387, 82.
- [50] González-Nuevo, J., Toffolatti, L. & Argüeso, F., "Predictions on the angular power spectrum of clustered extragalactic point sources at CMB frequencies from flat and all-sky 2D-simulations", 2005, *ApJ*, 621, 1.
- [51] González-Nuevo, J., Argüeso, F., López-Caniego, M., Toffolatti, L., Sanz, J. L., Vielva, P., Herranz, D., 2006, "The Mexican Hat Wavelet Family. Application to point source detection in CMB maps", *MNRAS*, 369, 1603.
- [52] Górski, K. M., Hivon, E., Banday, A. J., Wandelt, B. D., Hansen, F. K., Reinecke, M., Bartelmann, M., 2005, "HEALPix: A Framework for High-Resolution Discretization and Fast Analysis of Data Distributed on the Sphere", *ApJ*, 622, 759.
- [53] Granato, G. L., Silva, L., Monaco, P., Panuzzo, P., Salucci, P., De Zotti, G. Danese, L., 2001, "Joint formation of QSOs and spheroids: QSOs as clocks of star formation in spheroids", *MNRAS*, 324, 757.
- [54] Granato, G. L., De Zotti, G., Silva, L., Bressan, A. Danese, L., 2004, "A Physical Model for the Coevolution of QSOs and Their Spheroidal Hosts", *ApJ*, 600, 580.
- [55] Gregory P.C., Scott W.K., Douglas K., Condon J.J., 1996, "The GB6 Catalog of Radio Sources", *ApJS*, 103, 427.
- [56] Griffith M.R., Wright A.E., Burke B.F., Ekers R.D., 1994, "The Parkes-MIT-NRAO (PMN) surveys. 3: Source catalog for the tropical survey (-29 deg less than delta less than -9 deg .5)", *ApJS*, 90, 179.
- [57] Griffith M.R., Wright A.E., Burke B.F., Ekers R.D., 1995, "The Parkes-MIT-NRAO (PMN) surveys. 6: Source catalog for the equatorial survey (-9.5 deg less than delta less than +10.0 deg)", *ApJS*, 97, 347.
- [58] Guiderdoni, B., Hivon, E., Bouchet, F.R., Maffei, B., 1998, "Semi-analytic modelling of galaxy evolution in the IR/submm range", *MNRAS*, 295, 877.
- [59] Guth, A. H., 1981, "Inflationary universe: A possible solution to the horizon and flatness problems", *PhRvD*, 23, 347.
- [60] Haehnelt M. G., Tegmark M., 1996, *MNRAS*, "Using the Kinematic Sunyaev-Zeldovich effect to determine the peculiar velocities of clusters of galaxies", 279, 545.

- [61] Haffner, L. M., 1999, PhDT.
- [62] Haffner, L.M., Reynolds, R.J., Tufte, S.L., Madsen, G.J., Jaehnig, K.P., Percival, J.W., 2003, "The Wisconsin H-alpha Mapper Northern Sky Survey", ApJS, 149, 405.
- [63] Hansen, F.K., Branchini, E., Mazzotta, P., Cabella, P., Dolag, K., 2005, "A full-sky prediction of the Sunyaev-Zeldovich effect from diffuse hot gas in the local universe and the upper limit from the WMAP data", MNRAS, 361, 753.
- [64] Hanany, S., et al., 2000, "MAXIMA-1: A Measurement of the Cosmic Microwave Background Anisotropy on Angular Scales of $10' - 5''$ ", ApJ, 545, L5.
- [65] Haslam, C. G. T., Stoffel, H., Salter, C. J., Wilson, W. E., 1982, "A 408 MHz all-sky continuum survey. II - The atlas of contour maps", A&AS, 47, 1.
- [66] Henkel B., Partridge R. B., 2005, "Completing the Counts of Radio Sources at 8.5 GHz", ApJ, 635, 950.
- [67] Herranz, D., 2002, "Análisis de las anisotropías de la Radiación Cósmica del Fondo de Microondas mediante el uso de Filtros Adaptados", PhD Thesis. Universidad de Cantabria, Spain. ISBN 84-689-8244-X.
- [68] Herranz, D., Sanz, J. L., Barreiro, R. B., Martínez-González, E., 2002, "Scale-adaptive Filters for the Detection/Separation of Compact Sources", ApJ, 580, 610.
- [69] Herranz, D., Gallegos, J., Sanz, J. L., Martínez-González, E., 2002, "Point source detection and extraction from simulated Planck time-ordered data using optimal adaptive filters", MNRAS, 334, 533.
- [70] Herranz, D., Sanz, J. L., Hobson, M. P., Barreiro, R. B., Diego, J. M., Martínez-González, E., Lasenby, A. N., 2002, "Filtering techniques for the detection of Sunyaev-Zel'dovich clusters in multifrequency maps", MNRAS, 336, 1057.
- [71] Herranz D., Sanz J. L., Barreiro R. B., López-Caniego M., 2005, "The estimation of the Sunyaev-Zel'dovich effects with unbiased multifilters", MNRAS, 356, 944
- [72] Herranz D., Sanz J. L., López-Caniego M., González-Nuevo J., 2006, "A Bayesian approach to flux correction in extragalactic source detection", to appear in Proceedings of IEEE ISSPIT 2006.
- [73] Hinshaw, G., et.al., 2006, "Three-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Temperature Results", ApJ submitted.

- [74] Hobson, M.P., Jones, A.W., Lasenby, A.N., Bouchet F.R., 1998, "Foreground separation methods for satellite observations of the cosmic microwave background", 1998, MNRAS, 300, 1.
- [75] Hobson, M. P., Barreiro, R. B., Toffolatti, L., Lasenby, A. N., Sanz, J. L., Jones, A. W., Bouchet, F. R., 1999, "The effect of point sources on satellite observations of the cosmic microwave background", MNRAS, 306, 232.
- [76] Hobson, M. P., McLachlan, C., 2003, "A Bayesian approach to discrete object detection in astronomical data sets", MNRAS, 338, 765.
- [77] Hogg D. W., Turner E. L., 1998, "A Maximum Likelihood Method to Improve Faint-Source Flux and Color Estimates", PASP, 110, 727.
- [78] Jarosik N., et al., 2006, ApJS submitted (astro-ph/0603452)
- [79] Jackson, C. A., Wall, J. V., 1999, "Extragalactic radio-source evolution under the dual-population unification scheme", MNRAS, 304, 160.
- [80] Jonas, J. L., Baart, E. E., Nicolson, G. D., 1998, "The Rhodes/HartRAO 2326-MHz radio continuum survey", MNRAS, 297, 977.
- [81] Jones B.J.T. & Wyse R.F.G., 1985, "The ionisation of the primeval plasma at the time of recombination", A&A, 149, 144.
- [82] Kovac J.M., Leitch E.M., Pryke C., Carlstrom J.E., Halverson N.W., Holzapfel W.L., 2002, "Measurement of polarization with the Degree Angular Scale Interferometer", Nature, 420, 772.
- [83] Kogut A., 1999, "Anomalous Microwave Emission", ASPC, 181, 91.
- [84] Kühr H., Witzel A., Pauliny-Toth I. I. K., Nauber U., 1981, "A catalogue of extragalactic radio sources having flux densities greater than 1 Jy at 5 GHz" A&AS, 45, 367.
- [85] Leitch, E. M., Readhead, A. C. S., Pearson, T. J., Myers, S. T., 1997, "An Anomalous Component of Galactic Emission", ApJ, 486, L23.
- [86] López-Caniego, M., Herranz, D., Barreiro, R. B., Sanz, J. L., 2004, "A Bayesian approach to filter design: detection of compact sources", SPIE, 5299, 145L.
- [87] López-Caniego, M., Herranz, D., Barreiro, R. B., Sanz, J. L., 2005, "Filter design for the detection of compact sources based on the Neyman-Pearson detector", MNRAS, 359, 993.

- [88] López-Caniego, M., Herranz, D., Sanz, J. L., Barreiro, R. B., 2005, "Detection of Point Sources on Two-Dimensional Images Based on Peaks", EURASIP, Special Issue on "Applications of Signal Processing in Astrophysics and Cosmology", 15, 2426.
- [89] López-Caniego, M., Sanz, J. L., Herranz, D., González-Nuevo, J., Barreiro, R. B., Kuruoglu, E., 2005, "Non-linear fusion of images and the detection of point sources", Nonlinear Signal and Image Processing, IEEE-Eurasip, 1502222.
- [90] López-Caniego M., Herranz D., Gonález-Nuevo, J., Sanz, J. L., Barreiro, R. B., Vielva, P., Argüeso, F., Toffolatti, L., 2005, "Comparison of filters for the detection of point sources in Planck simulations", MNRAS, 370, 2047.
- [91] López-Caniego, M., Gonález-Nuevo, J., Massardi, M., Herraz, D., Sanz, J. L., De Zotti, G., Toffolatti, L., Argüeso, F., 2006, 'Non-blind catalogue of extragalactic point sources from the Wilkinson Microwave Anisotropy Probe (WMAP) first 3-year survey data', ApJS submitted.
- [92] Li, S., Kwok, J. T. and Wang, Y., 2001, "Combination of images with diverse focuses using the spatial frequency", Information Fusion, 2, 169.
- [93] Maino, D., Farusi, A., Baccigalupi, C., Perrotta, F., Banday, A. J., Bedini, L., Burigana, C., De Zotti, G., Górski, K. M., Salerno, E., 2002, "All-sky astrophysical component separation with Fast Independent Component Analysis (FastICA)", MNRAS, 334, 53.
- [94] Maino, D., Donzelli, S., Banday, J., Stivoli, F., Baccigalupi, C., , "CMB signal in WMAP 3yr data with FastICA", MNRAS, submitted.
- [95] Maitre, H. and Bloch, I., 1997, "Image fusion", Vistas in Astronomy, 41, 329.
- [96] Mandolesi, N. et al. 1998, proposal submitted to ESA for the Planck Low Frequency Instrument.
- [97] Martinez-Gonzalez, E., Sanz, J. L., 1990, "CMB Anisotropies Generated by Cosmic Voids and Great Attractors", MNRAS, 247, 473.
- [98] Mason B.S., Pearson T.J., Readhead A.C.S., et al., 2003, "The Anisotropy of the Microwave Background to $l = 3500$: Deep Field Observations with the Cosmic Background Imager", ApJ, 591, 540.
- [99] Mather, J. C., et al., 1994, "Measurement of the cosmic microwave background spectrum by the COBE FIRAS instrument", ApJ, 420, 439.

- [100] Mather, J. C., Fixsen, D. J., Shafer, R. A., Mosier, C., Wilkinson, D. T., 1999, "Calibrator Design for the COBE Far-Infrared Absolute Spectrophotometer (FIRAS)", *ApJ*, 512, 511.
- [101] Mauch T., Murphy T., Buttery H.J., Curran J., Hunstead R.W., Piestrzynski B., Robertson J.G., Sadler E.M., 2003, "SUMSS: a wide-field radio imaging survey of the southern sky - II. The source catalogue", *MNRAS*, 342, 1117.
- [102] McEwen, J. D., Hobson, M. P., Lasenby, A. N., Mortlock D. J., 2005, "A high-significance detection of non-Gaussianity in the Wilkinson Microwave Anisotropy Probe 1-yr data using directional spherical wavelets", *MNRAS*, 359, 1583.
- [103] Muxlow, T. W. B., et al., 2005, "High-resolution studies of radio sources in the Hubble Deep and Flanking Fields", *MNRAS*, 358, 1159.
- [104] Negrello, M., Magliocchetti, M., Moscardini, L., De Zotti, G., Granato, G.L., Silva, L., 2004, "Confusion noise at far-infrared to millimetre wavelengths", *MNRAS*, 352, 493.
- [105] Negrello, M., González-Nuevo, J., Magliocchetti, M., Moscardini, L., De Zotti, G., Toffolatti, L., Danese, L. 2005, "Effect of clustering on extragalactic source counts with low-resolution instruments", *MNRAS*, 358, 869.
- [106] de Oliveira-Costa, A., Kogut, A., Devlin, M. J., Netterfield, C. B., Page, L. A., Wollack, E. J., 1997, "Galactic Microwave Emission at Degree Angular Scales", *ApJ*, 482, L17.
- [107] Padmanabhan, T., 1993, "Structure formation in the universe", Cambridge Univ. Press, New York.
- [108] Page L., et al., 2003, "First-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Interpretation of the TT and TE Angular Power Spectrum Peaks", *ApJS*, 148, 233.
- [109] Paladini, R., Burigana, C., Davies, R.D., Maino, D., Bersanelli, M., Cappellini, B., Platania, P., Smoot, G., 2003, "A radio catalog of Galactic HII regions for applications from decimeter to millimeter wavelengths", *A&A*, 397, 213.
- [110] Patanchon G., Cardoso J.-F., Delabrouille J., Vielva P., 2005, *MNRAS*, "Cosmic microwave background and foregrounds in Wilkinson Microwave Anisotropy Probe first-year data", 364, 1185.

- [111] Penzias, A. A., Wilson R. W., 1965, "A Measurement of Excess Antenna Temperature at 4080 Mc/s.", *ApJ*, 142, 419.
- [112] Piella, G., "A general framework for multiresolution image fusion: from pixels to regions", *Information Fusion*, vol. 4, pp. 259-280, 2003.
- [113] Planck Collaboration, 2005, Planck: The Scientific Programme, ESA-SCI(2005)1, astro-ph/0604069.
- [114] Postman, M., Lubin, L. M., Gunn, J. E., Oke, J. B., Hoessel, J. G., Schneider, D. P., Christensen, J. A., 1996, "The Palomar Distant Cluster Survey: I. The Cluster Catalog", *AJ*, 615, 111.
- [115] Puget, J.L. et al. 1998, proposal submitted to ESA for the Planck High Frequency Instrument.
- [116] Reich, P., Reich, W., 1986, "A radio continuum survey of the northern sky at 1420 MHz. II", *A&AS*, 63, 205.
- [117] Rees M. J., Sciama D. W., 1968, "Larger scale Density Inhomogeneities in the Universe", *Nature*, 217, 511
- [118] Rice, S. O., 1954, "Mathematical Analysis of Random Noise" in "Selected papers on Noise and Stochastic Processes", N. Wax, ed., Dover Publ. Inc. (N. Y.), 133.
- [119] Ricci R., et al., 2004, "First results from the Australia Telescope Compact Array 18-GHz pilot survey" *MNRAS*, 354, 305.
- [120] Rybicki, G. B., Lightman, A. P., 1979, "Radiative processes in astrophysics", Wiley, N.Y..
- [121] Rybicki, G. B., Press W. H., 1992, "Interpolation, realization, and reconstruction of noisy, irregularly sampled data", *ApJ*, 398, 169.
- [122] Sachs, R. K., Wolfe, A. M., 1967, "Perturbations of a Cosmological Model and Angular Variations of the Microwave Background", *ApJ*, 147, 73.
- [123] Sanz, J. L., Argüeso, F., Cayón, L., Martínez-González, E., Barreiro, R. B. & Toffolatti, L. 1999, "Wavelets applied to cosmic microwave background maps: a multiresolution analysis for denoising", *MNRAS*, 309, 672.
- [124] Sanz, J. L., Herranz, D., Martínez-González, E., 2001, "Optimal Detection of Sources on a Homogeneous and Isotropic Background", *ApJ*, 552, 484.

- [125] Schlegel, D. J., Finkbeiner, D. P., Davis, M., 1998, "Maps of Dust Infrared Emission for Use in Estimation of Reddening and Cosmic Microwave Background Radiation Foregrounds", *ApJ*, 500, 525.
- [126] Schrater, P. R., 2003, "Bayesian data fusion and credit assignment in vision and fMRI data analysis", *SPIE*, 5016, 24.
- [127] Schulz A. E., White M., 2003, "Surveys of Galaxy Clusters with the Sunyaev-Zel'dovich Effect", *ApJ*, 586, 723.
- [128] Silk, J., 1968, "Cosmic Black-Body Radiation and Galaxy Formation", *ApJ*, 151, 459.
- [129] Smoot, G. F., et al., 1992, "Structure in the COBE differential microwave radiometer first-year maps", *ApJ*, 396, L1.
- [130] Smoot, G. F., Galactic Free-Free and H_α emission, *astro-ph/9801121*.
- [131] Smoot, G. F., 1999, "CMB Synchrotron Foreground", *ASPC*, 181, 61.
- [132] Spergel, D. N. et al., 2003, "First-Year Wilkinson Microwave Anisotropy Probe (WMAP)1 Observations: Preliminary Maps and Basic Results", *ApJS*, 148, 175.
- [133] Starck, J-L., Murtagh, F. and Bijaoui, A., *Image Processing and Data Analysis: the Multiscale Approach*, Cambridge University Press, U.K, 1998.
- [134] Stolyarov, V., Hobson, M.P., Ashdown, M.A.J., Lasenby, A.N., 2002, "All-sky component separation for the Planck mission", *MNRAS*, 336, 97.
- [135] Barreiro R. B., Hobson M. P., Banday A. J., Lasenby A. N., Stolyarov V., Vielva P., Górski K. M., 2004, "All-sky component separation in the presence of anisotropic noise and dust temperature variations", *MNRAS*, 351, 515
- [136] Sunyaev R. A., Zeldovich Y. B., 1972, "The Observations of Relic Radiation as a Test of the Nature of X-Ray Radiation from the Clusters of Galaxies", *CoASP*, 4, 173.
- [137] Tauber, J. A., 2004, *AdSpR*, 34, 491.
- [138] Tegmark, M., Efstathiou, G., 1996, "A method for subtracting foregrounds from multifrequency CMB sky maps", *MNRAS*, 281, 1297.
- [139] Tegmark, M., de Oliveira-Costa, A., 1998, "Removing Point Sources from Cosmic Microwave Background Maps", *ApJ*, 500, L83.

- [140] Tegmark, M., Eisenstein, D. J., Hu, W., de Oliveira-Costa A., 2000, "Foregrounds and Forecasts for the Cosmic Microwave Background", *ApJ*, 530, 133.
- [141] Toffolatti, L., Argüeso Gómez, F., de Zotti, G., Mazzei, P., Franceschini, A., Danese, L., Burigana, C., 1998, "Extragalactic source counts and contributions to the anisotropies of the cosmic microwave background: predictions for the Planck Surveyor mission", *MNRAS*, 297, 117.
- [142] Toffolatti, L., Argüeso, F., De Zotti, G., and Burigana, C., 1999, in *ASP Conf. Series*, Vol. 181, *Microwave Foregrounds*, A. de Oliveira-Costa and M. Tegmark eds., p.153
- [143] Tucci, M., Martínez-González, E., Toffolatti, L., González-Nuevo, J., De Zotti, G., 2004, "Predictions on the high-frequency polarization properties of extragalactic radio sources and implications for polarization measurements of the cosmic microwave background", *MNRAS*, 349, 1267.
- [144] Vielva, P., Martínez-González, E., Cayón, L., Diego, J. M., Sanz, J. L., Toffolatti, L., 2001a, "Predicted Planck extragalactic point-source catalogue", *MNRAS*, 326, 181.
- [145] Vielva, P., Barreiro, R. B., Hobson, M. P., Martínez-González, E., Lasenby, A. N., Sanz, J. L., Toffolatti, L., 2001b, "Combining maximum-entropy and the Mexican hat wavelet to reconstruct the microwave sky", *MNRAS*, 328, 1.
- [146] Vielva P., Martínez-González, E., Gallegos, J. E., Toffolatti, L., Sanz, J. L., 2003, "Point source detection using the Spherical Mexican Hat Wavelet on simulated all-sky Planck maps", *MNRAS*, 344, 89.
- [147] Vielva, P., Martínez-González, E., Barreiro, R. B., Sanz, J. L., Cayón, L., 2004, "Detection of Non-Gaussianity in the Wilkinson Microwave Anisotropy Probe First-Year Data Using Spherical Wavelets", *ApJ*, 609, 22.
- [148] Vio, R., Andreani, P., Wamsteker, W., 2004, "Some good reasons to use matched filters for the detection of point sources in CMB maps", *A&A*, 414, 17.
- [149] Waldram E.M., Pooley G.G., Grainge K., Jones M.E., Saunders R.D.E., Scott P.F., Taylor A.C., 2003, "9C: a survey of radio sources at 15 GHz with the Ryle Telescope", *MNRAS*, 342, 915.
- [150] Wang, Y. and Lohmann, B., "Multisensor image fusion: concept, method and applications", Institut für Automatisierungstechnik, Universität Bremen, Tech. Rep., Dec. 2000.

- [151] Watson R. A., Rebolo R., Rubiño-Martín J. A., Hildebrandt S., Gutiérrez C. M., Fernández-Cerezo S., Hoyland R. J., Battistelli E. S., 2005, "Detection of Anomalous Microwave Emission in the Perseus Molecular Cloud with the COSMOSOMAS Experiment", *ApJ*, 624, L89.
- [152] White, M., Scott, D., Silk, J., 1994, "Anisotropies in the Cosmic Microwave Background", *ARA&A*, 32, 319.
- [153] Wright A.E., Griffith M.R., Burke B.F., Ekers R.D., 1994, "The Parkes-MIT-NRAO (PMN) surveys. 2: Source catalog for the southern survey (delta greater than -87.5 deg and less than -37 deg)", *ApJS*, 91, 111.
- [154] Wright A.E., Griffith M.R., Hunt A.J., Troup E., Burke B.F., Ekers R.D., 1996, "The Parkes-MIT-NRAO (PMN) Surveys. VIII. Source Catalog for the Zenith Survey (-37.0 degrees < delta < -29.0 degrees)", *ApJS*, 103, 145.