

**ADVERTIMENT.** La consulta d'aquesta tesi queda condicionada a l'acceptació de les següents condicions d'ús: La difusió d'aquesta tesi per mitjà del servei TDX ([www.tesisenxarxa.net](http://www.tesisenxarxa.net)) ha estat autoritzada pels titulars dels drets de propietat intel·lectual únicament per a usos privats emmarcats en activitats d'investigació i docència. No s'autoritza la seva reproducció amb finalitats de lucre ni la seva difusió i posada a disposició des d'un lloc aliè al servei TDX. No s'autoritza la presentació del seu contingut en una finestra o marc aliè a TDX (framing). Aquesta reserva de drets afecta tant al resum de presentació de la tesi com als seus continguts. En la utilització o cita de parts de la tesi és obligat indicar el nom de la persona autora.

**ADVERTENCIA.** La consulta de esta tesis queda condicionada a la aceptación de las siguientes condiciones de uso: La difusión de esta tesis por medio del servicio TDR ([www.tesisenred.net](http://www.tesisenred.net)) ha sido autorizada por los titulares de los derechos de propiedad intelectual únicamente para usos privados enmarcados en actividades de investigación y docencia. No se autoriza su reproducción con finalidades de lucro ni su difusión y puesta a disposición desde un sitio ajeno al servicio TDR. No se autoriza la presentación de su contenido en una ventana o marco ajeno a TDR (framing). Esta reserva de derechos afecta tanto al resumen de presentación de la tesis como a sus contenidos. En la utilización o cita de partes de la tesis es obligado indicar el nombre de la persona autora.

**WARNING.** On having consulted this thesis you're accepting the following use conditions: Spreading this thesis by the TDX ([www.tesisenxarxa.net](http://www.tesisenxarxa.net)) service has been authorized by the titular of the intellectual property rights only for private uses placed in investigation and teaching activities. Reproduction with lucrative aims is not authorized neither its spreading and availability from a site foreign to the TDX service. Introducing its content in a window or frame foreign to the TDX service is not authorized (framing). This rights affect to the presentation summary of the thesis as well as to its contents. In the using or citation of parts of the thesis it's obliged to indicate the name of the author

**Distribution and population biology of  
pelagic decapod crustaceans of the western  
Mediterranean**

*Daniela Silveira Simão*

*Dr. Pere Abelló Sala*

Barcelona, Septiembre de 2013





## Acta de qualificació de tesi doctoral

Curs acadèmic: 2012-2013

Nom i cognoms

Daniela Silveira Simão

DNI / NIE / Passaport

Y 0024865Q

Programa de doctorat

Ciències del Mar

Unitat estructural responsable del programa

Departamento de Ingeniería Hidráulica, Marítima y Ambiental  
(EHMA)

## Resolució del Tribunal

Reunit el Tribunal designat a l'efecte, el doctorand / la doctoranda exposa el tema de la seva tesi doctoral titulada

Distribution and population biology of pelagic decapod crustaceans of the western Mediterranean

Acabada la lectura i després de donar resposta a les qüestions formulades pels membres titulars del tribunal, aquest atorga la qualificació:

APTA/E     NO APTA/E

(Nom, cognoms i signatura)		(Nom, cognoms i signatura)	
President/a		Secretari/ària	
(Nom, cognoms i signatura)	(Nom, cognoms i signatura)	(Nom, cognoms i signatura)	(Nom, cognoms i signatura)
Vocal	Vocal	Vocal	Vocal

Barcelona, \_\_\_\_\_ d'/de \_\_\_\_\_ de 2013

El resultat de l'escrutini dels vots emesos pels membres titulars del tribunal, efectuat per l'Escola de Doctorat, a instància de la Comissió de Doctorat de la UPC, atorga la MENCIÓ CUM LAUDE:

SI     NO

(Nom, cognoms i signatura)	(Nom, cognoms i signatura)
Presidenta de la Comissió de Doctorat	Secretària de la Comissió de Doctorat

Barcelona, \_\_\_\_\_ d'/de \_\_\_\_\_ de 2013



# *Distribution and population biology of pelagic decapod crustaceans of the western Mediterranean*

**PhD thesis of Daniela Silveira Simão**

**Supervisor:** Dr. Pere Abelló Sala

2013 September

Distribution and population biology of pelagic decapod  
crustaceans of the western Mediterranean

Simão, Daniela Silveira

Thesis presented to obtain the PhD title  
from the Universitat Politècnica de Catalunya

124 pages  
Barcelona, 2013 July

Supervisor: Abelló, Pere

Institut de Ciències del Mar  
Consejo Superior de Investigaciones Científicas (CSIC)

Programa de doctorado en Ciencias del Mar  
Departamento de Ingeniería Hidráulica, Marítima y Ambiental (EHMA)  
Universitat Politècnica de Catalunya

UNESCO codes: 240191 Invertebrates no Insects,  
250805 Hydrobiology,  
250501 Biogeography,  
240903 Population Genetics

*Distribution and population biology of pelagic  
decapod crustaceans of the western Mediterranean*





The PhD studies of the candidate were supported by a predoctoral AECID-MAE studentship.

This thesis is based on data from various sources and projects: the Medits trawl series surveys, run by the Instituto Español de Oceanografía within the EU Data Collection frame, and the research projects PONT09 (CGL2009-12912-C03-03), IDEADOS (CTM2008-04489-C03-02/MAR), DisMarGen\_2009-FBBVA, and BENTHOMICS (CTM2010-22218-C02).

# Acknowledgements

Primeramente agradezco a mi supervisor de tesis Dr. Pere Abelló Sala por la grande oportunidad de hacer mi tesis de doctorado sobre su supervisión. Gracias por la confianza depositada en mí al permitirme trabajar los datos de las campañas MEDITS-ES que usted viene participando en todo el sector peninsular desde el principio en 1994. Gracias por el constante compromiso que tiene conmigo desde el primera vez que vine a Barcelona a conocerte y a estudiar la posibilidad de hacer un doctorado contigo hasta ahora, el final de la redacción de mi tesis de doctorado. He tenido mucha suerte en encontrar a ti Pere como supervisor pues además de tu humanidad, optimismo y sinceridad priorizas la tranquilidad y el respeto como base para una buena relación profesional. Por eso hoy me alegro en decir que estos años de trabajo fluyeron muy bien. Muchísimas gracias por todo!

A lo largo de ese trayecto sobre su supervisión tuve la oportunidad de aprender con experiencias riquísimas. La primera fue estrenarme en la MEDITS-ES como mi primera campaña oceanográfica donde tuve la oportunidad de conocer la diversidad de la fauna Mediterránea 'ao vivo e a cores' -como se dice en Brasil- y observar como la comunidad del fondo se va cambiando desde aguas costeras hasta las aguas profundas y desde la proximidad del estrecho de Gibraltar hasta el Golfo de Lion, inolvidable! Aprovecho aquí para agradecer a los jefes de campaña Luis Gil de Sola y Mariano García por permitir mi participación en el muestreo y a todos los tripulantes científicos y marineros por la buena compañía y por la oportunidad de vivenciar sus costumbres y culturas. En especial agradezco a Emilia Zas por la ayuda en los muestreos biológicos.

Al desembarcar Pere me dio la iluminada misión de llevar algunas muestras para hacer el análisis genético junto a la Dra. Marta Pascual, de la Universidad de Barcelona. Marta, soy muy agradecida de haber aprendido dar mis primeros pasos en el análisis genético y filogenético a ti contigo. A lo largo de mis largas temporadas en tu laboratorio he tenido la oportunidad de dejarme contagiar con tu ánimo en trabajar y discutir temas novedosos y espero seguir siendo invitada a participar de las charlas que organizas ahí en la UB. Gracias por todo Marta! Victor Hugo y Aymeé, muchísimas gracias por la atenciosa ayuda en compartir sus conocimientos de metodología de laboratorio para la extracción y secuenciación de ADN. Además agradezco la alegre compañía y el apoyo de los demás compañeros del laboratorio Gemma, Pedro, Victor Hugo y Rosana.

Otra campaña muy interesante que Pere me brindó fue la campaña IDEADOS donde tuve la oportunidad de trabajar con un equipo multidisciplinar formado por algunos compañeros aquí del ICM y del IEO Balear. Gracias Pilar Olivar por permitir mi participación en la y a Ainoa, Adam y Clara por la ayuda en el muestreo biológico y a todo el equipo de científicos y marineros la

campana. En especial agradezco a Asvin Pérez por toda tu ayuda y complicidad en diferentes frentes: en la ayuda con el muestreo biológico en el mar y en tierra y por la atención en compartir datos e información sobre la campana y workshop. Além disso estou feliz de estar em colaboração contigo no novo artigo, obrigada por tudo menina.

Agradezco a los investigadores: Guillermo Guerao por la atención y recomendaciones con los dibujos científicos y a Francesc Maynou por la ayuda en quitar dudas en cuestiones de estadística. Gracias Pilar Sánchez, Ana Sabatés, Conchita, Cesar García y Núria pues con el apoyo de vosotros pude mantener mi espacio de trabajo dentro del ICM así como llevar a cabo actividades fuera de él como las campanas oceanográficas.

Un especial agradecimiento a Genoveva Comas de la UPC por toda su atención y compromiso en concretar todos los trámites administrativos en tiempo hábil y con muy buena voluntad. Además agradezco al coordinador del programa de doctorado en Ciências del Mar Dr. Agustín Sánchez-Arcilla por la atención en aceptar mi participación en el programa.

Gracias a mis queridos compañeros becarios del Departamento de recursos pesqueros por la complicidad y el apoyo en el trabajo y por la divertida compañía en las comidas y cafés mientras estamos reunidos en el ICM, lo que siempre viene bien para recargar las pilas y seguir currando un poco más. No olvidaré de nuestros clásicos encuentros entre becarios del departamento de recursos pesqueros donde presentábamos nuevas herramientas de análisis de datos y discutíamos. De los Chanclas que elige estar en la categoría los 'casi que no' solo para pasárselo bien y mantener el espíritu democrático e igualitario de nuestra pandilla. De las incontables cervecitas en el eterno Bitácora, donde sé que siempre habrá algún becario del ICM colgadito allí en medio a las bufandas.

Un agradecimiento especial a las chicas del despacho B-60 por toda la amistad, cariño, alegría y acogida. Hemos compartido buenos momentos jutas, las quiero mucho. Sé que siempre tendré un lugarcito ahí en el corazón (y en el despacho) de vosotras Amalia Manjabacas, Anabel Colmenero, Vanesa Raya y Gemma Quilez, las quiero mucho! A mi querida super Quely! por tu amistad, amor, complicidad, cariño, paciencia, alegría, viajes, paseos, risas y por compartir tus momentos en familia conmigo, te adoro Raquel Saez! Mis queridas vecinas Noelia Díaz y Dafni Anastasiadi siempre con sus sonrisas gracias por la amistad, apoyo y más risas. A Samuele Tecchio y Marco Martínez gracias a mantenerse presente mismo si pasamos meses sin vernos pues cuando aparecéis me alegran el día al hacerme recordarme que puedo contar con ustedes. Entre los pasillos y actividades del ICM tuve la oportunidad de conocer personas maravillosas como las queridas Paola Castelanos, Ana Gomes y Cristina Castillo, gracias por llenarme de alegría y positividad a cada vez que nos vemos, adoro vocês e já estou com saudades!

Ah Barcelona! Adoro essa cidade. Obrigada por acolherme tão bem. Aqui conheci gente muito bacana fora do meio academico. As minhas queridas Tryna, Lucile e Ingrid obrigada pela

amizade e alegria contagiante que só as bairns têm. Agora que terminei a tese posso voltar pra zueira, a levada ta louca pra dançar. Gracias Paula Leiva y Eva Todoli por la ayuda y el cariño. E às minhas amigas estrelinhas Rita, Lidia, Fiona, Alba y Nuria gracias por luz de vosotras y por todo lo que hemos aprendido juntas.

Mas a maior gratidão que sinto é por minha familia. Obrigada papai e mamãe por abençoars-me com a vida, amor e cuidado.

Barbara minha irma! Conto contigo sempre e todo momento bonito, divertido ou interessante que vivo desejo que voce o viva também. Tipo: Humm que sorvete gostoso! Cadê a Barbara pra provar isso?! Hahahha é assim o tempo todo 😊 Quero seguir minha vida compartilhando bons momentos contigo. Te amo!

Pai, obrigada pelos exemplos que voce me deu. Cresci te observando e aprendendo com sua curiosidade e teus inventos loucos, desengonçados mas sempre muito interessantes para mim. Saiba que voce foi meu maior inspirador na escolha dessa minha profissao. Sou muito orgulhosa de ser tua filha pois admiro sua grandeza de espírito. Eres um grande amigo, honesto, bom, generoso, inteligente e sei que posso contar contigo para sempre sempre! Te amo meu pai querido.

Obrigada minha querida mae. Te amo tanto. O teu amor é a minha força. Obrigada por cuidar da minha infância e formação com teu imenso amor, atenção, conversas, não me faltou nada. Voce é minha fonte inspiração e de energia para seguir adiante sempre. Tudo valeu a pena. Obrigada por tudo o que me ensinou e por confiar em mim. Por isso hoje sinto o teu apoio em tudo o que faço e todas as minhas conquistas dedico também a ti.

A minha tia Zoly obrigada por vir me visitar aqui em Barcelona. Obrigada por estar presente na minha vida cuidando de mim com todo seu amor e alegria sem fim. Você é sempre muito bem vinda. Te amo.

A mis antepasados y todo el amor que traspasó generaciones y generaciones y que me trajo a la vida. Gracias al amor incondicional de muchos estoy aquí hoy concretando y celebrando más una etapa en mi vida. Estoy muy feliz de haber hecho estos estudios de doctorado y sé que si yo estoy feliz vosotros todos están felices sea donde estén. Y es con mucho gusto que repaso ese mismo amor a las demás generaciones que vendrán y a las personas maravillosas que están a mi lado.

Gracias a ti mi marido Dario por el grande amor que vivimos. Gracias por compartir tu vida conmigo pues a tu lado cada día es una nueva alegría. Estoy muy feliz por empezar una nueva familia contigo. Hoy digo que el amor verdadero y puro existe si, solo hay que saber cuidarlo. Ti amo.

Y finalmente gracias por la vida. A todo, a lo bueno y a lo malo. Hoy soy quien soy gracias a todas las experiencias que he vivido. Y que vengan nuevos desafíos!

## *Dedicated to*

À minha maravilhosa mãe, Elian Silveira Simão, que sempre me apoiou e acreditou em mim. Obrigada pelo seu amor incondicional que segue comigo para sempre. Alegre, perseverante, carinhosa, dedicada, amiga, compreensiva, generosa e iluminada. Te agradeço por tudo e acredito sempre que você foi e continuará sendo a melhor mãe que eu poderia ter. Te amo tanto, cada dia penso em ti e sinto cada vez mais saudade. Mas em seguida busco me acalmar ao reconhecer que sou abençoada com muitas maravilhas. E me alegro pois apesar das homeopáticas lágrimas de saudade (que cada vez são mais raras) me sinto cada vez mais forte e centrada, todo o oposto do que sentia ao receber tuas últimas notícias. Mãe tudo valeu a pena, tudo. Obrigada pela vida e pela riqueza de teus ensinamentos. Eu te amo!

*No hay diferencias entre sueño y realidad, basta vivirlo.*

Daniela Silveira Simão

# Eterno Aprendiz

## Gonzaguinha

Eu fico  
Com a pureza  
Da resposta das crianças  
É a vida, é bonita  
E é bonita...

Viver!  
E não ter a vergonha  
De ser feliz  
Cantar e cantar e cantar  
A beleza de ser  
Um eterno aprendiz...

Ah meu Deus!  
Eu sei, eu sei  
Que a vida devia ser  
Bem melhor e será  
Mas isso não impede  
Que eu repita  
É bonita, é bonita  
E é bonita...

E a vida!  
E a vida o que é?  
Diga lá, meu irmão  
Ela é a batida  
De um coração  
Ela é uma doce ilusão  
Hê! Hô!...

Mas e a vida  
Ela é maravida  
Ou é sofrimento?  
Ela é alegria  
Ou lamento?  
O que é? O que é?  
Meu irmão...

Há quem fale  
Que a vida da gente  
É um nada no mundo  
É uma gota é um tempo  
Que nem dá um segundo...

Há quem fale  
Que é um divino  
Mistério profundo  
É o sopro do criador  
Numa atitude repleta de amor...

Você diz que é luta e prazer  
Ele diz que a vida e viver  
Ela diz que melhor é morrer  
Pois amada não é  
E o verbo é sofrer...

Eu só sei que confio na moça  
E na moça eu ponho a força da fé  
Somos nós que fazemos a vida  
Como der ou puder ou quiser...

Sempre desejada  
Por mais que esteja errada  
Ninguém quer a morte  
Só saúde e sorte...

E a pergunta roda  
E a cabeça agita  
Fico com a pureza  
Da resposta das crianças  
É a vida, é bonita  
E é bonita...

Viver!  
E não ter a vergonha  
De ser feliz  
Cantar e cantar e cantar  
A beleza de ser  
Um eterno aprendiz...

Ah meu Deus!  
Eu sei, eu sei  
Que a vida devia ser  
Bem melhor e será  
Mas isso não impede  
Que eu repita  
É bonita, é bonita  
E é bonita...





# List of contents

Summary	iv
Resum	vi
i. Introduction	1
i.i. Pelagic decapod crustaceans	1
i.ii. General aim	3
i.iii. Study area	3
i.iv. Specific Goals and Chapters	4
i.v. Presentation of the species	8
i.v.i. Infraorder Caridea	8
i.v.i.i. Family Pasiphaeidae	8
<i>Pasiphaea sivado</i>	10
<i>Pasiphaea multidentata</i>	13
i.v.i.ii. Family Pandalidae	15
<i>Chlorotocus crassicornis</i>	15
<i>Plesionika heterocarpus</i>	16
i.v.ii. Suborder Dendobranchiata	16
i.v.ii.i. Family <i>Sergestidae</i>	17
<i>Sergestes arcticus</i>	17
<i>Sergestes vigilax</i>	18
<i>Sergestes arachnipodus</i>	18
<i>Sergia robusta</i>	18
i.v.ii.ii. Family Benthescymidae	19
<i>Gennadas elegans</i>	19
Chapter 1	21
Pasiphaeid shrimp populations in the western Mediterranean: geographical variability in distribution and population patterns	
1.1. Introduction	22
1.2. Material and Methods	24
1.2.1. Study area and oceanographic context	24
1.2.2. Sampling and Analysis	26
1.3. Results	29
1.3.1. <i>Pasiphaea sivado</i>	29

1.3.1.1. Occurrences	29
1.3.1.2. Density and biomass	31
1.3.1.3. Size structure and size at maturity	34
1.3.1.4. Temperature and salinity	36
1.3.2. <i>Pasiphaea multidentata</i>	37
1.3.2.1. Occurrences	37
1.3.2.2. Density and biomass	38
1.3.2.3. Size structure	39
1.3.2.4. Temperature and salinity	40
1.4. Discussion	40
Chapter 2	45
Sexual dimorphism in the benthopelagic shrimp <i>Pasiphaea sivado</i> (Crustacea: Caridea: Pasiphaeidae)	
2.1. Introduction	46
2.2. Material and Methods	47
2.2.1. Pleopod morphology	47
2.2.2. Gender differences in size and sex ratio	49
2.3. Results	50
2.3.1. Pleopod morphology	50
2.3.2. Comparison of sizes and size-related sex-ratio among geographic areas	52
2.3.3. Size structure with depth by geographic area	53
2.3.4. Sex-ratio with depth	56
2.4. Discussion	56
2.4.1. Pleopod morphology	56
2.4.2. Size dimorphism and sex ratio	57
Chapter 3	61
Population genetics and phylogeny of pasiphaeid shrimps from the western Mediterranean Sea	
3.1. Introduction	62
3.2. Material and Methods	64
3.2.1. Sampling	64
3.2.2. DNA isolation, amplification and sequencing	65
3.2.3. Intraspecific genetic diversity	66
3.2.4. Phylogenetic analysis	67

3.3. Results	69
3.3.1. Population genetic structure	69
3.3.2. Phylogeny of <i>Pasiphaea</i>	72
3.4. Discussion	74
3.4.1. Population genetics	74
3.4.2. Phylogeny	79
Chapter 4	73
Vertical and seasonal distribution of pelagic decapod crustaceans over the shelf-break and middle slope in two contrasting zones around Mallorca (western Mediterranean Sea)	
4.1. Introduction	83
4.2. Material and Methods	84
4.2.1. Study area and hydrographic setting	84
4.2.2. Sampling	85
4.2.3. Data analysis	88
4.3. Results	89
4.3.1. Funistics	89
4.3.2. Day/night depth distribution	91
4.3.3. Pelagic trawl samples	92
4.3.4. IKMT samples	96
4.3.5. Population size structure	98
4.4. Discussion	101
4.4.1. Study area	101
4.4.2. Species and faunistic assemblages	103
4.4.3. Size structure	105
4.4.4. Colouration	106
Conclusions	110
References	115

# Summary

The present thesis provides new information about pelagic decapod crustaceans, specially so concerning the Mediterranean pasiphaeids. Remarkable differences in the pattern of distributions of the pasiphaeids were observed according to the influence of Atlantic waters. Overall both abundance and biomass of the pasiphaeids in Alboran Sea was higher than in the rest of the Iberian Mediterranean populations. The depth on which the pasiphaeids started to occur in appreciable density was also shallower in the Alboran Sea where the *P. sivado* presented the widest depth range of distribution as well as the biggest size of first maturity of females. This size was progressively smaller with increasing distance from the Atlantic water source, i.e the Gibraltar Strait area. Overall the two species presented a size structure that changed with depth, with younger individuals occupying shallower depths and biggest adults occupying deeper bathymetries. *P. multidentata* juveniles were only observed in the Catalan Sea while they were absent in the Alboran Sea.

Size and morphological dimorphism of *P. sivado* were reported for the first time in the present thesis. All the five pleopods of females were thinner and more elongated than male pleopods, which in turn were rounded anteriorly and had robust shapes. A critical size from which the population sex ratio biases changed from females to males dominance was found in all *P. sivado* populations of the Mediterranean Iberian coast. This critical size was different and progressively larger again from populations nearest the Gibraltar Strait versus those placed in the Catalan Sea, in concordance with the female first maturity size gradient.

The present thesis also provides a first attempt to investigate the influence of the Gibraltar Strait on the genetic population structure of the species by using congener benthopelagic shrimps that dwell in different depth strata. The Gibraltar Strait was shown to be a strong and unique geographical barrier to the genetic flux of both *P. sivado* and *P. multidentata*, due the presence of a marked genetic structure characterized by two main haplotypes: one Atlantic and another one Mediterranean. Other new information related to the phylogeny of the genus *Pasiphaea* was that *P. sivado* a much higher genetic divergence when compared to the rest of pasiphaeid species analysed, indicating that several genetic subgroups do occur within the genus *Pasiphaea*.

The study of the pelagic decapod community in the waters around the island of Mallorca, placed between the Algerian and the Catalano-Balearic basins, showed the occurrence of two main assemblages: one above the shelf break and another one above the middle slope. The shelf community was characterized by the dominance of almost transparent species, such as *Sergestes arcticus*, the species that reached the greatest abundances, and *P. sivado* that was omnipresent in all pelagic strata. The novel registration of *Chlorotocus crassicornis* and *Plesionika heterocarpus* in the

pelagic realm were restricted to the shelf-break at the Deep Scattering Layer. The red body species *Gennadas elegans*, *Pasiphaea multidentata*, and *Sergia robusta*, were restricted to the middle slope. No significant differences concerning the sampling geographic area or seasonality were found. Actually, the decapod pelagic community of the Balearic Sea was mainly structured by both the geomorphology (and associated hydrographic characteristics of the shelf/slope transition) and the influence of light in the water column. No decapod crustacean species occurred in epipelagic daytime samples. Size analysis showed the occurrence of species-specific patterns concerning the size/age movements into the water column throughout the day-night cycle. Moreover, the present work clearly confirms the presence of adult *P. multidentata* in the water column, evidencing that the adult fraction of the population has also the ability to perform vertical migrations.

In the present days many knowledge gaps still exist concerning the autoecology of pasiphaeids and several pelagic decapods. Many questions about population structure, distribution and phylogeny could be better endorsed by knowing the early life and dispersal histories of these species. The high sampling effort, and their associated high cost, still remains one of the principal reasons for the scarce information available nowadays.

Keywords:

Macrozooplankton, Micronekton, benthopelagic decapods, Necktobentonic crustaceans, Diel vertical migrations, Size structure, Sexual dimorphism, Population genetics, Phylogenetics, Pasiphaea, Sergestidae.

# Resum

Aquesta tesi presenta nova informació sobre crustacis decàpodes pelàgics, principalment sobre els pasifèids presents a la Mediterrània: *Pasiphaea sivado* i *Pasiphaea multidentata*. S'han trobat diferències importants en les pautes de distribució dels pasifèids en relació a la influència de les aigües atlàntiques a la Mediterrània. Globalment, tant l'abundància com la biomassa dels pasifèids del Mar d'Alboran van ser superiors que a la resta de poblacions mediterrànies mostrejades. La profunditat a la qual es comencen a trobar les diferents espècies de pasifèids de la Mediterrània és també menor a la Mar d'Alboran, on *P. sivado* va presentar el major rang de distribució en fondària, així com també la més gran talla de maduresa en femelles. Aquesta talla es va fent progressivament menor a mesura que augmenta la distància a la font d'aigües atlàntiques, és a dir, de la zona de l'Estret de Gibraltar. Ambdues espècies van presentar una estructura de talles variable en funció de la profunditat, amb individus juvenils ocupant profunditats més somes que els adults. Addicionalment, juvenils de *P. multidentata* només es van observar al Mar Català, mentre que van ser absents del Mar d'Alboran, fet que suggereix l'existència de diferències important en l'estacionalitat de la reproducció i/o de la dinàmica de les poblacions.

L'existència de dimorfisme sexual entre mascles i femelles de *P. sivado* es reporta per primera vegada en aquesta tesi, així com de diferències significatives en l'estructura de talles entre sexes. Els cinc pleopodis de les femelles adultes són clarament més fins i allargats que els pleopodis dels mascles adults, els quals són arrodonits anteriorment i tenen una forma més robusta. S'ha detectat una talla crítica a partir de la qual la proporció de sexes de les poblacions canvien cap a una dominància significativa dels mascles. Aquesta talla disminueix progressivament i significativa des de la població més propera a l'Estret de Gibraltar, la del Mar d'Alboran, passant per la població d'Alacant, fins a la població del Mar Català, en concordança amb els canvis observats a la talla de maduresa de les femelles.

Aquesta tesi també presenta una primera aproximació a investigar el paper de l'Estret de Gibraltar sobre l'estructura genètica de les poblacions de pasifèids, utilitzant les dues espècies utilitzades en els capítols precedents, les quals representen espècies congenèriques, simpàtriques geogràficament, però amb una marcada distribució batimètrica diferencial. Les anàlisis efectuades han mostrat que l'Estret de Gibraltar constitueix una forta i única barrera geogràfica per a la connectivitat genètica entre les poblacions d'ambdues espècies situades a una banda i altra de l'Estret. En ambdues espècies s'ha detectat l'existència d'una marcada estructuració genètica caracteritzada per dos haplotips principals, un que es troba només en poblacions atlàntiques, i un altre únicament en poblacions mediterrànies. Basant-se també en característiques genètiques, s'ha

obtingut nova informació referent a la filogènia del gènere *Pasiphaea*, que mostra l'existència d'una important divergència genètica entre *P. sivado* i la resta d'espècies de la família *Pasiphaeidae* de les quals en l'actualitat es té informació, fet que indica que dins del gènere *Pasiphaea* es troben diferents subgrups genètics ben delimitats.

L'estudi de la comunitat de crustacis decàpodes pelàgics en aigües al voltant de l'illa de Mallorca, localitzada entre les conques Algeriana i Catalano-Balear ha mostrat la presència de dues comunitats faunístiques diferenciades, no geogràficament sinó en funció de la batimetria: una en aigües localitzades sobre el límit plataforma-talús i una altra al damunt del talús mig. La comunitat més soma, la del límit plataforma-talús, es va caracteritzar per la dominància d'espècies pràcticament transparents, com *Sergestes arcticus*, l'espècie que va assolir les abundàncies més elevades, i *Pasiphaea sivado*, espècie omnipresent a tots els nivells pelàgics mostrejats. Els registres a l'ambient pelàgic de *Chlorotocus crassicornis* i *Plesionika heterocarpus*, no citades anteriorment en aquest ambient, es van restringir a les capes de difusió profunda del límit plataforma-talús. Les espècies de coloració més vermellosa (*Gennadas elegans*, *Pasiphaea multidentata* i *Sergia robusta*) es van localitzar únicament sobre fons del talús mig. No es van trobar diferències significatives en funció de la localització geogràfica o de les dues estacions de l'any mostrejades. Es considera que la comunitat de crustacis decàpodes pelàgics del Mar Balear està fonamentalment estructurada tant per la geomorfologia i característiques hidrogràfiques associades a la transició entre la plataforma i el talús continentals, com per la influència de la llum a la columna d'aigua. No es van trobar espècies de crustacis decàpodes en mostres epipelàgiques preses durant el dia. L'anàlisi de les talles dels individus capturats ha mostrat que els moviments de les diferents classes de talla (edat) a través de la columna d'aigua varien al llarg del cicle dia-nit d'una manera diferencial en funció de cada espècie. Adicionalment, aquest treball confirma clarament que la fracció adulta de la població de *Pasiphaea multidentata* presenta també la capacitat de dur a terme migracions verticals.

S'és conscient que existeixen avui dia encara molts forats en el coneixement de l'autoecologia i cicle vital tant dels pasifèids com de la resta de decàpodes pelàgics. Moltes qüestions romanents per a comprendre millor l'estructura i dinàmica de poblacions, distribució i filogènia, podrien ser millor enteses i discutides si hagués més coneixença sobre el cicle vital, els estadis larvaris i post-larvaris, així com sobre la dispersió i connectivitat de les poblacions. L'elevat esforç de mostreig, i el seu alt cost associat, són encara avui dia una de les raons principals de l'escassetat d'informació disponible.

Paraules clau:



Macrozooplàncton, Micronècton, decàpodes bentopelàgics, crustacis nectobèntics, migracions diàries verticals, estructura de talles, dimorfisme sexual, genètica de poblacions, filogènia, Pasiphaea, Sergestidae.





# Introduction





# Introduction

## i.i. Pelagic decapod crustaceans

Pelagic decapod shrimps have a widespread geographic and bathymetric distribution from intertropical waters to high latitudes in both hemispheres (Casanova and Judkins 1977, Fasham and Foxton 1979, Clarke and Holmes 1987, Burghart *et al.* 2007). They comprise an important biomass fraction of the faunistic communities of the shelf break and slope (Long *et al.* 1995, Sumida and Pires-Vanin 1997, Abelló *et al.* 2002, Company *et al.* 2004, Serejo *et al.* 2007, Callaway *et al.* 2007, Escobar-Briones *et al.* 2008, Delgado *et al.* 2013), which are continental margin areas of intense and increasing exploitation by the trawl fishery around the world (Orensanz *et al.* 1998, Callaway *et al.* 2007). These shrimps are relatively small and usually have no commercial interest. However, benthopelagic shrimps are very frequent in fishery hauls (Abelló *et al.* 2002), but, with some exceptions (e.g. *Pasiphaea japonica*) where they have become the targets of a specific fishery (Nanjo and Ohtomi 2009) or when they are sold as an accompanying by-catch in non-industrial fisheries (Ramirez-Llodra *et al.* 2007), virtually all their captures are discarded and consequently they are not landed nor sold. Notwithstanding they are a substantial, when not fundamental, part of epibenthic food chains on the continental slope (Rosecchi *et al.* 1988, Hassani, *et al.* 1997, Polunin *et al.* 2001, Serrano *et al.* 2003, Layman *et al.* 2005, Fanelli and Cartes 2008, Battaglia *et al.* 2013).

An inherent characteristic of the benthopelagic decapods inhabiting the slope is their marked behavioural rhythms. During the day hours these shrimps dwell on or close to the sea floor, in the so called *Benthic Boundary Layer* (BBL), whereas during the night they ascend in the water column to feed in surface or sub-surface layers of the ocean (Aguzzi *et al.* 2007). From the energetic point of view these shrimps act as exporters of primary production from the superficial photic zones to deep sea aphotic water masses (Maynou and Cartes 1998).

In oligotrophic zones, the primary production of the organisms that dwell near bottom areas of the slope and bathyal zones depends, almost exclusively, on the epipelagic planktonic production

of the photic zone (Eppley and Peterson 1979, Pauly and Christensen 1995, Cartes and Maynou 1998, Cartes *et al.* 2001, Cartes *et al.* 2002, Cusson and Bourget 2005, Tecchio *et al.* 2013). In this light, the role of the vertical migrators such as pelagic shrimps is the key for the functioning of the slope and bathyal ecosystems (Cartes and Maynou 1998). There are some families of pelagic decapods, with cosmopolitan distribution, such as the Pasiphaeidae or Sergestidae that are well adapted to inhabit the deep sea as well as to perform vertical migrations (Cartes *et al.* 1993, Cartes 1993, Aguzzi *et al.* 2007).

Trophic relationship studies based on the stomach content analysis in fishes (Rosecchi *et al.* 1988, Hassani *et al.* 1997, Polunin *et al.* 2001, Serrano *et al.* 2003, Layman *et al.* 2005) showed that pasiphaeid shrimps are preyed upon by a variety of nektonic organisms with both commercial and ecological interest (Rosecchi *et al.* 1988, Orsi-Relini and Relini 1990, Cartes 1994, Hassani *et al.* 1997, Garison and Link 2000, Polunin *et al.* 2001, Serrano *et al.* 2003, Layman *et al.* 2005, Fanelli and Cartes 2008, Tecchio *et al.* 2013, Battaglia *et al.* 2013). The high frequency of occurrence and abundance associated to the vertical migratory behaviour in this family (Aguzzi *et al.* 2007) has demonstrated their key role in the trophic ecology of slope and demersal environments (Skjoldal and Bamstedt 1977, Maynou and Cartes 1998, Polunin *et al.* 2001, Cartes *et al.* 2007a, Tecchio *et al.* 2013).

In the Mediterranean Sea the Pasiphaeidae and Sergestidae are the most common families of pelagic decapods. In special, the congeneric pasiphaeid species *Pasiphaea sivado* and *Pasiphaea multidentata*, together with the sergestid *Sergestes arcticus*, are the most frequent and abundant benthopelagic decapod shrimps along the western Mediterranean coasts (Abelló *et al.* 1988, 2002, García-Rodríguez *et al.* 2011).

## **i.ii. General aim**

The main aim of the present thesis is to provide new knowledge on pelagic decapod crustaceans of the western Mediterranean continental slope ecosystem, focusing on distribution characteristics and their association with geomorphology and water masses, population biology, genetics and benthopelagic behaviour. Specific goals are detailed below and in the presentation of each chapter. The main target species of the study are those belonging to the caridean shrimp family Pasiphaeidae, particularly *Pasiphaea sivado* and *Pasiphaea multidentata* since they are the most common and abundant benthopelagic species in the study area when sampled with epibenthic and demersal gears.

Since scientific knowledge constitutes the basis for fisheries and conservation management, it is expected that the information provided will be useful both as an increase in basic knowledge and as a contribution for the sustainable management of the exploited ecosystems of the continental slope. The cosmopolitanism of the crustacean families focus of this study implies that a specific approach in a particular pelagic/benthopelagic ecosystem can be widely extrapolated to similar ecosystems functioning in geographically distant areas. In this light, the description of the ecology of pelagic decapods provides also a basis for the knowledge of the dynamics, functioning and interrelationships between the pelagic and deep sea systems.

## **i.iii. Study area**

The overall study area encompassed the western Mediterranean Sea along the coasts of the Iberian Peninsula from the Gibraltar Strait in the SW to Cape Creus in the NE, including the Balearic Islands. For the genetic studies, the study area was enlarged as to comprise the Gulf of Cadiz, west of Gibraltar, in the NE Atlantic. The semi-arid climatology of the Mediterranean Sea area is characterised by an excess of evaporation in relation to pluviometry and river input. This disequilibrium, together with the seasonal increase of sea surface temperatures in summer, confers to these semi-closed



Mediterranean Sea high values of both salinity and density. High evaporation rates due to strong, extremely dry, cold, northerly winds from the European subcontinent also contribute to the salinity increase, especially so in

winter. The Gibraltar Strait is the most important entrance of the Mediterranean Sea with receives large amounts of Atlantic surface water (in an order of  $106 \text{ m}^3 \text{ s}^{-1}$ ) to compensate the water loss due to evaporation. The western Mediterranean Sea is composed by two



Figure 1 : Map of the Mediterranean iberic Sea showing the morphology of the hidrographic basins and the conection with the Atlantic ocean.

geomorphologic basins: The Algerian and the Balearic (or

Catalano-Balearic) basins. The Algerian basin is located east of the Gibraltar Strait, becoming the first Mediterranean basin that receives Atlantic water. Density differences imply that Atlantic water enters the Mediterranean Sea through the surface water layer towards the east, while denser Mediterranean water leaves this sea through deep layers in smaller volumes. Incoming Atlantic water is slowly altered during their trajectory by mixing with resident Mediterranean water masses and small river input, and also by evaporation (Tizperman and Malanotte-Rizzoli 1991, Cacho *et al.* 1999, Millot and Taupier-Letage 2005) (Figure 1). Details about the oceanography of the area will be presented in the next chapters as specific information will be required.

## i.iv. Specific Goals and chapters

The present thesis is structured in four main chapters, in agreement with four main specific goals.

i.iv.i. Chapter 1:

*Pasiphaeid shrimp populations in the western Mediterranean: geographical variability in distribution and population patterns*

---

The aim of this first chapter is to analyse the geographic and bathymetric distribution of the congeneric species *P. sivado* and *P. multidentata* along the Mediterranean coasts of the Iberian Peninsula in terms of both density and biomass, to determine some population characteristics, namely size structure and size at maturity along 10 geographic subareas, as well as to analyse the relationships between abundance and hydrological characteristics, namely bottom temperature and salinity. The study is based on an extensive data set of 15 annual fishery research cruises (1994-2008) along the study area. The specific objectives that guided this chapter were:

- To determine the depth range and preferential depth of occurrence and abundance (density and biomass) of both species along the Iberian Mediterranean coast (10 geographic/geomorphologic subareas) and along the sampled depth gradient (40-800 m).
- To analyse the population size structure of the species along the Iberian Mediterranean coast and the sampled depth gradient.
- To assess the variability in size of maturity of *P. sivado* along the Iberian Mediterranean coast.
- To analyse the relationships between abundance and hydrological characteristics, namely bottom temperature and salinity, taking into account the ontogeny of the species (juveniles vs. adults).

i.iv.ii. Chapter 2:

## Sexual dimorphism in the benthopelagic shrimp *Pasiphaea sivado* (Crustacea: Caridea: Pasiphaeidae)

Indications of the existence of sexual dimorphism in pleopod shape were observed in *P. sivado* during field biological data acquisition. Accordingly, pleopod shape was tested using morphological and geometrical morphometrics techniques. Gender differences in sex size structure were also tested to verify the occurrence of additional sexual dimorphism characteristics. The main objectives of this chapter were:

6

- To verify whether the pleopod protopod overall shape, as well as its length-width ratio, varies between genders in adult individuals.
- To assess whether size structure varies between genders, globally, geographically and bathymetrically.
- To verify whether the sex ratio varies as a function of size, depth and geographic sector.

### i.iv.iii. Chapter 3:

## Population genetics and phylogeny of pasiphaeid shrimps from the western Mediterranean Sea

This chapter is a first attempt to investigate the influence of the Gibraltar Strait on the genetic population structure of both *P. sivado* and *P. multidentata*. Additionally, the chapter also aims at assessing the internal phylogeny of the Pasiphaeidae using own data and all the available sequences of pasiphaeid species in Genbank database. The goals intended were:

- To characterize the genetic structure of pasiphaeid species populations (*P. sivado* and *P. multidentata*) across the influence of the Atlantic-Mediterranean transition.

- To verify whether there is a genetic gradient within the Mediterranean in the target species as a function of the Atlantic water influence in this sea.
- To quantify the genetic distance within the Pasiphaeidae.
- To provide new insights on the phylogenetic relationships within the Pasiphaeidae.

i.iv.iv. Chapter 4:

*Vertical and seasonal distribution of pelagic decapod crustaceans over the shelf-break and middle slope in two contrasting zones around Mallorca (western Mediterranean Sea)*

The aim of this chapter was to characterize the pelagic/benthopelagic decapod community, as well as to describe their diel vertical movements and determine their size distribution across the pelagic environment in two extreme photoperiod seasons in two areas around Mallorca characterized by different oligotrophic regimes as well as water mass characteristics and dynamics. The main objectives that conducted this chapter were:

- To characterize the faunistic composition of the pelagic decapod species according to the different sampling cells (SCs): season (late autumn / summer), geographic sector (Cabrera / Sóller), bathymetry (shelf edge / slope) and day time (day / night).
- To quantify the abundance of pelagic decapod species in the different SCs, and therefore estimate their role in the benthopelagic environment.
- To determine the depth range of each pelagic decapod species.
- To verify whether and how the depth distribution of each species varies along the day time.

- To assess the occurrence of differential species assemblages as a function of the SCs.
- To estimate the diversity indexes of the pelagic decapod assemblages identified.
- To describe the size structure of the pelagic decapods according to depth and the significant assemblages identified.

## ***i.v. Presentation of the main benthopelagic species target of this thesis***

### ***i.v.i. Infraorder Caridea***

#### ***i.v.i.i. Family Pasiphaeidae***

The Pasiphaeidae family (Dana 1852) (Pleocyemata, Caridea) is distributed along a wide latitudinal range, from tropical to arctic and subantarctic waters, mainly on and above continental slopes and in the open sea, including deep-sea habitats (Clarke and Holmes 1987, Kaartvedt *et al.* 1988, Gibbons *et al.* 1994, d'Udekem & d'Acoz 1999, Tavares and Cardoso 2006). This family is represented by seven genera and 98 species: *Alainopasiphaea* (Hayashi 1999) (2), *Eupasiphae* (Wood-Mason 1893) (4), *Glyphus* (Filhol 1884) (1), *Leptochela* (Stimpson 1860)(15), *Parapasiphae* (Smith 1884)(4), *Pasiphaea* (Savigny 1816) (67) and *Psathyrocaris* (Wood-Mason 1893)(5) (De Grave and Fransen 2011, Komai *et al.* 2012).

Morphological features in this family, such as lateral compression of the body and pleopods as well as reduced rostrum, are considered to be the result of selective pressures on adaptation to the pelagic environment (Cartes *et al.* 1993, Johnsen 2005, Sardà *et al.* 2005, Aguzzi *et al.* 2007). These body characteristics confer to these shrimps the ability to maintain their position in the pelagic

environment as well as to migrate promoting fast and wide jumps horizontally above the substrate or even vertically in relation to the sea bed.

Notwithstanding the large amount of taxonomic and faunistics work on this family (e.g. Stephensen 1923, Zariquiey Álvarez 1957, Iwasaki 1990, Hayashi 1999, Tavares and Cardoso 2006) there are still few detailed studies on biological aspects of species of the family, and most of them refer mainly to the European species *P. sivado* and *P. multidentata* (Cartes 1993, Company and Sardá 2000, Frank 2000, Company *et al.* 2001, Ramirez-Llodra *et al.* 2007), but also to Pacific Ocean species such as *Pasiphaea japonica* and *Leptochela gracilis* (Nanjo 2007, Oh *et al.* 2006). In the literature, the majority of the studies on Pasiphaeidae shrimps concern information on the distribution and abundance in relatively restrict geographic and bathymetric scales, almost always in the context of deep benthic communities analysis (Cartes *et al.* 1994, Maynou *et al.* 1996, Koukouras *et al.* 2000, Abelló *et al.* 2002, Serejo *et al.* 2007). Overall, information on the biology of the Mediterranean pasiphaeid species is not scarce (Casanova 1976, Cartes 1993, Maynou & Cartes 1998, Frank *et al.* 2000, Company & Sardá 2000, Company *et al.* 2001, Ramirez-Llodra *et al.* 2007, Aguzzi *et al.* 2007). A specific work on the biologies of *Pasiphaea sivado* and *P. multidentata* in the northwestern Mediterranean was published by Company *et al.* (2001) (Table 1).

Table 1: Summary of aspects related to the life history of *Pasiphaea sivado* and *P. multidentata*. The minimum body size (Min.), maximum (Max.) and asymptotic length ( $L_{\infty}$ ) are referred to cephalothorax length in millimetres. Adapted from Company and Sardá (2000) and Company *et al.* (2001).

Population biology												
	Sex ratio fits 1:1	Sex	Min	Max	$L_{\infty}$	K	$\phi$	age ( $L_{\infty}$ )	smallest ovigerous female	Reproductive period	Recruitment	No. eggs by Females ( $\pm$ SD)
<i>P. sivado</i>	no	female	10	23.2	26.0	0.55	0.057	2	14.7	February and October	late spring	49
		Male	10	24.3	27.5	0.62	0.126	2	( $\pm$ 21.5)			
		All	7	24.3	29.5	0.55	0.162					
<i>P. multidentata</i>	no	female	21	46.7	48.5	0.85	0.833	3 to 4	29.9	February and November	late spring	12
		Male	19	42.7	44.4	0.77	0.807	3 to 4	( $\pm$ 4.5)			
		All	6	46.7	50.0	0.80	0.687					

Among all the species studied in the present thesis the pasiphaeid species show the largest mean body length ( $CL_{\min-\max} = 7 - 24.3$  mm for *P. sivado* and 6.5 - 46.7 mm for *P. multidentata*) which

in addition to their high abundance provide a very high biomass. In the Japan Sea, *Pasiphaea japonica* is a key commercial species in Toyama Bay. Recently the annual catches of *P. japonica* in this bay have increased from 400-500 t annually up to an average of 600-700 t which consequently called for implementation of appropriate stock management strategies in order to ensure the sustainability of the fishery (Nanjo and Ohtomi 2009). Nevertheless, in the western Mediterranean both *P. sivado* and *P. multidentata* are very common and abundant species, particularly *P. sivado*, but hardly no economic interest nor policy rules exist concerning these species.

### *Pasiphaea sivado* (Risso, 1816)

Also known as glass shrimp, this species has an overall transparent body with small red spots or aggregation of little spots restricted to a few edge areas of the extremities of the somites and pereopods. The species performs diel vertical migrations, inhabiting near the bottom during the day and moving upwards at night to near surface layers (Stephensen 1923, Franqueville 1971) (Figure 2). *P. sivado* is commonly caught by demersal trawling during the day on the upper slope along the Mediterranean Sea and northeast Atlantic from Morocco to Norway (Sund 1913, Stephensen 1923, Maurin 1961, González-Gurriarán and Olaso 1987; Abelló *et al.* 2002). It is also a very common species when sampled by pelagic/micronekton trawl surveys (Franqueville 1971, Koukouras *et al.* 2000).

*P. sivado* is a typical temperate species (Udekem d'Acoz 1999). In the Mediterranean the species inhabits both the eastern and western basins (Stephensen 1923). Environmental variations due to seasonality appear to have no large effect in the distribution structure and abundance of the species in Atlantic and Mediterranean populations at around 40°N of latitude (Fariña *et al.* 1997, Maynou *et al.* 1996). *P. sivado* is often the dominant species of bottom trawl surveys rather than other species target of the fisheries, including fishes, molluscs, and other crustaceans. In south Portugal the species occurred in so high densities that was classified as the typical species of the slope zone (Fariña *et al.* 1997), being also the dominant species in the Gulf of Cadiz (Delgado *et al.* 2013). In the Mediterranean it was the dominant species in the Spanish coast between 300-400 m strata

(524 ind.h<sup>-1</sup>) (Abelló *et al.* 1988, 2002) and in the central-western coast of Italy at 300-500 m (Fanelli *et al.* 2007).

The widespread geographic and bathymetric distributions, the daily vertical migration pattern together with their high abundance provides to *P. sivado* a very important role in the flux of matter and energy between the epipelagic productive zone and the demersal aphotic zone. Stomach content studies on Atlantic specimens have shown that *P. sivado* preys on euphausiids and calanoid copepods (Lagardère 1972). In the Catalan Sea the feeding activity of *P. sivado* is more intense during the night when suprabenthic crustaceans such as mysids and amphipods are the main food items. During the day mysids remain one of the most important food items together with mesopelagic fishes (Cartes 1993). Pasiphaeid shrimps are preyed upon by several crustaceans and pelagic species including fishes, cephalopods, dolphins and even fin whales (Rosecchi *et al.* 1988, Orsi-Relini and Relini 1990, Cartes 1994, Hassani *et al.* 1997, Lorentsen *et al.* 1998, Serrano *et al.* 2003, Fanelli and Cartes 2008, Battaglia *et al.* 2013).



Figure 2: A fresh sampled *Pasiphaea sivado* female bearing eggs in the abdomen. Photography by Pere Abelló.

In the Japan Sea the sister species *P. japonica* feeds mainly on mysids and euphausiids, but also on other crustaceans, fishes and cephalopods. Euphausiids were preyed in higher amounts during the summer, following their seasonal abundance bloom in Toyama Bay. These euphausiids were the exclusive prey item of the youngest size class of *P. japonica* (15 mm CL). Cephalopods occurred in foregut contents only in July and November, being preyed mainly by small individuals (16 mm CL) (Nanjo 2007).

Overall, the majority of the published data about *P. sivado* species reports their presence in the pelagic and epibenthic communities. However, due to its relevant ecological importance, *P. sivado* biology and life history has been often studied in the western Mediterranean. In the Catalan



Sea *P. sivado* sampled near the bottom has been reported to range from 7 to 24 mm of cephalothorax length (CL), its longevity was estimated to be around 2 years, the smallest ovigerous female sampled had 14.7 mm CL, and their reproduction is continuous throughout the year, although peaking in autumn-winter (Company and Sardà 2000, Company *et al.* 2001, see table 1). *P. sivado* females of the Ligurian Sea also reproduce continuously, however females parasitized by *Amallocystis* (Protozoa: Ellobiopsidae) were never seen reproducing (Orsi-Relini and Pinca 1990).

In the Irish Sea the species shows two reproductive peaks. According to Williamson (1960) final stage embryos occurred from June to December with two peaks, in July and November. In the same work the author reported for the first time the morphology of embryos, four zoeal stages and megalopa or post-larvae. The distribution of larvae and early juveniles remains unknown, except for the finding of about 20 post-larvae close to the bottom at around 100 m in the Irish Sea (Williamson 1960).

Within the genus *Pasiphaea*, *P. sivado* is the main representative of a group of nine pasiphaeid species that is distinguished from the rest of pasiphaeids by having a reduced branchial formulae (Hayashi 1999, 2006). The *P. sivado* group is composed by those species having five pleurobranchiae (*P. sivado* (Risso, 1816), *P. propinqua* De Man, 1916, and *P. japonica* Omori, 1976) while the remaining group is composed by species that present four pleurobranchiae (*P. debitusae* Hayashi, 1999, *P. fragilis* Hayashi, 1999, *P. gracilis* Hayashi, 1999, *P. laevis* Hayashi, 1999, *P. marisrubri* Iwasaki, 1989, *P. philippinensis* Hayashi, 1999, and *P. mclaughlinae* Hayashi, 2006).

The report of morphological differences between Atlantic and Mediterranean Sea populations of pasiphaeids was firstly reported by Sivertsen and Holthuis (1956) and Zariquiey Álvarez (1957). In 1976, Omori related that Mediterranean specimens of *P. sivado* presented a higher number of spinules on the merus of the first and second pereopods when compared to Atlantic specimens.

*Pasiphaea multidentata* Esmark, 1866

*Pasiphaea multidentata* is a cold temperate species (Udekem d'Acoz 1999) that inhabits benthic boundary layers in the middle and lower slope down to 2000 m depth (Cartes 1993a; Abelló *et al.* 2002; Tecchio *et al.* 2011) (Figure 3). The species also performs vertical migrations into upper water layers up to 50-200 m of the water column at night (Franqueville 1971), although vertical displacements have been apparently restricted to juvenile individuals (Aguzzi *et al.* 2007, Cartes 1993c) (see also Chapter 4). The species has a trans-oceanic distribution comprising northern latitudes of both the eastern and western north Atlantic from the Gulf of Maine (69° W) to the west coast of Europe (18° E) and latitudinally from Tromsø (69° N), Norwegian Sea, down to Mauritania (20° N), including the Mediterranean Sea (Udekem d'Acoz 1999). Concerning the bathymetric range of the species in the benthic environment, the upper limit of distribution overlaps with that of *P. sivado*. Younger individuals of *P. multidentata* with about 10-20 mm CL inhabit shallower waters (together with *P. sivado*) than adults, which can reach about 47 mm CL (Company *et al.* 2001).

In the Mediterranean Catalan Sea the reproductive cycle of *P. multidentata* is markedly seasonal (Company *et al.* 2001). However differences concerning the reproductive period and seasonality of the species have been reported along their geographical distribution. In the gulf of Maine the spawning period takes place in two seasons: spring and autumn (Apollonio 1969), while in the western coast of Norway ovigerous females occur all year round (Matthews and Pinnoi 1973). In the Catalan Sea the size structure of *P. multidentata* ranges from juveniles with about 6 mm CL to adults with about 47 mm CL (Company *et al.* 2001). Population dynamics estimations showed that *P. multidentata* can live as long as 3.5 years (Table 1) (Company *et al.* 2001; Ramirez-Llodra *et al.* 2007). The species present a very marked seasonal reproduction with ovigerous females being only present from September to February. A well defined recruitment period takes place in early spring, showing a clearly different pattern to that of the shallower congeneric species *P. sivado* that reproduces continuously in the same area. Different reproduction periodicity has been reported for both species depending on the geographic study area (Stephensen 1923, Williamson 1960, Apollonio 1969, Matthews and Pinnoi 1973).

Feeding habits studies show that in the Gulf of Maine, northwest Atlantic, the main food items of *P. multidentata* are based on pelagic copepods and euphausiids (Apollonio 1969). In the Gulf of Gascogne, northeastern Atlantic, the diet of *P. multidentata* is based on mesopelagic fishes and juvenile cephalopods, decapods such as *S. arcticus* and euphausiids; no predatory activity was found during the day (Lagardère 1977). In the Catalan Sea *P. multidentata* specimens sampled near the bottom, preyed mainly on mesopelagic fishes, euphausiids and isopods (Cartes 1993). Nocturnal feeding activity was just observed in large individuals (>28 mm CL) that stayed near the bottom during the night time, which fed on suprabenthic organisms such as *Gennadas elegans*, *P. sivado*, *P. multidentata*, sergestid shrimps and little mesopelagic fish as myctophids and *Cyclothone* spp., while individuals sampled during the day evidenced a foraging on pelagic hyperiids, fishes, euphausiids and chaetognaths ingested the night before (Cartes 1993). Different

dietary composition was also found by the same author depending on the sampled depth. Thus, on the middle slope, *P. multidentata*, *P. sivado* and *S. arcticus* were the main decapod prey items of *P. multidentata*, while in the lower slope *Gennadas elegans* and *Sergia robusta* were the most common decapod prey of *P. multidentata*. *P. sivado* was the second most important food item of *P. multidentata* individuals captured between 380-540 m (Cartes 1993), with correspond to the preferential occurrence depth of this prey (Abelló *et al.* 2002). Overall, low dietary overlap was observed among *P. sivado* and *P. multidentata* by Cartes (1993), who supposed that it could be due the different preferential size spectra of the food items preyed by the these pasiphaeids. In this sense,



Figure 3: (a) *Pasiphaea multidentata* individuals sampled by a micronekton trawl. One adult individual is placed vertically while juvenile individuals are placed horizontally. (b) a typical adult captured by a demersal trawl.

Lagardère (1977) stated that the different depth range among the same pasiphaeids and other sergestid species prevents any trophic competition between them.

Morphological divergences between western Mediterranean and Atlantic (North Sea) populations of *Pasiphaea multidentata* were described by Zariquiey Alvarez (1957), mainly concerning the postrostral spine shape (with Mediterranean specimens showing a longer and more pointed rostrum). The number of spinules on the merus of the first pereiopod, and on the basis and merus of the second pereiopod, showed also differences, as in *P. sivado*. The Mediterranean form of *P. multidentata* was even described as a different species, under the name of *Pasiphaea* (Phye) *sicula* Riggio, 1896, later on considered as a subspecies of *P. multidentata* (Zariquiey Álvarez 1946). Sivertsen and Holthuis (1956) did not accept the species status of *P. sicula*. After that Zariquiey Alvarez in 1957 reported that his father R. Zariquiey Cenarro had a paper in press detailing the differences between Mediterranean and Atlantic *P. multidentata* populations that was unfortunately lost during an air raid that destroyed the press during the Spanish civil war (1936-1939). He concluded that Mediterranean specimens could be assigned to *P. multidentata*, but clearly remarked the occurrence of these morphological differences that could in the future assign the Mediterranean specimens to a different "race", as then reported by him, or significantly different populations.

### i.v.i.ii. Family *Pandalidae*

#### *Chlorotocus crassicornis* (A. Costa, 1871)



*Chlorotocus crassicornis* is a nektobenthic species with an almost cosmopolitan distribution ranging from the North Atlantic through the Mediterranean, south-east Africa and Indo-Pacific region with a reported depth distribution ranging from 3-597 m (d'Udekem d'Acoz 1999, Li 2006). In the Mediterranean the species occurs from the mid continental shelf to the upper slope (D'onghia *et al.* 1998, Abelló *et al.* 2002), information about their biology and distribution is scarce (Carbonell and Abelló 1998, Vafidis *et al.* 2004, 2008). It shows a marked rhythmic night pattern of

catchability on continental shelf waters that shifts towards a dusk and dawn pattern on the upper slope (Aguzzi *et al.* 2007).

### *Plesionika heterocarpus* (A. Costa, 1871)



16

*Plesionika heterocarpus* is a nektobenthic species distributed along the eastern Atlantic and Mediterranean Sea, being a very common species of epibenthic trawl surveys. In the western Mediterranean, its depth distribution is comprised between the deep continental shelf to the middle slope (Abelló *et al.* 1988, 2002). In the Catalan Sea the species reproduces continuously around the year (Company and Sardà 2000, Company *et al.* 2003). Overall most knowledge on the species concerns its main life history characteristics, distribution, population genetics, role as a fishery bycatch, etc. (e.g. Maynou *et al.* 1996, Carbonell *et al.* 2003, Vafidis *et al.* 2005, Fanelli and Cartes 2008, Vafidis *et al.* 2008, Muñoz *et al.* 2012, Guijarro *et al.* 2012, García-Merchán *et al.* 2012).

## i.v.ii. Suborder *Dendrobranchiata*

Species of the suborder Dendrobranchiata comprise a group of phylogenetically basal decapods that, among other characteristics, differ from the Caridea and Stenopodidea shrimps by the branching form of the gills and by the fact that they release their eggs directly to the water instead of brooding them under the abdomen. Among the Dendrobranchiata, shrimps of the family Sergestidae and Benthesicymidae are among the commonest pelagic decapod species (Figure 5).




Figure 4: Scanner image of (a) *Sergestes arcticus*, (b) *Sergestes arachnipodus*, (c) *Sergia robusta* and (d) *Gennadas eleoans*.

### i.v.ii.i. Family *Sergestidae*

Judkins and Kensley (2008) made a thorough systematic study of this family and assigned some species to new genera. In the present thesis the proposed nomenclature of those authors was not followed since there is still some controversy about it and it is not completely accepted by the crustacean taxonomists. New changes are envisaged in the near future due to a wider use of genetic analyses, therefore the “classical” nomenclature was followed throughout this thesis.

### *Sergestes arcticus* Krøyer, 1855



*Sergestes arcticus* is an almost transparent species with red pigmentation restricted to some areas of its cephalothorax (Figure 4) (Vestheim and Kaartvedt 2009). The species performs vertical migrations where adults are distributed from 80 to 2100 m (Koukouras *et al.* 2000, Gartner *et al.* 2008) and preferentially from 500 m below the surface during the day and below around 150 m during the night (Franqueville 1971, Koukouras 2000). In the northwestern Mediterranean *S. arcticus* is one of the most abundant pelagic decapod species captured by pelagic IKMT trawls representing 65.4 % of the total capture (Franqueville 1971) and reaching high densities when sampled in their preferential depth. However, the species is not very abundant when sampled by demersal commercial gears in the Mediterranean Sea, due the big mesh size of the gear in relation to the species body size. This small pelagic crustacean feeds mainly on copepods, ostracods and euphausiids (Lagardere 1977), having a very important role within the continental slope and deep sea trophic nets (Cartes *et al.* 1998, 2002a, 2009). However, much information about their biology is still lacking (Company and Sardá 2000).

The species has a cosmopolitan distribution occurring in the Indo- Pacific south hemisphere (Australia and New Zealand), western and eastern Atlantic Ocean (from Canada, to Magellan Strait, and from Norway to South Africa) including also the whole Mediterranean Sea. Casanova (1977) identified morphological differences among south and north Atlantic and Mediterranean populations, concerning the shape of the petasma, and also among Atlantic and Mediterranean populations, concerning the presence/absence of a protuberance or spine in the first two pereopods.

*Sergestes vigilax* Stimpson, 1860

*S. vigilax* is an almost all transparent body similar to *S. arcticus* in terms of coloration, and size, but not on its morphology which is characteristic due to its highly developed third maxillipeds. Information on its biology is scarce while some on its depth distribution is available (Casanova and Judkins 1977, Hopkins et al 1994). The geographical distribution of *S. vigilax* comprises the Indo-Pacific, western Atlantic, north-eastern Atlantic and Mediterranean, being mainly sampled near the surface (Udekem d' Acoz 1999). In the Mediterranean Sea the species is rarely collected by demersal gears, while it is more common in samples taken by pelagic trawls (Koukouras 2000). Its densities, however never reach as high values as those of *S. arcticus* (Fasham 1979).

18

*Sergestes arachnipodus* (Coco, 1832)

*S. arachnipodus* is a morphologically similar species to *S. arcticus*, which can be identified by the presence of little red spots on its dorsal area, both on its cephalothorax and abdomen, and by their slightly larger size (Figure 4). Its global distribution comprises the Mediterranean and the western and eastern North Atlantic from temperate to tropical waters. Its depth distribution ranges from the surface to 2300 m depth (d'Udekem d'Acoz 1999, Koukouras 2000, Abelló *et al.* 2002, Company *et al.* 2004, Cartes *et al.* 2009), but general information about their biology is lacking.

*Sergia robusta* (S.I. Smith, 1882)

*S. robusta* is a species with an overall red colouration that is more linked the deep sea regions compared to the before mentioned Sergestidae species (Figure 4 and 5). It is present across the temperate north-west and north-east Atlantic from the Faroe Islands (61° N) to Cabo Verde (14° N) and Mediterranean (d'Udekem d'Acoz 1999). Overall its depth distribution comprises the pelagic environment between around 200 m and almost 5000 m (Koukouras *et al.* 2000). However, its preferential depth during the night is located around 200-500 m while during the day it is found in deeper waters down to 2000 m (Crosnier and Forest 1973, Froggia and Giannini 1982, Vereshchaka

1994). There are no gaps in the distribution of the species across the Atlantic-Mediterranean transition, even so slight morphological differences between Atlantic and Mediterranean males were reported by Casanova and Judkins (1976).

In the Mediterranean, the species is often found in demersal trawl hauls targetting the red shrimp *Aristeus antennatus*, as also are *P. multidentata* and *Gennadas elegans* (Fanelli *et al.* 2007, Guijarro *et al.* 2011, García-Rodríguez *et al.* 2011) (Figure 4). It shows a trend of increasing size with depth (Morales-Nin *et al.* 2003), and it is one of the largest species within the Sergestidae (Company and Sardá 2000). *S. robusta* has also an important role in the throphic dynamics of the demersal system being preyed by both nektonic and benthic species and feeding on copepods, ostracods, chaetognathes and euphausiids (Lagardere 1977, Cartes *et al.* 2007a, 2009).

### i.v.ii.ii. *Benthescymidae* family

#### *Gennadas elegans* (S.I. Smith, 1882)

*Gennadas elegans* is an all red pigmented shrimp that has the smallest body length compared to the before presented species (Figure 4 and 5). The species has also been shown to possess a differential high lipid storage which has been attributed to contribute to buoyancy and therefore maintain position in the water column (Childress and Nygaard 1974). The distribution of the species comprises the north-west and east Atlantic from Greenland to east Sahara and Mediterranean, dwelling in the pelagic environment from 30 to 2000 m below the surface (Franqueville 1971, Casanova and Judkins 1976, d'Udekem d'Acoz 1999). It mainly occurs on the bottom deeper than around 500 m but it can occur in much shallower waters in the pelagic environment during the night (Andersen *et al.*




Figure 5: Drawings of: (A) *Gennadas elegans*, (B) *Sergia robusta*, (C) *Pasiphaea multidentata*. Source: Würtz (2010).



1998, Koukouras 2000).

# Chapter

# I

Pasiphaeid shrimp populations in the western Mediterranean: geographical variability in distribution and population patterns



# Pasiphaeid shrimp populations in the western Mediterranean: geographical variability in distribution and population patterns

## 1.1. Introduction

Nekto-benthic decapod crustacean shrimps have a widespread geographic and bathymetric distribution from high latitudes in both hemispheres to intertropical waters (Crosnier and Forest 1973, Casanova and Judkins 1976, Fasham and Foxton 1979, Clarke and Holmes 1987, Hayashi 1999, Burghart *et al.* 2007, Serejo *et al.* 2007). Many continental slope shrimps show cyclic movements associated to the photoperiod as shown in some pasiphaeid and sergestid shrimps (Frogliia and Giannini 1982, Cartes *et al.* 1993, Aguzzi *et al.* 2007). This vertical daily migration ability present in the behaviour of these species (Naylor 2010) provides them with an important role in the transference of matter and energy from the upper primary productive layers of the ocean, where these species tend to feed, down to the continental slope epibenthic community, where they dwell during day-time hours (Cartes 1993c, Herring and Roe 1988, Naylor 2010). Nekto-benthic/benthopelagic decapod crustaceans constitute a fundamental food item to fish, other crustaceans and cephalopods with epibenthic habits on the continental slope and deep sea (Garrison and Link 2000, Fanelli and Cartes 2008). Some of these species have a relevant ecological importance as well as an exploitation potential as commercial target species (Nanjo and Ohtomi 2009). Relevant studies have however been mainly conducted only in a few regions, such as the Mediterranean Sea (e.g. Casanova and Judkniš 1977, Orsi-Relini and Relini 1990, Cartes *et al.* 1993, Company *et al.* 2001, 2003, Ramirez-Llodra *et al.* 2007), the Japan Sea (Nanjo 2007, Nanjo and Ohtomi 2009), or the Northeast and Southeast Atlantic Ocean (Matthews and Pinnoi 1973, Gibbons *et al.* 1994, Kensley and Schotte 2006).

The genus *Pasiphaea* has a worldwide distribution and over sixty species are known to date (Hayashi 1999, Tavares and Cardoso 2006, De Grave and Fransen 2011). Members of this family show a strong lateral compression of both cephalotorax and abdomen (Zariquiey Álvarez 1968), which may improve their swimming and dispersal abilities (Cartes *et al.* 1993, Aguzzi *et al.* 2009). While up to 18 species of the family Pasiphaeidae are known to occur in the northeast Atlantic and Mediterranean waters (eight of them belonging to the genus *Pasiphaea* (d'Udekem d'Acoz, 1999), only two of them are present in the Mediterranean Sea, namely *Pasiphaea sivado* and *Pasiphaea multidentata* (Zariquiey Álvarez 1968, Casanova and Judkins 1977, Koukouras 2000).

*Pasiphaea sivado* is a benthopelagic caridean shrimp commonly caught by demersal trawling on the upper slope across the East Atlantic and Mediterranean Sea (González-Gurriarán and Olaso 1987, Abelló *et al.* 2002). Biological studies on this species have been mainly performed in the western Mediterranean, where it has been reported to reproduce continuously throughout the year, although peaking in autumn-winter (Company *et al.* 2001). The smallest ovigerous female sampled was 14.7 mm CL and its longevity has been estimated to be around 2 years. *P. sivado* is a night feeder on pelagic crustaceans such as euphausiids and calanoid copepods (Lagardère 1972), as well as on mobile epibenthic organisms such as peracarid crustaceans (Cartes 1993b).

*Pasiphaea multidentata* inhabits benthic boundary layers in the middle and lower slope down to 2000 m depth (Cartes 1993a, Abelló *et al.* 2002, Tecchio *et al.* 2011), with juveniles inhabiting shallower waters than adults (Company *et al.* 2001). It also performs night vertical migrations into upper water layers, although apparently mainly restricted to juvenile individuals (Aguzzi *et al.* 2007, Cartes 1993c). In the northwestern Mediterranean the species shows a marked seasonality in reproduction with ovigerous females being only present from September to February, and with an estimated maximum longevity of around 3.5 years (Company *et al.* 2001, Ramirez-Llodra *et al.* 2007). It is an active nocturnal feeder on pelagic crustaceans such as *Gennadas elegans*, *P. sivado*, *P. multidentata*, sergestid shrimps and little mesopelagic fish as myctophids and *Cyclothone* spp. (Cartes 1993b).

Comparative studies of both species have addressed different life history processes. Company *et al.* (2001) suggested that the greater fecundity output of *P. sivado* supports its higher

population densities when compared with *P. multidentata*. Both species perform daily vertical migrations, with the apparent exception of the adults of *P. multidentata* (CL > 30 mm), which are assumed to perform epibenthic depth displacements (Cartes *et al.* 1993, Aguzzi *et al.* 2007). The dietary overlap between these species is low due the different size spectra of their respective prey items (Cartes 1993b).

This paper aims to analyze the main characteristics of the bathymetric and geographic distribution, population size structure and some reproductive characteristics of both *Pasiphaea sivado* and *P. multidentata* along the Mediterranean coasts of the Iberian Peninsula, and to relate the patterns obtained with geomorphological and hydrographic characteristics. The data base was obtained in a series of annual demersal fisheries research surveys performed throughout the continental shelf and upper slope of the western Mediterranean (Bertrand *et al.* 2002).

## 1.2. Material and methods

### 1.2.1. Study area and oceanographic context

The study area encompassed the continental shelf, upper and middle slope down to 800 m along the Iberian Peninsula Mediterranean coasts from Gibraltar in the SW to Cape Creus in the NE (Figure 1). Overall, the continental shelf is very narrow in the Alboran Sea and Vera Gulf, south of Cape Palos, and widens to the north, reaching a maximum width of up to 70 km in the Ebro Delta - Columbretes area. North of Barcelona, the continental shelf is heavily indented by several submarine canyons.

The Western Mediterranean is influenced by the inflow of Atlantic water through the Strait of Gibraltar (Hopkins 1985, Millot 2005), where lighter Atlantic water inflows towards the Mediterranean on surface waters, and higher density Mediterranean water outflows towards the Atlantic Ocean in depth. This surface inflow of Atlantic waters generates two anticyclonic gyres between the Strait of Gibraltar and Cape Gata, and adjacent upwelling cells in the vicinity of the

Strait (Vargas-Yáñez and Sabatés 2007). The main current of inflowing Atlantic waters is directed from Cape Gata towards the North African coast, generating the Almeria-Oran front (AOF), therefrom continuing its inflow along the North African coasts towards the Central and Eastern Mediterranean. The AOF is a strong thermohaline density front confined to the upper layers of the



Figure 1: Iberian Peninsula Mediterranean coast showing the study regions: Western Alboran (WALB); Eastern Alboran (EALB); Alboran Island (ALBO); Vera Gulf (VERA); Alicante (ALIC); Ibiza Island (IBIZ); Valencia (VALE); Ebro Delta region (DELT); Central Catalonia (CCAT) and North Catalonia (NCAT). 200 m and 1000 m isobaths are shown.

water column and shows important seasonal and interannual variability in strength (Tintoré *et al.* 1988). The Atlantic water also flows north-eastwardly, due to the detachment of anticyclonic gyres which reach the Balearic Islands and generate a second density front along the northeastern part of the archipelago, associated to the NE flowing Balearic Current (Tintoré *et al.* 1988, López-Jurado *et al.*, 2008, Monserrat *et al.* 2008). The interaction between the strong Northern Current, flowing southwestwards along the continental slope from the Gulf of Lions, and the Balearic Current in the Ibiza Channel region (between Cape La Nao and the island of Ibiza) generates a cyclonic gyre over the Balearic basin enclosing the older resident waters in its center (Salat 1995, Sabatés *et al.* 2007).

## 1.2.2. Sampling and Analysis

The material analyzed in the present work was obtained from the annual MEDITS\_ES surveys performed between 1994 and 2008. The aim of the Mediterranean International Trawl Surveys project (MEDITS) is to obtain density, biomass and recruitment indexes of the main target species exploited by the demersal fishery throughout the European Union and adjacent Mediterranean countries, based in a common sampling protocol (Bertrand *et al.* 2000, 2002). The Spanish surveys were performed on board R/V 'Cornide de Saavedra'. These cruises took place in spring, centered in the month of May and had duration of about 30 days. All hauls were performed during day-time hours. Overall, samples were taken at depths from around 40 m down to 800 m depth based on a sample design randomly stratified by geographical sector and depth stratum. Ten geographic sectors were established according to geomorphology characteristics of the overall area and previous biogeographic results (Abelló *et al.* 2002, Rufino *et al.* 2005): Western Alboran (WALB) from Gibraltar to Cape Sacratif, Eastern Alboran (EALB) from Cape Sacratif to Cape Gata, Alboran Island (ALBO), Vera Gulf (VERA) from Cape Gata to Cape Palos, Alicante (ALIC) from Cape Palos to Cape La Nao, Ibiza Island (IBIZ), Valencia (VALE) from Cape La Nao to Castelló/Columbretes Islands, Ebro Delta region (DELTA) from Castelló/Columbretes Islands to Cape Salou/Tarragona, Central Catalonia (CCAT) from Cape Salou/Tarragona to Barcelona, and North Catalonia (NCAT), from Barcelona to Cape Creus. The bottom trawl used was a GOC-73 model (Fiorentini *et al.* 1999). It has a large, up to 3 m vertical opening allowing the capture of epibenthic and benthopelagic fish and crustaceans, and a codend stretched mesh size of 20 mm. The trawls were performed at a speed of 3 knots with duration of one hour, except for the hauls performed above the 200 m isobath that had duration of 30 minutes. A total of 1741 valid hauls were performed along the studied period (Table 1).

On board, after each haul, the total amount of each species sampled was weighed and counted. Random sub-samples were taken whenever necessary. Density and biomass were later standardized by swept area, to obtain number of individuals and weight (in g) per square kilometer. The swept area was calculated taking into account the horizontal opening of the trawl, measured with Scanmar devices, and distance from starting point of the haul (net on the bottom) to end of the effective haul (net off the bottom), measured from GPS latitude and longitude readings. The

frequency of occurrence, density, biomass and size structure was analyzed in terms of bathymetrical and geographical distribution in the two species. Mean density and biomass values

Depth stratum (m)	WALB	EALB	ALBO	VERA	ALIC	IBIZ	VALE	DELT	CCAT	NCAT	Total
0-50	22	15	-	8	31	-	13	43	26	6	164
51-100	47	35	2	23	86	-	76	172	62	33	536
101-150	23	11	3	5	46	6	35	44	24	49	246
151-200	9	10	-	18	23	-	5	5	4	7	81
201-250	10	11	-	-	7	5	-	-	10	6	49
251-300	7	6	-	20	20	8	6	1	3	9	80
301-350	21	10	2	6	12	4	10	1	5	15	86
351-400	14	5	5	1	4	2	2	1	5	3	42
401-450	16	14	-	8	14	1	-	2	6	10	71
451-500	1	3	1	2	19	11	-	-	3	11	51
501-550	21	14	4	9	18	4	-	-	6	11	87
551-600	13	10	3	3	19	10	3	-	6	5	72
601-650	22	15	2	1	5	-	2	-	5	11	63
651-700	12	1	2	3	1	11	-	-	6	8	44
701-750	15	1	1	7	-	2	10	-	2	2	40
751-800	13	7	3	-	-	-	5	-	1	-	28
<b>Total</b>	<b>266</b>	<b>168</b>	<b>28</b>	<b>114</b>	<b>305</b>	<b>64</b>	<b>167</b>	<b>269</b>	<b>174</b>	<b>186</b>	<b>1741</b>

were calculated by averaging positive haul density values over total number of hauls made within each geographic sector – depth interval cell. One-way analysis of variance was used to test for the occurrence of significant interannual differences in density (natural logarithmic transformation), after testing for normality of data and homogeneity of variances. If no significant differences were detected, a two-way ANOVA was used considering geographical sectors and 100 m depth strata as factors to test for the occurrence of significant differences among these factors. As a requisite for ANOVA, only factor cells with positive values were used in the analyses, so that only the most relevant sectors and strata were used.

From 1998, all individuals, or a subsample of up to 60 individuals, of each species, *Pasiphaea sivado* and/or *P. multidentata*, for each haul, were sexed and measured (carapace length, CL), with an accuracy of 0.1 mm. Ovigerous females were also noted down. Size frequency distributions (SFD), weighted by the density of the corresponding sample, were obtained for each combination of geographic sector and depth stratum. SFD based on <15 individuals per cell have not been shown. Normal-distributed components were identified in SFD using the Bhattacharya method implemented in Fisat II (Gayanillo *et al.* 2005). Mean sizes identified with this method for each depth stratum were later pooled in a frequency distribution by sector in order to identify the commonest mean size classes within each geographic sector. This would provide greater precision on the actual number of age classes present in each geographic sector and on their average size (Yamasaki 1988, Queiroga 1993). The cumulative frequency of ovigerous females per sector was adjusted to a logistic distribution in order to calculate the mean size of occurrence of ovigerous



females of *P. sivado* per sector. These were tested with a t-test.

Table 2: Mean bottom temperature (°C) and salinity  $\pm$  SD in each 50 m depth stratum in the five most relevant geographic sectors for *Pasiphaea* spp. Light grey cells show the preferential depth strata for *P. sivado*; dark grey cells show the preferential depth strata for *P. multidentata*

Depth stratum (m)	TEMPERATURE					SALINITY				
	WALB	EALB	ALIC	CCAT	NCAT	WALB	EALB	ALIC	CCAT	NCAT
001-050	13.96 $\pm$ 0.75	14.60 $\pm$ 1.28	14.15 $\pm$ 0.45	14.75 $\pm$ 1.42	14.61 $\pm$ 1.08	38.01 $\pm$ 0.45	37.78 $\pm$ 0.58	38.08 $\pm$ 0.17	38.00 $\pm$ 0.15	37.91 $\pm$ 0.17
051-100	13.57 $\pm$ 0.28	14.09 $\pm$ 0.84	13.61 $\pm$ 0.61	13.94 $\pm$ 0.47	13.31 $\pm$ 0.26	38.15 $\pm$ 0.16	37.94 $\pm$ 0.45	38.14 $\pm$ 0.11	38.03 $\pm$ 0.12	38.07 $\pm$ 0.19
101-150	13.32 $\pm$ 0.09	13.41 $\pm$ 0.22	13.15 $\pm$ 0.24	13.35 $\pm$ 0.41	13.25 $\pm$ 0.29	38.31 $\pm$ 0.07	38.26 $\pm$ 0.05	38.24 $\pm$ 0.13	38.10 $\pm$ 0.11	38.13 $\pm$ 0.10
151-200	-----	13.30 $\pm$ 0.13	13.08 $\pm$ 0.12	13.61 $\pm$ 0.43	-----	-----	38.35 $\pm$ 0.09	38.28 $\pm$ 0.12	38.06 $\pm$ 0.10	-----
201-250	13.27 $\pm$ 0.03	13.38 $\pm$ 0.17	13.24 $\pm$ 0.39	13.15 $\pm$ 0.16	13.18 $\pm$ 0.14	38.37 $\pm$ 0.12	38.29 $\pm$ 0.10	37.58 $\pm$ 1.33	38.23 $\pm$ 0.07	38.34 $\pm$ 0.05
251-300	-----	13.72 $\pm$ 0.95	13.14 $\pm$ 0.07	13.24 $\pm$ 0.01	-----	-----	38.34 $\pm$ 0.22	38.38 $\pm$ 0.06	38.32 $\pm$ 0.09	-----
301-350	13.26 $\pm$ 0.02	13.29 $\pm$ 0.07	13.14 $\pm$ 0.07	13.04 $\pm$ 0.03	13.30 $\pm$ 0.14	38.45 $\pm$ 0.07	38.41 $\pm$ 0.09	38.42 $\pm$ 0.07	38.36 $\pm$ 0.02	38.39 $\pm$ 0.15
351-400	13.27 $\pm$ 0.01	13.25 $\pm$ 0.08	-----	13.26 $\pm$ 0.00	-----	38.50 $\pm$ 0.02	38.42 $\pm$ 0.01	-----	38.35 $\pm$ 0.04	-----
401-450	13.24 $\pm$ 0.02	13.26 $\pm$ 0.03	13.22 $\pm$ 0.09	13.20 $\pm$ 0.03	13.31 $\pm$ 0.00	38.50 $\pm$ 0.02	38.50 $\pm$ 0.01	38.50 $\pm$ 0.03	38.43 $\pm$ 0.06	38.50 $\pm$ 0.02
451-500	-----	13.36 $\pm$ 0.09	13.20 $\pm$ 0.15	-----	13.24 $\pm$ 0.14	-----	38.39 $\pm$ 0.17	38.45 $\pm$ 0.13	-----	38.47 $\pm$ 0.06
501-550	13.22 $\pm$ 0.05	13.20 $\pm$ 0.08	13.16 $\pm$ 0.09	-----	13.29 $\pm$ 0.06	38.48 $\pm$ 0.04	38.49 $\pm$ 0.02	38.42 $\pm$ 0.14	-----	38.51 $\pm$ 0.04
551-600	13.23 $\pm$ 0.03	13.20 $\pm$ 0.07	13.12 $\pm$ 0.09	13.12 $\pm$ 0.22	-----	38.50 $\pm$ 0.03	37.86 $\pm$ 1.37	38.44 $\pm$ 0.12	38.35 $\pm$ 0.15	-----
601-650	13.18 $\pm$ 0.04	13.25 $\pm$ 0.086	14.33 $\pm$ 2.45	-----	13.17 $\pm$ 0.13	38.48 $\pm$ 0.04	38.38 $\pm$ 0.12	38.37 $\pm$ 0.16	-----	38.50 $\pm$ 0.02
651-700	-----	-----	-----	13.13 $\pm$ 0.07	-----	-----	-----	-----	38.50 $\pm$ 0.01	-----
701-750	13.14 $\pm$ 0.04	-----	-----	13.14 $\pm$ 0.18	-----	38.48 $\pm$ 0.02	-----	-----	38.40 $\pm$ 0.09	-----
751-800	-----	13.15 $\pm$ 0.06	-----	-----	-----	-----	38.49 $\pm$ 0.02	-----	-----	-----

Information on mean temperature and salinity on the bottom during each trawl was recorded with a CTD SBE-37 placed at the float-line of the net (Table 2). Data from the 2001-2006 cruises were used to calculate mean temperature and salinity by depth stratum for each geographic sector. In order to assess the optimal temperature and salinity window for each species, the range, 25 and 75 percentiles and median values were calculated. For each species each sample was additionally categorized as juvenile or adult depending on its mean CL: For *P. sivado*, samples with mean CL $\leq$ 16 mm were considered as juvenile samples, and those with CL >16 mm as adult; for *P. multidentata*, a size break at 20 mm CL was considered (see below).

## 1.3. Results

### 1.3.1. *Pasiphaea sivado*

#### 1.3.1.1. Occurrences

The overall depth distribution of *P. sivado* along the study area ranged between 141 and 765 m, however the highest frequency of occurrence was located between the 350 and 500 m depth strata (Table 3a). At depths both shallower and deeper the occurrence of *P. sivado* was progressively rarer (Figure 2a).




Figure 2: Overall percentage occurrence ( $\pm$  95% confidence intervals) of *Pasiphaea sivado* (a) and *P. multidentata* (b) by depth intervals along the Iberian Peninsula Mediterranean coast.

Taking into account both the frequency of occurrence within the preferential depth interval of the species (350-500 m, Table 3a), and the sampling effort within each geographic sector (i.e. excluding those sectors where the number of samples taken in the preferential depth strata of the species were  $< 20$ , Table 1), the highest frequencies of occurrence of *P. sivado* were located in Western and Eastern Alboran, Alicante and Northern Catalonia (Table 3a).

Table 3: Frequency of occurrence of (a) *Pasiphaea sivado* and (b) *Pasiphaea multidentata* (in %) within each geographic sector and depth stratum. Occurrences larger than 75% are highlighted in bold. Cells with (\*) correspond to data with less than 3 hauls (see Table 1), not shown since considered not enough informative due to low sampling size. Values in the lowest line ((350 to 500 m in *P. sivado* and >500 m in *P. multidentata*) correspond to the overall percentage occurrence by sector restricted to the depth strata with overall percentage occurrence higher than 75%.

<b>(a) <i>Pasiphaea sivado</i></b>											
Depth stratum (m)	WALB	EALB	ALBO	VERA	ALIC	IBIZ	VALE	DELT	CCAT	NCAT	Total
0-50	0.0	0.0	-	0.0	0.0	-	0.0	0.0	0.0	0.0	0.0
50-100	0.0	0.0	0.0	0.0	0.0	-	0.0	0.0	0.0	0.0	0.0
100-150	4.3	0.0	0.0	0.0	4.3	0.0	0.0	0.0	0.0	0.0	1.2
150-200	22.2	0.0	-	11.1	0.0	-	20.0	0.0	0.0	0.0	6.2
200-250	60.0	0.0	-	-	14.3	0.0	-	-	20.0	0.0	18.4
250-300	57.1	50.0	-	30.0	45.0	0.0	66.7	*	33.3	33.3	38.8
300-350	81.0	90.0	*	50.0	75.0	0.0	70.0	*	100.0	80.0	73.3
350-400	100.0	100.0	40.0	*	75.0	*	*	*	100.0	100.0	88.1
400-450	93.8	100.0	-	62.5	92.9	*	-	*	83.3	100.0	90.1
450-500	*	100.0	100.0	50.0	73.7	100.0	-	-	66.7	54.5	76.5
500-550	90.5	50.0	0.0	44.4	61.1	25.0	-	-	50.0	81.8	62.1
550-600	69.2	0.0	0.0	0.0	5.3	20.0	0.0	-	50.0	40.0	23.6
600-650	50.0	26.7	0.0	*	0.0	-	50.0	-	0.0	27.3	31.7
650-700	41.7	*	*	33.3	*	0.0	-	-	16.7	0.0	15.9
700-750	6.7	*	*	0.0	-	*	0.0	-	*	*	10.0
750-800	15.4	0.0	0.0	-	-	-	0.0	-	*	-	7.1
No. samples (350-500 m)	31	22	6	11	37	14	2	3	14	24	164
Mean frequency (350 to 500 m)	96.8	100.0	50.0	63.6	81.1	85.7	100.0	100.0	85.7	79.2	85.4
<b>(b) <i>Pasiphaea multidentata</i></b>											
Depth stratum (m)	WALB	EALB	ALBO	VERA	ALIC	IBIZ	VALE	DELT	CCAT	NCAT	Total
200-250	0.0	0.0	-	-	0.0	0.0	-	-	0.0	0.0	0.0
250-300	14.3	0.0	-	5.0	0.0	0.0	0.0	*	0.0	0.0	3.8
300-350	0.0	0.0	0.0	0.0	8.3	0.0	20.0	*	0.0	0.0	3.5
350-400	0.0	20.0	*	*	0.0	*	0.0	*	0.0	0.0	2.4
400-450	6.3	35.7	-	37.5	35.7	*	-	*	50.0	50.0	32.4
450-500	*	66.7	*	*	52.6	18.2	-	-	100.0	36.4	43.1
500-550	71.4	71.4	100.0	100.0	88.9	100.0	-	-	83.3	54.5	79.3
550-600	61.5	90.0	66.7	100.0	84.2	90.0	66.7	-	66.7	100.0	80.6
600-650	77.3	100.0	*	*	100.0	-	*	-	60.0	81.8	84.1
650-700	91.7	*	*	100.0	*	100.0	-	-	66.7	75.0	88.6
700-750	86.7	*	*	100.0	-	*	100.0	-	*	*	92.5
750-800	92.3	100.0	100.0	-	-	-	100.0	-	*	-	100.0
No. samples (>500 m)	96	48	15	23	43	27	20	0	26	37	334
Mean frequency (>500 m)	79.2	87.5	86.7	100.0	88.4	96.3	95.0	-	73.1	75.7	85.1

Table 3a also clearly shows that *P. sivado* in Western Alboran, contrasting with the rest of geographical sectors, presented both a much shallower and a deeper distribution, since the species occurred from the 100-150 m stratum down to the deepest sampled stratum (750-800 m), whereas in the rest of areas the distribution of *P. sivado* usually started from the 200-250 m stratum down to maximum depths of around 600-700 m.

### 1.3.1.2. Density and biomass

Positive densities of *P. sivado* per haul ranged between 8 and 186,109 ind.km<sup>-2</sup>. No significant differences were found concerning interannual variability in densities (ANOVA,  $F_{14,340}=0.7313$ ,  $p>0.5$ ). The interaction between depth and geographical sector was not significant. Two-way ANOVA showed that densities across both geographical sectors ( $F_5=4.331$ ,  $p<0.001$ ) and depth strata ( $F_3=20.4$ ,  $p<0.0001$ ) differed significantly among them. No significant interaction was found ( $p>0.1$ ). The highest mean density values were found at depths between 250-500 m, i.e. within the preferential occurrence depth strata for the species (350-500 m) and also at the two adjacent upper strata. The Western Alboran population showed the widest density distribution range in depth for all sampled populations. Sizeable densities were found in this sector at depths between 150 and 600 m. In the rest of sampled areas, densities rose sharply only from 250 m downwards to around 500-600 m. Overall, the areas with highest mean density values were Western and Eastern Alboran and Central and North Catalonia. Ibiza also showed very high mean densities, but the relatively low sample size and the large variability in densities in this area did not allow to take these values as fully representative (Table 3a). Both density and biomass (Tables 4a and 5a) followed a similar bathymetric and geographic pattern along the studied geographic area.

31

The relationship between percentage occurrence and mean density per depth strata (Figure 3) showed that high densities were associated to depths with a high percentage occurrence of the




Figure 3: Relationship between mean density and percentage occurrence in *Pasiphaea sivado* and *P. multidentata*.

species, especially so in depth strata comprised between 350 and 500 m, which could therefore be categorised as the optimal depth ranges for *P. sivado* in the study area for both occurrence and density. It also showed that densities gradually increased with increasing depth and percentage occurrence up to the 350-400 m depth stratum, and sharply decreased deeper than 500 m. Also, the ratio between biomass and density (i.e. an index of mean weight per individual within each depth stratum) followed a significant increasing trend with depth ( $P < 0.01$ ), clearly indicating the occurrence of a marked size increase with depth (Figure 4).

Table 4: Mean density (ind.km<sup>-2</sup>) of (a) *Pasiphaea sivado* and (b) *Pasiphaea multidentata* within each depth stratum and geographic sector along the Iberian Peninsula Mediterranean coast. Mean density per geographic sector corresponds to densities within the 350-500 m depth strata for *P. sivado* and >500 m for *P. multidentata* (i.e. those with the overall highest frequency of occurrence).

(a) <i>Pasiphaea sivado</i>											
Depth stratum (m)	WALB	EALB	ALBO	VERA	ALIC	IBIZ	VALE	DELT	CCAT	NCAT	Total
0-50	0	0	-	0	0	-	0	0	0	0	0
50-100	0	0	0	0	0	-	0	0	0	0	0
100-150	1	0	0	0	3	0	0	0	0	0	1
150-200	411	0	-	3	0	-	4	0	0	0	47
200-250	3347	0	-	-	9	0	-	-	11	0	687
250-300	7146	1048	-	750	3145	0	295	3167	61	56	1748
300-350	8663	3453	0	119	4288	0	1027	176	3328	2400	3857
350-400	9200	2847	81	2350	426	1588	458	436	18358	2483	5983
400-450	13614	7387	-	187	1782	0	-	108	6879	8916	6737
450-500	5867	110	588	296	442	26368	-	-	299	987	6227
500-550	494	55	0	158	110	3	-	-	61	380	220
550-600	156	0	0	0	0	3	0	-	23	40	33
600-650	17	4	0	657	0	-	5	-	0	7	19
650-700	10	0	0	18	0	0	-	-	128	0	21
700-750	2	0	0	0	-	0	0	-	44	44	5
750-800	3	0	0	-	-	-	0	-	0	-	1
No. samples (350-500 m)	31	22	6	11	37	14	2	3	14	24	164
Mean density (350-500 m)	9560	3448	335	944	883	9319	458	272	8512	4129	6316
(b) <i>Pasiphaea multidentata</i>											
Depth stratum (m)	WALB	EALB	ALBO	VERA	ALIC	IBIZ	VALE	DELT	CCAT	NCAT	Total
0-50	0	0	-	0	0	-	0	0	0	0	0
50-100	0	0	0	0	0	-	0	0	0	0	0
100-150	0	0	0	0	0	0	0	0	0	0	0
150-200	0	0	-	0	0	-	0	0	0	0	0
200-250	0	0	-	-	0	0	-	-	0	0	0
250-300	74	0	-	0	0	0	0	11	0	0	7
300-350	0	0	0	0	11	0	25	0	0	0	4
350-400	0	2	0	0	0	0	0	0	0	0	0
400-450	4	22	-	56	117	0	-	199	141	80	63
450-500	0	80	0	50	97	18	-	-	142	379	137
500-550	28	42	92	281	124	94	-	-	199	11	92
550-600	34	37	28	59	281	502	58	-	127	263	190
600-650	124	38	70	213	178	-	102	-	16	46	85
650-700	24	48	99	1029	439	295	-	-	21	85	185
700-750	87	0	66	101	-	95	93	-	158	138	95
750-800	36	35	37	-	-	-	186	-	154	-	67
No. samples (>500 m)	96	48	15	23	43	27	20	0	26	37	335
Mean density (>500 m)	56	33	65	337	256	247	110	-	113	109	119

Table 5: Mean biomass (g km<sup>-2</sup>) of (a) *Pasiphaea sivado* and (b) *Pasiphaea multidentata* within each depth stratum and geographic sector along the Iberian Peninsula Mediterranean coast. Mean biomass per geographic sector corresponds to values within the 350-500 m depth strata for *P. sivado*) and >500 m for *P. multidentata* (i.e. those with the overall highest frequency of occurrence)

(a) <i>Pasiphaea sivado</i>											
Depth stratum (m)	WALB	EALB	ALBO	VERA	ALIC	IBIZ	VALE	DELT	CCAT	NCAT	Total
0-50	0	0	-	0	0	-	0	0	0	0	0
50-100	0	0	0	0	0	-	0	0	0	0	0
100-150	1	0	0	0	4	0	0	0	0	0	1
150-200	150	0	-	4	0	-	4	0	0	0	18
200-250	3979	0	-	-	17	0	-	-	18	0	818
250-300	9174	1020	-	524	2656	0	155	4642	29	32	1749
300-350	13853	2961	0	139	4023	0	1423	301	3146	2811	5140
350-400	12988	4967	131	1224	490	2299	664	185	25558	2739	8396
400-450	22070	7192	-	287	2021	0	-	155	5306	9323	8588
450-500	15496	178	1098	477	568	35611	-	-	436	1159	8523
500-550	1141	91	0	266	184	3	-	-	94	650	444
550-600	343	0	0	0	0	2	0	-	29	54	68
600-650	34	6	0	1137	0	-	11	-	0	6	33
650-700	22	0	0	25	0	0	-	-	72	0	17
700-750	3	0	0	0	-	0	0	-	67	44	7
750-800	6	0	0	-	-	-	0	-	0	-	3
No. samples (350-500 m)	31	22	6	11	37	14	2	3	14	24	164
Mean biomass (350-500 m)	16851	4112	614	663	1026	12637	664	170	10433	4407	8502
(b) <i>Pasiphaea multidentata</i>											
Depth stratum (m)	WALB	EALB	ALBO	VERA	ALIC	IBIZ	VALE	DELT	CCAT	NCAT	Total
0-50	0	0	-	0	0	-	0	0	0	0	0
50-100	0	0	0	0	0	-	0	0	0	0	0
100-150	0	0	0	0	0	0	0	0	0	0	0
150-200	0	0	-	0	0	-	0	0	0	0	0
200-250	0	0	-	-	0	0	-	-	0	0	0
250-300	88	0	-	2	0	0	0	22	0	0	8
300-350	0	0	0	0	33	0	145	0	0	0	21
350-400	0	14	0	0	0	0	0	0	0	0	2
400-450	26	130	-	361	693	0	-	945	370	102	281
450-500	0	501	0	317	466	88	-	-	266	209	295
500-550	167	198	577	2009	575	449	-	-	773	59	507
550-600	230	252	165	386	1551	2823	338	-	726	1351	1069
600-650	696	256	413	1314	999	-	577	-	114	271	492
650-700	165	402	577	7568	2718	1779	-	-	118	351	1183
700-750	672	0	485	584	-	587	656	-	917	944	653
750-800	245	281	861	-	-	-	1064	-	1121	-	437
No. samples taken (> 500 m)	96	48	15	23	43	27	20	0	26	37	335
Mean biomass (>500 m)	362	231	513	2372	1461	1410	659	-	628	595	723




Figure 4: Mean density (black bars), biomass (grey bars) and biomass/density ratio (dots) of *Pasiphaea sivado* and *P. multidentata* by depth strata.

### 1.3.1.3. Size structure and size at maturity

Overall, sizes of *P. sivado* ranged from 9.2 to 26.1 mm CL. Size frequency distributions per 100 m depth strata and geographic sector (Figure 5) showed that the populations in the Alboran Sea reached larger sizes than those in the northwestern Mediterranean and the intermediate Alicante sector. The western Alboran population showed a clear size-increasing trend with depth, with juvenile individuals being restricted to the upper 100-300 m depth strata; the largest individuals were recorded in the two deepest strata (500-700 m). Populations along the Catalan coast did not show such a marked increasing trend, but rather showed a similar size structure with depth, except for the dominant presence of juveniles in the upper depth occurrence stratum (200-300 m).




Figure 5: Size frequency distributions of *Pasiphaea sivado* per depth interval in the five best sampled geographic sectors. Total N: WALB = 1502; EALB = 592; ALIC = 999; CCAT = 653; NCAT = 462

Based on the analysis of significant normally-distributed cohorts on the several SFD by depth stratum using the Bhattacharya method (Figure 7), the frequency distribution of the mean sizes of the normally-distributed identified cohorts per sector (Figure 8) showed that most populations appeared to be structured in two main cohorts, broadly corresponding to juvenile (mainly placed around 14 mm CL) and adult individuals (located mainly around 19 mm CL); the populations in Alboran Sea also showed additional significant cohorts around 22 mm CL which were not identified in the rest of sampled sectors.

To calculate the mean size of ovigerous females per geographic sector, a logistic curve was fitted to the cumulative frequency of ovigerous females per sector (Figure 9). This showed that the populations from western and eastern Alboran significantly overlapped (Table 6). These sizes were significantly larger than those in the rest of sampled regions, which distributions also broadly and significantly overlapped.






Figure 7: Mean size of the different normally-distributed identified components (using the Bhattacharya method implemented in Fisat II) in the size frequency distributions of *Pasiphaea sivado* and *P. multidentata* by geographic sector (NCAT: North Catalonia; CCAT: Central Catalonia; ALIC: Alicante; EALB: Eastern Alboran; WALB: Western Alboran)

**Table 6** (a) Minimum, maximum, and mean (SD and N) (obtained by fitting a logistic curve) sizes of ovigerous female *Pasiphaea sivado* per geographic sector. (b) Pairwise comparisons of mean ovigerous female sizes by geographic sector (t-test values).

(a)					
Carapace Length	WALB	EALB	ALAC	CCAT	NCAT
Min	14.5	17.2	17.5	16.4	17.2
Max	24.8	23.2	21.5	22.1	22.7
Mean	20.7	20.5	19.1	19.6	19
SD	1.8	1.7	1.2	1.6	1.6
N	243	59	49	37	20

(b)				
	EALB	ALAC	CCAT	NCAT
WALB	n.s.	***	***	***
EALB		***	*	***
ALAC			n.s.	n.s.
CCAT				n.s.




Figure 8: Frequency distribution of normally-distributed components identified in the size frequency distributions of *Pasiphaea sivado* and *P. multidentata* by geographic sector (NCAT: North Catalonia; CCAT: Central Catalonia; ALIC: Alicante; EALB: Eastern Alboran; WALB: Western Alboran). Size class intervals of 1 mm in *P. sivado* and 2 mm in *P. multidentata*.

### 1.3.1.4. Temperature and salinity

Occurrences of *P. sivado* took place at bottom temperatures comprised between 12.98 and 13.42°C and salinities between 38.18 and 38.54. The temperature-salinity (TS) window was




Figure 10 Temperature and salinity windows (25-75 percentiles) for juvenile (dashed lines) and adult (solid lines) of *Pasiphaea sivado* (grey) and *P. multidentata* (black). Only temperature/salinity data from CTD samples taken in the depth range 200–800 m are shown.

narrower for adults than for juveniles (Figure 10). In particular, adults occurred (percentiles 25-75) at temperatures between 13.17 and 13.27°C and salinities of 38.43 and 38.51, while juveniles occurred at temperatures between 13.04 and 13.28°C and salinities between 38.36 and 38.48.

## 1.3.2. *Pasiphaea multidentata*

### 1.3.2.1. Occurrences

The overall depth distribution of *P. multidentata* along the study area ranged between 265 and 799 m. Its frequency of occurrence was null at strata shallower than 250 m and very low down to 400 m. Then it increased sharply down to 500 m, and reached occurrences higher than 75% from 500 m down to the deepest sampled stratum (750-800 m) (Figure 2b), clearly showing that the sampling strategy did not fully encompass the overall bathymetric range of the species.

As in *P. sivado*, the upper occurrence limit of *P. multidentata* in WALB and VERA took place at depths shallower (250-300 m) than in the rest of Mediterranean geographic sectors sampled (Table 2). At depths greater than 500 m (where the frequency of occurrence was > 75%) the species was common throughout the geographic area studied. However, and differing from *P. sivado*, the overall frequency of occurrence values were lower (73-76%) in the Balearic basin (excluding the Gulf of Valencia sector) than in the rest of geographic sectors sampled, all of them located in the Algerian basin, where overall frequencies of occurrence were markedly higher (most of them being higher than 85%).

### 1.3.2.2. Density and biomass

Positive density values of *P. multidentata* ranged between 7.5 and 3,696 ind.km<sup>-2</sup>. No significant differences were found concerning interannual variability in densities (ANOVA,  $F_{14,321}=1.266$ ,  $p>0.1$ ). The interaction between depth and geographical sector was not significant. Two-way ANOVA showed that densities across geographical sectors ( $F_6=7.489$ ,  $p<0.0001$ ) differed significantly among them but not among depth strata ( $F_3=0.5484$ ,  $p>0.5$ ). Significant interaction was detected ( $p<0.01$ ). The highest mean density values were generally found deeper than 500 m in those sectors belonging to the Algerian basin (from Western Alboran to Ibiza), whereas in the northernmost sectors the highest mean densities were found slightly shallower (400-500 m) (Table 3b). Overall, the highest mean densities were found in the intermediate sectors, from Vera to Ibiza; densities were also high in the northernmost sectors (Central and Northern Catalonia), while the lowest density values were detected in the Alboran Sea sectors. Biomass however showed a slightly different pattern, since the depth of highest biomass values was found below 500 m in all sampled sectors (except Eastern Alboran), including those in the Balearic basin, a fact related to the different size structure found in the two basins, with a higher occurrence of juveniles, occurring in shallower waters, in the northern sectors, as compared to the rest of sampled sectors (see below).

The relationship between percentage occurrence and mean density per depth strata (Figure 3b) showed that high densities were associated to depths with a high percentage occurrence of the species, especially so in strata deeper than 450 m, which would accordingly be categorised as the

optimal depth ranges for *P. multidentata* in the study area for both occurrence and density, taking always into account that the depth distribution range of the species was not fully covered by the samplings. Occurrences of the species at depths shallower than 400 m were always associated to very low densities.

Concerning the ratio between biomass and density by depth (i.e. an index of mean weight per individual within each depth stratum), it also showed, as in *P. sivado*, a significant increasing trend with depth ( $P < 0.01$ ), clearly suggesting the occurrence of a marked size increase with depth (Figure 4b).

### 1.3.2.3. Size structure

Sizes of *P. multidentata* ranged between 7.7 and 47.9 mm CL. The most remarkable feature that size frequency distributions per depth and sector showed (Figure 6) is that the occurrence of juveniles was practically restricted to the sectors of the Balearic basin (CCAT and NCAT), where the population size structure was clearly bimodal, with the juvenile cohort being present at sizes comprised between 8 and 16 mm CL and adults found at sizes between around 24 and 32 mm CL. In these northern sectors, the occurrence of individuals with sizes around or larger than 40 mm CL was practically anecdotal. In contrast, very few or hardly any juvenile was present in the populations from the Alboran Sea and Alicante, the sectors belonging to the Algerian basin. In these sectors, most of the population was comprised of adult individuals of sizes ranging between 24 and 32 mm CL, but also a third, larger, cohort was discernible at sizes of around 40 mm in most samples.

The frequency distribution of the mean sizes of the cohorts identified in the SFD by depth strata (Figures 7 and 8), showed that most populations were structured in 2-3 main cohorts, broadly corresponding to juvenile (placed around 13-19 mm CL) and adult individuals, with two main cohorts, one around 30 mm CL, and another at around 38 mm CL, this last not present in NCAT. Juvenile cohorts were only identified in the Balearic basin (NCAT, CCAT) and in Alicante (ALIC); they were not identified in the Alboran Sea.




Figure 6: Size frequency distributions of *Pasiphaea multidentata* per depth interval in the five best sampled geographic sectors. Total N: WALB = 337; EALB = 118; ALIC = 668; CCAT = 225; NCAT = 205)

### 1.3.2.4. Temperature and salinity

Occurrences of *P. multidentata* took place at bottom temperatures comprised between 12.91 and 13.43°C and salinities between 38.20 and 38.54. The temperature window was narrower for juveniles than for adults (Figure 10), while concerning salinities the range for juveniles was slightly broader. In particular, adults occurred (percentiles 25-75) at temperatures between 13.10 and 13.25°C and salinities of 38.44 and 38.50, while juveniles occurred at temperatures between 13.25 and 13.30 and salinities between 38.42 and 38.50.

## 1.4. Discussion

The information obtained during the studied series of trawl surveys has allowed the analysis of the distribution patterns of density and biomass of the two species of the genus

*Pasiphaea* present in the western Mediterranean throughout the southern and eastern coasts of the Iberian Peninsula down to depths of around 800 m. Furthermore their population size structure was also described. While the overall depth range distribution of both *P. sivado* and *P. multidentata* found in the present study fall within the ranges described in the literature for these species (Cartes 1993a, Koukouras *et al.* 2000, Fanelli *et al.* 2007), several patterns could be studied in detail given the broad geographic range of the surveys (Bertrand *et al.* 2002). This has allowed delimiting up to 10 smaller geographic sectors, based on the heterogeneous geomorphological and oceanographic characteristics of the western Mediterranean.

Most of the information available on distribution patterns, size structure and other population characteristics of the two species of *Pasiphaea* studied herein was restricted to the northernmost sampled area, the Catalan coasts (Abelló *et al.* 1988, 2002, Cartes 1993a,b,c, Company *et al.* 2001, 2003, Aguzzi *et al.* 2007, Ramirez-Llodra *et al.* 2007). Orsi-Relini and Pinca (1990) and Orsi-Relini and Relini (1990) respectively provided information on reproduction and trophic interaction in the Ligurian Sea (NE of the western Mediterranean basin). No biological or population information was to date available from other areas within the distribution range of the two species. Present results have shown the occurrence of marked differences in both distribution and population characteristics between the southwestern areas in the Alboran Sea, located in the Algerian basin of the western Mediterranean, and those present in the northeastern area, the Catalan Sea in the Balearic basin.

Thus, the geographic distribution of densities of *P. sivado* has firstly shown a heterogeneous distribution pattern, with two main nuclei, one in the Alboran Sea, and especially in its western area, and another along the Catalan coasts in the Northeast. Additionally, the bathymetric distribution in the West Alboran sector was markedly different to that found in the rest of geographic sectors. In this area, the species occurred in both much shallower and deeper waters reaching markedly higher densities at depths between 100 and 250 m, where this species was practically absent from the rest of sectors. The bathymetric distribution of *P. multidentata* reached also shallower waters in the western Alboran sector. This pattern found in both *Pasiphaea* species, although more marked in *P. sivado*, is in agreement with the occurrence of temporal upwellings located along the northwesternmost region of the Alboran Sea, in the area around Malaga (Vargas-

Yáñez and Sabatés 2007). These upwellings are responsible for the occurrence of high primary production cells which have been shown to allow delayed coupling of epibenthic shrimp populations with respect of plankton blooms associated to these upwelling cells (Fanelli and Cartes 2004). The occurrence of these upwelling cells close to the coast is a consequence of the interaction of the permanent strong eastward inflow of Atlantic water into the Mediterranean through the nearby Strait of Gibraltar with the steep continental slope of the southern Iberian continental margin, and intense local westerly winds (Millot 2005, Vargas-Yáñez and Sabatés 2007). Moreover, densities of *P. sivado* have been shown to be much higher than those of *P. multidentata* in accordance with Company *et al.* (2001) who suggested that the greater fecundity output of *P. sivado* would support its higher population densities when compared with *P. multidentata*.

Both juvenile and adult *P. sivado* are known to perform daily vertical migrations into the pelagic realm during night-time hours, while remaining on the bottom or close to it (epibenthic layer) during the day (Cartes1993c, Aguzzi *et al.* 2007), in this way, they can be considered as bentopelagic or nektobenthic species. Night-time daily migrations into the water column have also been reported for *P. multidentata*, but mainly restricted to juveniles, with the apparent exception of the adults (CL > 30 mm), which are assumed to perform epibenthic depth displacements (Cartes1993c, Cartes *et al.* 1993, Aguzzi *et al.* 2007). The day-time sampling schedule of the present trawl surveys would accordingly be adequate to sampling both juvenile and adults of the two species.




Figure 9: Logistic curves adjusted to the cumulative distribution by size of ovigerous female *Pasiphaea sivado* by geographic sector (NCAT: North Catalonia; CCAT: Central Catalonia; ALAC: Alacant; EALB: Eastern Alboran; WALB: Western Alboran)

The relationship between mean density and percentage occurrence by depth stratum allowed delimitating those depth strata showing high figures of both density and occurrence, which could be assumed to be the optimal depth ranges for the species. These were clearly located

between 300 and 500 m in *P. sivado*, and deeper than 500 m (down to the deepest sampled depth, 800 m) in *P. multidentata*. It must be emphasized that the sampling schedule clearly encompassed the whole bathymetric distribution range of *P. sivado* in the study area (Abelló *et al.* 1988, 2002, Cartes *et al.* 1994), whereas in *P. multidentata* it did not reach the deepest distribution of the species, which has been reported to occur down to 2261 m depth (Cartes 1993a) in the Mediterranean.

Both species showed a marked size increasing trend with depth, as shown in previous papers (Company *et al.* 2001), with juveniles being found in much shallower waters than adults. In *P. sivado*, populations in the Alboran Sea reached larger sizes than those present in the northwestern Mediterranean and in the intermediate Alicante sector. This implies that population dynamics are different in these areas, probably linked to the high productivity of the Alboran Sea (Fanelli and Cartes 2004). Recruitment in *P. sivado* was detected throughout the study area in the present surveys, performed in spring, in accordance with the main autumn-winter reproductive season of the species (Company *et al.* 2001), and was mainly located at depths shallower than 300-400 m. On the contrary, recruitment in *P. multidentata* was not present in the Algerian basin sectors (Alboran Sea and Alicante regions), while it was marked in the Balearic basin. The reproductive season in *P. multidentata*, from studies made in the Catalan Sea (Company *et al.* 2001, Ramirez-Llodra *et al.* 2007), is centered in late autumn – winter. The absence of recruitment in spring in the geographic sectors belonging to the Algerian basin may imply that seasonality of reproduction in this species significantly differs between the populations inhabiting the two basins.

The analysis of the environmental information available clearly detected differences between the occurrence patterns of the two species with respect to temperature and salinity on the bottom, as well as between juveniles and adults of the two species. Temperature and salinity windows (25-75 percentiles) of adults of the two species clearly overlapped, but the salinity range of *P. multidentata* was narrower than that of *P. sivado*, and much narrower than that shown by juveniles of the species, while the temperature range was contiguous but non-overlapping at a limiting temperature of 13.25 °C. In contrast, the temperature-salinity window of *P. sivado* juveniles was much larger, and widely overlapping, than that of the adults, which reached, however, higher salinities. This clearly showed that in both species, but particularly in *P. sivado*, juveniles showed a



wider thermohaline window, being able to cope with a wider variability in temperature and salinity conditions. The ability of juveniles to inhabit a wider range of salinity is common to other marine species, a physiological feature that has been mainly studied in coastal waters (McGaw and Naylor 1992, Abelló *et al.* 1997, Reid *et al.* 1997) related to the ability to cope with osmotic stress and the energetic balance involved. A close dependence on a restricted salinity range has also been shown in deep-sea and continental slope crustaceans (Guijarro *et al.* 2008, 2009, Cartes *et al.* 2011).


The differential distribution pattern between juveniles and adults of both *P. sivado* and *P. multidentata* (Company *et al.* 2001) could be related, as in other continental slope caridean shrimps (Company and Sardá 1997, Carbonell and Abelló 1998), to trophic resource partitioning between the two ontogenetic phases, which would allow a lesser degree of intraspecific competition. The dietary overlap between these species has been shown to be low due the different size spectra of their respective prey items (Cartes 1993b). Adults of *P. sivado* and juvenile *P. multidentata* share a similar size spectrum and they are both important food items to *P. multidentata* adults (Cartes 1993b). Size segregation, although implying a higher degree of osmotic regulation and its concomitant higher energetic expenses, would also be helpful to avoid intraspecific predation by juvenile *P. multidentata*.

The present study showed that in both *P. sivado* and *P. multidentata* Alboran Sea populations, their bathymetric distribution, density, maturity size, and population size structure were clearly differentiated from those in the Balearic basin. The populations in the intermediate Alicante sector showed more affinities with the Alboran Sea populations, especially so in *P. multidentata*, than with the northern populations, in agreement with their belonging to the Algerian basin.

# Chapter

# 2

Sexual dimorphism in the benthopelagic shrimp *Pasiphaea sivado* (Crustacea: Caridea: Pasiphaeidae)



# Sexual dimorphism in the benthopelagic shrimp *Pasiphaea sivado* (Crustacea: Caridea: Pasiphaeidae)

## 2.1 Introduction

Pasiphaeid shrimps are a group of benthopelagic and pelagic species that presently comprises a total of 98 species grouped in seven genera: *Alainopasiphaea* (2), *Eupasiphae* (4), *Glyphus* (1), *Leptochela* (15), *Parapasiphae* (4), *Pasiphaea* (67), and *Psathyrocaris* (5) (De Grave and Fransen 2011, Komai *et al.* 2012). In particular, within the genus *Pasiphaea* up to four morphologically-related species groups have been recognized, among them the *Pasiphaea sivado* group, to which a total of nine species belong (Hayashi, 1999, 2006). Species of this family are distributed along a wide latitudinal range, from tropical to arctic and subantarctic waters, mainly on and above continental slopes and in the open sea, including deep-sea habitats (Clarke and Holmes 1987, Kaartvedt *et al.* 1988, Gibbons *et al.* 1994, d'Udekem and d'Acoz 1999, Tavares and Cardoso 2006). Morphological features in this family, such as lateral compression of the body and pleopods, are considered to be the result of selective pressures on adaptation to the pelagic environment (Cartes *et al.* 1993a, Johnsen 2005, Aguzzi *et al.* 2007). Notwithstanding the large amount of systematics and faunistics work on this family (e.g. Stephensen 1923, Zariquiey Álvarez 1957, Iwasaki 1990, Hayashi 1999, Tavares and Cardoso 2006) there are still few detailed studies on biological aspects of species of the family, and most of them refer mainly to the European species *P. sivado* and *P. multidentata* (Cartes 1993, Company and Sardá 2000, Frank 2000, Company *et al.* 2001, Ramirez-Llodra *et al.* 2007), but also to Pacific Ocean species such as *Pasiphaea japonica* and *Leptochela gracilis* (Nanjo 2007, Oh *et al.* 2006).

No reference is made to the occurrence of macroscopic differences in shape in relation to gender in this family (Stephensen 1923, Zariquiey Álvarez 1968, Iwasaki 1990, Hayashi 1999, Tavares and Cardoso 2006), except for the noted occurrence of differences in pleopod shape in *Pasiphaea japonica* (Doi 1975, Nanjo and Ohtomi 2009). Concerning size dimorphism, males have

been reported to be slightly larger than females in *P. sivado*, and slightly smaller in *P. multidentata* (Company *et al.* 2001), based in overall size frequency distributions, but sexual dimorphism was so far considered to be not significant in neither *P. sivado* nor *P. multidentata*.

After careful macroscopic examination of previously sexed individuals of *P. sivado* (based on the presence of appendix masculina in the second pleopods of males), marked differences in pleopod shape were observed between adult male and female individuals of the species. We herein aim to report the occurrence of this morphological dimorphism in *P. sivado*, as well as to quantify it, based in both traditional and geometric morphometrics, which may help to further evidence and characterize the occurrence of this dimorphism (Rufino *et al.* 2006). Size-related population structure and sex-ratio are also analyzed in relation to geographic and bathymetric characteristics.

## 2.2 Material and Methods

### 2.2.1 Pleopod morphology

Ten adult specimens of *Pasiphaea sivado* (five males and five females) were selected from a sample taken by demersal trawling in the western Mediterranean Sea (Alboran Sea: 36.4908 N, -4.2885 W) at 424-428 m depth during daytime on 17/05/2010. The haul was performed on board R/V “Cornide de Saavedra” at a speed of 3 knots during 1h using a demersal GOC-73 otter trawl with a 2.5-3.0 m vertical opening and a codend stretched mesh size of 20 mm (Fiorentini *et al.* 1999). The cruise was part of the MEDITS\_ES project (Bertrand *et al.* 2000, 2002).

The selected specimens had a cephalothorax length (CL) > 18 mm and were clearly morphologically considered as adults (Company *et al.* 2001; present observations). CL was measured using a calliper rule with a precision of 0.01 mm. Sizes of the examined specimens ranged between 18.8–23.5 mm CL in females, and between 21.6-24.0 mm CL in males. The individuals were sexed by checking the presence/absence of appendix masculina on the endopod of the second pleopod, as usual in caridean shrimps.

In a first step, the five pleopod protopods of one of the males and one of the females of similar size were figured in order to graphically show the a priori differences observed between adult males and females. Then, with the aid of a “Leica MZ12” magnifier provided with ocular micrometer, the length and maximum width of all right pleopod protopods of all ten specimens studied were measured to obtain a first index of pleopod shape based on the ratio between length and width of this segment.

Pictures were also taken of all right second pleopods (external view) of all five males and females studied using the same magnifier (which has an integrated video camera Leica DC100 and DC200 digital imaging systems) to further assess the occurrence of sexual dimorphism using geometric morphometric techniques. The first landmark was placed on the dorsal maximum convex curvature of the protopod, the second and fourth were placed at the anterior maximum curvature where the exopod and endopod insert on the protopod, the third at the maximum concavity between these two landmarks, the fifth at the maximum perpendicular width of the protopod in relation to the first landmark, the sixth at the ventral maximum curvature in the posterior part of the protopod and the seventh at the maximum concave curvature on the posterior dorsal margin of the protopod (Figure 1).




Figure 1: Landmark positions in the digital images obtained with stereoscope using a 16x magnifier with optical zoom. The width and length of each *Pasiphaea sivado* pleopod protopod were measured from the scalar distance (in mm) between the landmarks 1-5 and 4-6, respectively.

Using Tps.Dig2 v2.16 software (Rohlf 2010a), the  $x,y$  coordinates of the landmark dataset were obtained. Generalized procrustes superimposition analysis (GPA) was performed to align and to eliminate scale and orientation distortions before proceeding to shape analysis using tpsRelw v1.49 software (Rohlf, 2010b). Statistical analysis of shape variance was assessed using the MorphoJ software (Klingenberg 2011). Then procrustes ANOVA was applied, which is a two

factorial ANOVA that performs analysis of both size (univariate) and shape (multivariate). This procedure therefore is a useful method to discover localized variation (Klingenberg and McIntyre 1998). Finally a principal component analysis (PCA) was applied to quantify the contribution of the principal factor of landmark displacement that explains the shape variance between genders. This PCA was run using the covariance matrix generated from shape variation components, which were previously derived from the coordinates landmark dataset, pooled within-group with sex as covariate (Viscosi and Cardini 2011).

### 2.2.2 Gender differences in size. Sex-ratio

In order to assess whether there occurred significant differences between males and females concerning overall size dimorphism, data were obtained from a total of 4010 *P. sivado* individuals. The samples were collected during a series of demersal fisheries research cruises (MEDITS\_ES) between 2004 and 2011 at depths from around 40 to 800 m along the Mediterranean coasts of the Iberian Peninsula (Bertrand *et al.* 2002). In these cruises, *P. sivado* was only collected at depths between 150-750 m.

All specimens were sexed as described above and males were classified as juveniles when the appendix masculina was not fully formed, which usually appeared as small buds. This was evident in individuals with sizes ranging between 9 and 16 mm CL, which were therefore classified as juvenile males. However, the proportion of individuals showing appendix masculina buds was very small at sizes smaller than 11-12 mm CL. This implies that based only in this character the correct assignation of gender was not possible at these small sizes, there probably being a larger proportion of males with still undeveloped appendix masculina. Therefore, sex-ratio analysis was restricted to individuals exceeding 12 mm CL.

Sex-ratio, as proportion of males, was calculated within each 1 mm CL size class, separately for samples taken in the Alboran Sea (the westernmost basin of the Mediterranean Sea), the intermediate Alacant region (between Cape Palos and Cape La Nao) and the Catalan Sea (in the NW Mediterranean). These sectors were established according to previous results (Simão *et al.*

submitted; see Chapter 1 on population characteristics of pasiphaeid shrimps) in which differences in size at maturity were observed between Alboran Sea populations and the rest of sampled populations along the Iberian Peninsula Mediterranean coasts. Sex-ratio was not calculated in size classes with less than 25 individuals. The significance of the deviation from an expected 1:1 sex-ratio was calculated using a heterogeneity G-test (Sokal and Rohlf 1981, Abelló *et al.* 1990). The relationship of sex-ratio with depth was also calculated; only samples  $\geq 25$  sexed individuals larger than 15 mm CL were considered for this analysis; this cutting size was chosen to ensure that sex-ratio was calculated over adult specimens to minimize error in gender assessment (see above), and also to encompass the whole adult cohort as visually assessed through size frequency distributions.

In order to verify whether size frequency distributions (SFD) differed between geographic populations and between depth strata, the Kolmogorov-Smirnov test within VITMAN fisheries software (Leonart and Salat 1992) was used. Combinations of geographic area and depth strata resulting in less than 100 individuals measured (i.e. at both extremes of the bathymetric range of the species in each area) were not used in the analyses.

## 2.3 Results

### 2.3.1 Pleopod morphology

The difference in protopod morphology between genders of *Pasiphaea sivado* can be easily identified by sight (Figure 2). Overall, in all five pleopod pairs, adult female protopods were clearly much thinner than those in adult males. Additionally, the anterior edge of the male protopods was clearly much more convex than in females, while the posterior edge was approximately straight in both males and females. As shown in Figure 3, the ratio between pleopod protopod length and width (L/W), which can be considered a protopod thinness index, was much higher in females than in males, thus characterizing the thinner, rectangular, elongated shape in females versus the more robust shape present in males. A significant difference between genders was obtained for all five pleopod pairs (Table 1). Additionally, and concerning overall morphology, no setae were present on

the anterior edge of the first pleopod protopod in any of the adult males examined. Also, in the rest of pleopods, anterior setae were much scarcer in males than in females.

The use of geometric morphometrics techniques further evidenced the differences between genders in pleopod protopod shape. Thus, both *P. sivado* males and females grouped as two well-differentiated clusters when analyzing the shape of their second pleopod protopods (Figure 4a). The overall consensus shape (Figure 4b) of the pleopod protopods, which is based on the

geometric mean position of each landmark, clearly differentiated between the thinner, elongated shape of the females versus the heavier found in males. Principal Component Analysis showed that the first principal component (PC1), which accounted for 56.1% of the total shape variability,




Figure 3: Mean pleopod (PL) protopod thinness index (length/width) ( $\pm$ SEM) of female (white circles) and male (black circles) *Pasiphaea sivado*.




Figure 2: Lateral view of the protopods of each five pleopods of a 22.7 mm CL *Pasiphaea sivado* female and of a 22.6 mm CL male.

clearly differentiated males from females. The shape variance between genders explained by the PC1 was significantly different (ANOVA  $F_{10,80} = 27.24$   $p < 0.0001$ ) and was not affected by the centroid size effect (ANOVA  $F_{1,8} = 1.87$   $p = 0.209$ ). The next five principal components explained 16.5%, 10.4%, 6.2%, 3.5% and 1.26% of the total shape variance, respectively.



Table 1: Number of individuals examined (N), mean, standard deviation, maximum and minimum length (L) and width (W) of the pleopod protopod (PL1-5), and mean ratio L/W of adult *Pasiphaea sivado* females (CL<sub>range</sub> = 18.8–23.5 mm) and males (CL<sub>range</sub> = 21.6–24.0 mm). Significance of differences of the L/W ratio between genders is also shown.

Females	Length					Width				Mean L/W ratio	t Test p
	N	Mean	+SD	Max	Min	Mean	+SD	Max	Min		
PL 1	4	4.65	0.38	5.0	4.1	1.25	0.14	1.4	1.1	3.73	p < 0.05
PL 2	5	5.13	0.47	5.8	4.7	1.73	0.38	2.3	1.3	3.07	p < 0.02
PL 3	5	4.97	0.38	5.5	4.5	1.79	0.41	2.4	1.4	2.90	p < 0.05
PL 4	5	4.73	0.43	5.2	4.2	1.99	0.38	2.6	1.5	2.44	p < 0.05
PL 5	4	4.19	0.40	4.5	3.6	1.73	0.20	1.9	1.4	2.42	p < 0.001
Males	Length					Width				Mean L/W ratio	
N	Mean	+SD	Max	Min	Mean	+SD	Max	Min			
PL 1	5	4.89	0.22	5.2	4.6	1.79	0.22	2.0	1.4	2.76	
PL 2	5	5.09	0.09	5.2	5.0	2.33	0.22	2.5	2.0	2.21	
PL 3	5	5.08	0.36	5.6	4.8	2.52	0.16	2.8	2.4	2.01	
PL 4	5	4.86	0.30	5.3	4.5	2.60	0.09	2.7	2.5	1.87	
PL 5	5	4.16	0.23	4.4	3.8	2.46	0.16	2.7	2.3	1.70	




Figure 4: Shape variation analysis (a) and thin plate consensus shapes (b) for adults *Pasiphaea sivado*: females (white circles) and males (black circles). The contribution percentages of the two principal components are presented in the correspondent axis.

### 2.3.2 Comparison of sizes and size-related sex-ratio among geographic areas

The overall size frequency distributions of sexed individuals by geographic area (Figure 5) were biased towards adult individuals in all populations. The Alboran Sea population showed a higher proportion of individuals in the largest sizes, when compared with those obtained in the other areas: Alacant and Catalan Sea. The comparison between male and female SFDs within each

area showed that they were not significantly different in any case (main diagonal in Table 2). Comparison of male or female SFDs among areas also showed non-significant values (Table 2).

The analysis of sex-ratio by size class showed, in all sectors, that females were dominant at sizes smaller than around 18-19 mm CL, while the proportion of males was significantly higher at larger sizes, therefore evidencing

Table 2: Comparison of SFD between geographic populations (Kolmogorov-Smirnov test). The  $p$  values for the comparison between females are presented above the diagonal, those between males below the diagonal, and those comparing males and females within each population along the main diagonal.

		Kolmogorov-Smirnov (p value)			
		Alboran	Alacant	Catalan	
Alboran		0.869	0.308	0.537	♀
Alacant		0.522	0.263	1.000	
Catalan	♂	0.182	1.000	0.243	

the occurrence of a slight sexual size dimorphism in *P. sivado* (Figure 5). When paying especial attention to the precise size at which the dominance of males occurred, a decreasing gradient was found along the study area, with the proportion of males being significantly higher at sizes larger than 21 mm CL in the Alboran Sea, 20 mm in Alacant and 19 mm in the Catalan Sea (Figure 5).




Figure 5: Size frequency distributions by sex in *Pasiphaea sivado* from the Alboran Sea, Alacant coast and Catalan Sea geographic sectors. The Sex-ratio, as proportion of males is represented by circles, with indication of values showing a significant ( $p < 0.05$ , dark circles) or non-significant (n.s., light circles) deviation from an expected 1:1 ratio. The total number of individuals analysed by geographic area is presented.

### 2.3.3 Size structure with depth by geographic area

When the shape of the SFDs was analyzed by depth, a high frequency of juveniles in relation to adults was evident in the shallowest depth stratum (200-300 m), especially so in the Alboran Sea (Figure 6). In this area, the population showed a marked SFD displacement towards larger sizes with increasing depth; no noticeable variation in SFD displacement was evident in the Alacant and Catalan Sea populations (Figure 6 and Table 3). The comparison of the SFDs with the Kolmogorov-

Smirnov test showed the occurrence of significant differences only in the Alboran Sea population, in both males and females (Table 4). No significant differences of SFD variation with depth were detected in the Alacant and Catalan Sea populations.




Figure 6: Size frequency distributions of male and female *Pasiphaea sivado* by 100 m depth strata. Strata with less than 100 measured individuals were not taken into account. The number of measured individuals by gender (N) is shown.

Table 3: Comparison of median sizes among 100 m depth strata within each geographic population. The obtained Mann-Whitney *p* values for the comparison between females are presented above the diagonal, between males below the diagonal and between genders within each depth stratum along the diagonal.

	Median CL ♂	Depth Strata (m)	200-300	300-400	400-500	500-600	Median CL ♀
	Alboran Sea	17.0	200-300	0.301	0.000	0.000	
	19.4	300-400	0.000	0.000	0.000	0.000	18.4
	21.0	400-500	0.000	0.000	0.000	0.000	19.9
	22.3	500-600	♂ 0.000	0.000	0.000	0.000	22.4
Alacant Sea	19.9	200-300	0.001	0.294	0.970	-	♀ 17.8
	19.0	300-400	0.298	0.004	0.083	-	18.1
	19.7	400-500	0.626	0.008	0.000	-	17.6
	20.0	500-600	♂ -	-	-	-	20.0
Catalan Sea	21.8	200-300	-	-	-	-	♀ 12.9
	19.1	300-400	-	0.000	0.412	0.030	17.8
	19.4	400-500	-	0.036	0.000	0.959	17.9
	19.8	500-600	♂ -	0.001	0.023	0.000	17.8

55

Table 4: Comparison of size frequency distributions by 100 m depth strata within each geographic population. The Kolmogorov-Smirnov *p* values for the comparison between females are presented above the diagonal, between males below the diagonal, and between genders along the diagonal (values in italics); significant values are shown in bold.

		200-300	300-400	400-500	500-600	
		Alboran Sea	200-300	<b>1.000</b>	0.650	
	300-400	0.278	<b>0.931</b>	0.685	0.013	
	400-500	0.001	0.272	<b>0.517</b>	0.184	
	500-600	♂ 0.000	0.004	0.171	<b>0.830</b>	
Alacant coast	200-300	<b>0.105</b>	0.978	1.000	-	♀
	300-400	0.957	<b>0.687</b>	1.000	-	
	400-500	0.997	1.000	<b>0.208</b>	-	
	500-600	♂ -	-	-	-	
Catalan Sea	200-300	-	-	-	-	♀
	300-400	-	<b>0.240</b>	1.000	0.772	
	400-500	-	1.000	<b>0.261</b>	0.992	
	500-600	♂ -	0.736	0.985	<b>0.199</b>	

### 2.3.4 Sex-ratio with depth

A total of 84 samples were used to analyze the sex ratio across the bathymetric distribution of the species (Figure 7). Most of the samples (70.2%) did not show a significant deviation from a 1:1 sex-ratio. However, significant sex segregation was found in the rest of samples, with 11.9% of them being biased toward females and 17.9% towards males. The highest proportion of significant deviation toward females was found at the shallowest extreme of the bathymetric distribution of the species but also in the deepest strata. On the contrary, samples with a significantly higher proportion of males were located in the central depth strata. Overall, no significant tendency of gender segregation across the bathymetric distribution was found ( $r=0.0803$ ,  $p=0.4677$ ).




Figure 7: Sex-ratio distribution of *Pasiphaea sivado* as function of depth by geographic area: Alboran Sea (circles), Alacant coast (squares) and Catalan Sea (triangles). The full symbols correspond to samples significantly different to the expected 1:1 sex-ratio, while the empty circles indicate no significant difference in sex-ratio.

## 2.4 Discussion

### 2.4.1 Pleopod morphology

The occurrence of sexual dimorphism in pleopod shape was detected in *Pasiphaea sivado* by both the traditional linear measurement method and by the geometric morphometric analysis. An important, useful application of the present observations and description of the occurrence of sexual dimorphism in pleopod shape in adult *P. sivado* is, from now on, the ability to quickly assess the gender of individuals when performing population biology studies.

A similar sexual dimorphism in pleopod shape has also been reported in *Pasiphaea japonica*, but it also seems to occur in *Alainopasiphaea nudipeda* as evidenced by figures 1 and 2d in Hayashi (1999), as well as in *Pasiphaea semispinosa* (authors personal observations based in specimens deposited in collections: CBR at ICM-CSIC in Barcelona and CCDE at IEO-Cádiz). Other personal observations, not yet quantified, on the congeneric species *Pasiphaea multidentata* - a species that coexists with *P. sivado* across the Mediterranean Sea (Company *et al.* 2001, Abelló *et al.* 2002)- indicate that this species does not show such dimorphism.

The phylogenetic relationships within the family *Pasiphaeidae* are far from being properly understood (Hayashi 1999, 2006, Matzen da Silva *et al.* 2011; see also the section on genetics in this thesis). Based on present findings, the occurrence of sexual dimorphism in pleopod shape may well have a significant role in determining phylogenetic relationships within the family *Pasiphaeidae*, in particular within the *Pasiphaea sivado* group (Hayashi 1999, 2006), since both *P. sivado* and *P. japonica* (who also belongs to this group) have been shown to present this morphological characteristic (Doi 1975, Nanjo and Ohtomi 2009). Not enough information is available concerning other species in the genus or the family.

### 2.4.2 Size dimorphism and sex ratio

The present study provides also evidence on the occurrence of significant size dimorphism in *P. sivado*, a fact previously unreported in this species. In all geographic areas and depth strata, males were significantly larger than females. *P. sivado* presented a marked size structure pattern where females accumulated in size classes smaller than around 20 mm CL while males dominated in the largest size classes. Indeed, no strong size dimorphism is usually found in species with pelagic habits, while strong differences in size usually occur in strictly benthic/epibenthic species such as brachyuran crabs or lobsters.

In *P. sivado*, sex-ratio as a function of size clearly suggests the occurrence of differential growth rates between the genders, as reported in many species of crustaceans (Wenner 1972, Abelló and Cartes 1992, Guerao *et al.* 1994, Ben Mariem 2004, Oh *et al.* 2002, Kim 2005, Kevrekidis and Thessalou-Legaki 2006, Oh *et al.* 2006), a fact that is usually attributed to a higher

investment of females on reproduction rather than on growth. This pattern appears to follow the so-called anomalous pattern described by Wenner (1972), in which no differences in sex-ratio would occur in juveniles, then a predominance of females would take place due to a lower female growth rate, with males finally accumulating their frequency in the largest sizes. Several explanations may account for this pattern, with different growth rates between the genders from the puberty moult or size at maturity being the most likely (Ben Mariem 2004, Fantucci *et al.* 2009). Some biological characters estimated by Company *et al.* (2001) for *P. sivado* in the Catalan Sea support the sex-ratio pattern with size reported in the present study. In that work males showed a faster growth rate ( $k=0.62$ ), a larger asymptotic size ( $L_{\infty}=27.5$  mm) and a higher growth performance index ( $\Phi=0.13$ ) than females (0.55, 26 mm and 0.06, respectively). Reproductive aspects of the species such as the continuous reproduction throughout the year and the higher volumetric index of females (2.92) compared to males (2.76) (Company *et al.* 2001) also suggest the smaller amount of energy investment in size growth by *P. sivado* females in relation to males. Another factor that may also limit the growth in females is the moulting delay of ovigerous females in order to avoid loss of attached embryos, a phenomenon characteristic of crustacean species with egg-carrying biological features (Guerao *et al.* 1994, Oh *et al.* 1999, Viegas *et al.* 2007).

Differential mortality rates between the genders may also contribute to reaching the observed pattern of sex-ratio variation with size (Abelló and Macpherson 1992). It is however so far not possible to assess the occurrence of different mortality rates between the sexes given the lack of behavioural observations on *P. sivado*. The size at which differences in the sex-ratio become significant has also been suggested to be an indirect indicator of the size at puberty moult and/or maturity size/age of the species (Abelló *et al.* 1990). In the present study the precise size at which males outnumbered females progressively decreased from the Alboran to the Catalan Sea through the Alacant area, corroborating the size and size at maturity pattern observed in Chapter 1, and suggesting the occurrence of environmental and ecological processes affecting growth rates.

Overall, no significant gender segregation in *P. sivado* populations as a function of depth was verified, suggesting that this species does not perform differential migration between sexes over the sea bottom. In the Catalan Sea the absence of gender segregation by depth was also reported by Company *et al.* (2001) throughout the year, which could be related to the gregarious behaviour of

macroplankton/micronekton organisms (McManus and Woodson 2012). During the day time, *P. sivado* inhabits the epi-/suprabenthic environment over the upper and medium slope, with high densities recorded when sampled by demersal trawl nets during fishery research surveys (see Chapter 1) (Fariña *et al.* 1997; Fanelli *et al.* 2007; Abad *et al.* 2007; Cartes *et al.* 2007b; García-Rodríguez *et al.* 2011), while during the night the species has been reported to migrate several hundred meters up the water column to waters near the surface to feed on plankton (Froggia and Giannini 1982, Cartes 1993a, Aguzzi *et al.* 2007, see also Chapter 4 on vertical migrations in this thesis).

Controversially, a relatively high number of samples showed significantly deviated sex ratios, both towards a higher proportion of males and of females. In the absence of direct observations, sex-ratio deviated groups, indicative of aggregative behaviour within genders, are suggestive of differential reproductive behaviour and/or selective feeding in relation with the energetics and quality needs depending on gonadal developmental stages. Thus, an increase in stomach fullness and dietary energy intake has been recorded in pre-reproductive females of the deep-sea shrimp *Aristeus antennatus*, parallel to a significant increase of the gonado-somatic index and fecundity (Cartes *et al.* 2008). Differences in diet composition between genders and maturity stages have also been found in other shrimps such as *Plesionika heterocarpus* and *Plesionika martia* (Fanelli and Cartes 2008). Aggregative behaviour within genders has also been related to other causes, such as the need to look for appropriate habitat for moulting (Reid *et al.* 1994, 1997, Sampedro and González-Gurriarán 2004) or for hatching (i.e females look for adequate sites to improve dispersal and survival of recently hatched larvae) (Hicks 1985; Zeng and Naylor 1997; Carr *et al.* 2004).

In attention to the concept of operational sex ratio (OSR), defined as the proportion of receptive females to sexually active males (Correa and Thiel 2003), the results of the present study suggest that the populations of *P. sivado* may overall be close to the OSR. The histograms of Figure 6 show that the overall shape of the SFDs of males and females is very similar, but males are displaced towards the right (largest sizes) by just one size class. If we quantify the total adult males and females bigger than the size class precedent of the one where the sex ratio starts to be biased towards males, we will find a sex-ratio very similar to the equilibrium. Even with the existence of a




size sexual dimorphism or some samples with biased sex-ratio, no overall sex segregation at population level was found.

# Chapter

# 3

Population genetics and phylogeny of pasiphaeid shrimps from the western Mediterranean Sea



# Population genetics and phylogeny of pasiphaeid shrimps from the western Mediterranean Sea

## 3.1. Introduction

Pasiphaeid shrimps are a worldwide distributed family of pelagic and benthopelagic shrimps. They comprise so far a total of 97 species grouped in seven genera: *AlainoPasiphaea* (2), *Eupasiphae* (4), *Glyphus* (1), *Leptochela* (15), *Parapasiphae* (4), *Pasiphaea* (66), and *Psathyrocaris* (5) (De Grave & Fransen, 2011). Most pasiphaeid species are benthopelagic species occurring on the continental slope and show important diel vertical movements, being present on the bottom or benthic boundary layer during the day, while rising several hundred meters into the water column during the night (Aguzzi *et al.* 2007, Naylor 2010). The wide geographical and bathymetrical distribution as well as their important role in linking the pelagic and benthic trophic webs confer to the *Pasiphaeidae* a wide presence on marine community and trophic studies (Fashan and Foxton 1979, Cartes 1993, Vereshchaka 1995, Company & Sardá 2000, Frank 2000, Company *et al.* 2001, Oh *et al.* 2006, Ramirez-Llodra *et al.* 2007, Nanjo and Otomi 2009). However there still lacks substantial information about the population biology and genetic dynamics of this family.

In the Mediterranean Sea the Pasiphaeoidea superfamily is represented by just two, but very common species *Pasiphaea sivado* and *Pasiphaea multidentata*. *Pasiphaea sivado* is a widely distributed species along the Mediterranean Sea, northeastern Atlantic and western African coasts being commonly sampled by trawling during demersal fisheries research surveys (Politou *et al.* 2000, Monteiro *et al.* 2001, Ungaro *et al.* 2005, Fanelli *et al.* 2007, García-Rodríguez *et al.* 2011). The species has a marked diel vertical migration occurring up to the first 50 m below the surface during the night, while occurring on the upper and middle slope bottom during the day (see Chapter 3 of this thesis), where it inhabits preferentially on bottom depths of around 350-500 m (Abelló *et al.* 1988, 2002, Company *et al.* 2001, Muñoz *et al.* 2012). In the Mediterranean *P. sivado*

reproduces continuously around the year, a pattern common in tropical and shallow water species (Orsi-Relini and Pinca 1990, Company *et al.* 2001).

*Pasiphaea multidentata*, on the contrary, is a much scarcer species, with a much deeper habitat (deeper than 500 m) (Cartes *et al.* 1993, Biagi *et al.* 2002, Company *et al.* 2004) and more linked to the benthic environment (Cartes 1993, Cartes *et al.* 2007, 2009) than *P. sivado*. Its pelagic behaviour has been mainly reported for juvenile individuals, which show a shallower distribution than adults (Company *et al.* 2001, Aguzzi *et al.* 2007). However adults are also able to perform wide vertical movements but their distance range is more restricted to the benthic boundary layer (BBL), being absent in pelagic strata shallower than 400 m depth (see the Chapter 3 of this thesis). Moreover the species presents seasonal reproduction placed during the boreal autumn and winter, producing fewer but bigger yolked eggs than *P. sivado* (Company *et al.* 2001), in the Mediterranean Sea, indicating that *P. multidentata* is a typical demersal / temperate species (Ramirez-Llodra *et al.* 2007).

Considering the daily night occurrence of both species in the pelagic realm and their relatively wide vertical distribution, their populations would a priori tend to present panmitic characteristics. However, the pelagic environment is not a homogeneous system. Different water masses, with their own temperature, salinity and density characteristics are present in the ocean basins and species can evolutionarily evolve to preferentially associate with different physico-chemistry characteristics (Cartes *et al.* 2011). Also, the occurrence of geographic and geomorphologic barriers may prevent population mixing and dispersion (Palero *et al.* 2008, 2011, García-Merchán *et al.* 2012). Strong oceanographic currents, usually associated to water mass fronts, may also play both a population restriction role across the front, and a dispersion effect along their preferential direction (Patarnello *et al.* 2007, Carreras *et al.* 2011, Schunter *et al.* 2011).

In order to investigate the genetic characteristics along the Atlantic-Mediterranean transition, the two congeneric species of pelagic shrimps, *Pasiphaea sivado* and *Pasiphaea multidentata*, were selected to verify the occurrence of any population genetic differentiation between eastern Atlantic and Mediterranean populations which could be linked to oceanographic

and/or geographic characteristics. Intraspecific population variability is also linked to interspecific variability in this wide group of species.

The precise placement of the family Pasiphaeidae within the Caridea, as well as the phylogenetic relationships among the genera and species within the family are not yet fully understood (Christoffersen 1990, Bracken *et al.* 2009, Chan *et al.* 2010, Matzen da Silva *et al.* 2011, Li *et al.* 2011). Recent taxonomic studies of Hayashi (1999) proposed differentiation of the *Pasiphaea sivado* group, composed by *P. sivado* (Risso 1816) and nine other species from the rest of the *Pasiphaeidae* family which present branchial reduction (see Hayashi 1999, 2006). Detailed studies on phylogeny and molecular aspects of *Pasiphaeidae* are still missing. In this way the selection of these two congeneric species may also allow to perform phylogenetic studies. We therefore aimed to present additional information on the genetic phylogeny of the Atlantic-Mediterranean species of *Pasiphaea*, taking into account the populations inhabiting both the Mediterranean and eastern Atlantic basins.

## 3.2. Material and Methods

### 3.2.1. Sampling

The material analysed in the present work was obtained from two demersal fishery research surveys run by the Instituto Español de Oceanografía (IEO). The Mediterranean Sea material was sampled across the Spanish coast during the 2009 and 2010 MEDITS\_ES surveys (Bertrand *et al.* 2002) while the Atlantic Ocean material was obtained from the 2009 ARSA survey (Lopez de la Rosa 1997, Silva *et al.* 2011) in the Gulf of Cadiz. The Mediterranean samples were obtained from three previously defined zones established according to previous oceanographic and biogeographic results (Abelló *et al.* 2002, Rufino *et al.* 2005, Palero *et al.* 2009, García-Merchán 2012): Northern Alboran Sea (between the Strait of Gibraltar and Cape Gata), Alacant coast (from Cape Palos to Cape La Nao), and Catalan Sea (from Barcelona to Palamós).

On board, after each haul, the species were identified according to Zariquiey Álvarez (1968). The number of sampled individuals by species and geographic sector is shown in Table 1. For each individual a section of the abdominal muscle (of approximately 20 mg) was preserved in absolute ethanol at 4 °C for posterior genetic analysis in the laboratory on land.

Table 1: Sampled areas and number of COI sequences used for the intra-population variability analysis. Five sequences from the Gulf of Cadiz, and those from western Scotland and Gulf of St. Lawrence were obtained from GenBank.

Sampled area	Sea	<i>P. sivado</i>	<i>P. multidentata</i>
Catalan Sea	W Mediterranean	16	3
Alacant coast	W Mediterranean	6	-
Alboran Sea	W Mediterranean	16	4
Gulf of Cadiz	NE Atlantic	11	-
Western Scotland	NE Atlantic	-	2
Gulf of St. Lawrence	NW Atlantic	-	3
Total		49	12

### 3.2.2. DNA isolation, amplification and sequencing

The mitochondrial DNA was extracted by the Estoup *et al.* (1996) method that consists in incubating the dried tissue (ca. 2 µg) in a mix of 200 µl of Chelex 10% with 2 µl of K proteinase (20 mg/µl). The minimum incubation time was 3 hours, blending the samples each 30 minutes. Fragments of cytochrome oxidase I (COI) gene from the mitochondrial DNA were amplified using the universal primers designed by Folmer *et al.* (1994): LCOI-1490 (GGT CAA CAA ATC ATA AAG ATA TTG) and HCOI-2198 (TAA ACT TCA GGG TGA CCA AAA AAT CA). The amplification reactions were performed using 40 ng of mitochondrial DNA in a solution of 1U of polimerase Taq (Amersham), 1x PCR buffer (Amersham), 0.2 mM of each primer, 0.12 mM of dNTPs and 1.5 mM of MgCl<sub>2</sub> in a final volume of 13 µl. After the DNA denaturalization at 94°C during 4 minutes, the PCR program starts 36 cycles of incubations: 1 minute at 94°C, 1 minute at 54°C, 1 minute at 72 °C and a final extension of 7 minutes at 72°C. Two microliters of the amplification product were purified using the EXO-SAP enzymatic method with 0.34 µl of Exonuclease I (ThermoScientific 20U/µl) and 0.66 µl of S.A.P. (Shrimp Alkaline Phosphatase) (PROMEGA 1U/µl) by two incubation steps:

first 15 minutes at 37°C, then another 15 minutes at 80°C. The sequences were obtained through the sequencing kit Big-Dye v3.1 (Applied Biosystems) by the assumptions of the automatic sequencing ABI PRISM® 3770 from the technical and scientific service of the Universitat de Barcelona: 0.75 µl of BigDye, 1.25 µl of BigDye's Buffer, 1.0 µl of the forward primer (0.2 µM), 3.0 µl of the purified product in final volume of 10 µl.

Prior to perform population and phylogenetic analysis the obtained sequences of COI fragments were manually checked for ambiguous base calls, assembled and aligned in BioEdit v7.0.1 software (Hall 1999). In the DnaSP v5 (Librado and Rozas 2009) software the aminoacid translation of COI fragments were examined to ensure that no gaps or stop codons were present.

### 3.2.3. Intraspecific genetic diversity

To investigate the effect of the Gibraltar Strait in the genetic population structure of the benthopelagic *Pasiphaeidae* dwelling across the Atlantic-Mediterranean gradient, a total of 44 *Pasiphaea sivado* individuals from four different populations were sequenced: Gulf of Cadiz in the eastern central Atlantic, and Alboran Sea, Alacant, and Catalan, coasts along the western Mediterranean Sea (Figure 1). Five additional sequences from GenBank were added to our data from Atlantic waters off Algarve, Portugal, in the western part of the Gulf of Cadiz (GenBank access numbers: JQ306261-65) (Matzen da Silva *et al.* 2011).



Figure 1: Atlanto-Mediterranean transition showing the sampling locations of *P. sivado* (circles) and *P. multidentata* (squares): Gulf of Cadiz, Alboran Sea, Alacant coast and Catalan Sea.

Samples of the scarcer *Pasiphaea multidentata* were obtained from the two Mediterranean extremes of the sampled area, namely Catalan Sea and Alboran Sea (seven sequences); two

additional sequences from the NE Atlantic (Scotland) (Matzen da Silva *et al.* 2011), and three more from the NW Atlantic (Gulf of St. Lawrence) (Radulovici *et al.* 2009) were obtained from Genbank (access numbers: JQ305977-78 and FJ581853-55, respectively). No *P. multidentata* could be sampled in any of the surveys performed in the Gulf of Cadiz.

The haplotype and nucleotide diversities, as well as the gene flow measurements ( $\Gamma_{st}$ ,  $K_{st}$  and  $S_{nn}$ ) and the neutral tests (Tajimas's  $D$  and Ramos-Onsins and Rozas,  $R^2$ ) were estimated. The  $S_{nn}$  statistic was obtained by Hudson's method (2000). The haplotype networks were constructed using the "median joining network" method by using the Network v.4.5.1.6 (Fluxus Thecnology) program.

### 3.2.4. Phylogenetic analysis

To provide an approach to the phylogenetic reconstruction of the *Pasiphaea* genus we used one sequence for each known haplotype of the genus, taken from our own present data (from the above population genetics analysis) and the information available in GeneBank (25/04/2013): *Pasiphaea hoplocerca*, *Pasiphaea multidentata*, *Pasiphaea pacifica*, *Pasiphaea sivado* and *Pasiphaea tarda*. As outgroup, the closely related caridean shrimps *Acanthephyra pelagica*, *Alvinocaris muricola* and *Glyphocrangon neglecta* (Li *et al.* 2011) were selected, with sequences obtained either from GeneBank, or from the University of Barcelona Genetics Department Database (Table 2). The caridean species were selected as outgroup to avoid the noise due to the high substitution rates of the COI fragment compared to another genes (Xia *et al.* 2013). In those cases when due to methodological requisites only one outgroup could be selected *G. neglecta* was used as such. The nucleotide substitution saturation and transitions over transversions ratios were analysed by using the DAMBE5 software (Xia *et al.* 2013).

To elucidate the phylogenetic relationships within the *Pasiphaea* genus two methods were used to reconstruct the phylogenetic trees: the Maximum likelihood estimation (ML) performed in MEGA v. 5.1Beta3 (Tamura *et al.* 2011) and the Bayesian Inference (BI) method by using MrBayes program (Ronquist *et al.* 2012). The best fitted evolutive method GTR+G+I, adjusted by using the



jModelTest (Posada 2008), was used in both phylogenetic reconstructions with 1000 bootstraps. Eleven independent ML reconstructions were performed to better identify the tree topology and the branch sustentation probability. The BI method was conducted using the Markov chain Monte Carlo (MCMC) algorithm with four Markov chains run for 100000 generations and sampled every 100 generations. The resulting phylogenetic data generated was visualized and edited in the FigTree v.1.4 (<http://tree.bio.ed.ac.uk/software/figtree/>).

Table 2: Species and locality used to reconstruct the phylogeny tree of Pasiphaea genus. UB – University of Barcelona

Species	Sampled area	Haplotype	Geographic coordinates (Lat.Long.)	GenBank Access number
<i>Pasiphaea sivado</i>	Mediterranean – Alacant	PS_hap1	38°08'N -0°04'W	UB (unpublished)
	Mediterranean – Alboran Sea	PS_hap3	36°35'N -4°15'W	UB (unpublished)
	Central E Atlantic – Gulf of Cadiz	PS_hap2	36°15'N -6°43'W	UB (unpublished)
	Atlantic – south Portugal	PS_hap4	36°80'N 7°77'W	JQ306265.1
<i>Pasiphaea multidentata</i>	Mediterranean – Alboran Sea	PM_hap1	36°16'N -4°56'W	UB (unpublished)
	Mediterranean – Catalan Sea	PM_hap3	41°23'N 3°18'W	UB (unpublished)
	Mediterranean – Alboran Sea	PM_hap4	36°16'N -4°56'W	UB (unpublished)
	NE Atlantic – Scotland	PM_hap2	58°29'N 9°W	JG305978.1
	NW Atlantic - Gulf of St. Lawrence	PM_hap2	49°90'N 65°06'W	FJ581854.1
<i>Pasiphaea pacifica</i>	NE Pacific - Gulf of Alaska	PP_hap1	51°54'N 128°21'W	DQ882134.1
		PP_hap2		DQ882132.1
		PP_hap3		DQ882133.1
<i>Pasiphaea tarda</i>	NE Pacific - Gulf of Alaska	PT_hap1	53°42'N 133°24'W	DQ882136.1
		PT_hap2		JQ305937.1
		PT_hap3		DQ882134.1
	NW Atlantic - Gulf of St. Lawrence	PT_hap4	59°21'N 10°W	DQ882139.1
		PT_hap5		JQ305981.1
		PT_hap6		JQ305979.1
<i>Pasiphaea hoplocerca</i>	Central E Atlantic - south Portugal	Unique	36°55'N 9°07'W	JQ306169.1
<i>Acanthephyra pelagica</i>	NE Atlantic – Scotland	Outgroup	58°29'N 9°W	JQ305961
<i>Alvinocaris muricola</i>	not available	Outgroup	not available	EU031814.1
<i>Glyphocrangon neglecta</i>	Caribbean Sea	Outgroup	not available	UB (unpublished)

Pairwise comparison among significant haplotype groups identified by the phylogenetic tree analysis was performed to analyse the degree of difference (as a proportion) among the species/haplotype groups. Within groups p-distances were computed as the proportion of nucleotide sites at which two sequences being compared are different (MEGA v5, Tamura *et al.* 2011).

## 3.3. Results

### 3.3.1 Population genetic structure

The successfully amplified COI fragments presented around 550 and 600 bp for *P. sivado* and *P. multidentata*, respectively. After the joining of the present sequenced data with the extra sequences from GenBank the final COI data set obtained had 518 and 576 bp for *P. sivado* and *P. multidentata* respectively (Table 3).

Table 3: Haplotype frequencies of *Pasiphaea sivado* and *P. multidentata* by sampled area: Catalan Sea, Alacant coast, Alboran Sea (westernmost Mediterranean Sea), Gulf of Cadiz (includes samples taken in Portugal and in Spain).


<i>Pasiphaea sivado</i> (518 bp)	Psiv_01	Psiv_02	Psiv_03	Psiv_04	Total
Catalan Sea - NW Mediterranean	15	1	-	-	16
Alacant coasts - Central W Mediterranean	6	-	-	-	6
Alboran Sea - W Mediterranean	15	-	-	1	16
Gulf of Cadiz - Central E Atlantic	-	10	1	-	11
Total	36	11	1	1	49

<i>Pasiphaea multidentata</i> (576 bp)	Pmul_01	Pmul_02	Pmul_03	Pmul_04	Total
Catalan Sea - NW Mediterranean	2	-	1	-	3
Alboran Sea - W Mediterranean	3	-	-	1	4
Eastern Scotland - NE Atlantic	-	2	-	-	2
Gulf of St. Lawrence - NW Atlantic	-	3	-	-	3
Total	5	5	1	1	12

#### *Pasiphaea sivado*

N = 49



#### *Pasiphaea multidentata*

N = 12




Figure 1: Atlanto-Mediterranean transition showing the sampling locations of *P. sivado* (circles) and *P. multidentata* (squares): Gulf of Cadiz, Alboran Sea, Alacant coast and Catalan Sea.

In *P. sivado*, four different haplotypes were found, of which three sites were polymorphic. The main haplotype was representative of the Mediterranean population (81.82% of total individuals) while the second most frequent haplotype (13.64%) was practically only found in the Atlantic population sampled. The other two haplotypes were unique and derived through a variation in just one polymorphic position from each main haplotype, i. e. one from the Mediterranean haplotype and the other from the Atlantic population (Table 3, Figure 2). Overall, the three populations sampled within the Mediterranean Sea were genetically homogeneous and no gradient was found across the Mediterranean Sea. The gene flow in the western Mediterranean Sea, between Alboran and Catalonia populations, was high and characterized by low values of  $F_{st}$  and no significant values of  $S_{nn}$  (0.484 and 0.763 respectively) (Table 4). However it is important to point out that the nucleotide change observed in the Mediterranean haplotype does translate into an aminoacid change. The nucleotide shift occurred in the second position base and generated a translation from alanine to valine.

Table 4: Population genetic variability of *Pasiphaea sivado* populations across the Atlantic-Mediterranean transition. The statistical significance of  $p < 0.001$  is expressed as (\*\*\*),  $p < 0.01$  (\*\*),  $p < 0.05$  (\*) and non-statistical significance (ns).

<i>Pasiphaea sivado</i>	Within Mediterranean	Within Atlantic	Between Atlantic-Mediterranean
Genetic diversity			
N. of sequences	38	11	49
Sequence size, bp	518	518	518
No. of haplotypes	3	2	4
No. polymorphic sites	2	1	3
Haplotype diversity, $H_d$ ( $\pm$ standard deviation)	0,107 ( $\pm 0.07$ )	0,182 ( $\pm 0.14$ )	0.424 ( $\pm 0.70$ )
Nucleotide diversity, $\Pi$	0.0002	0.0004	0.0009
Number of nucleotide differences, $k$	0.108	0.1818	0.466
Neutral tests			
Tajima's D	-1.493 ns	-1.129 ns	-0.647 ns
Ramos-Onsins and Rozas, $R^2$	0.113	0.288	0.092
Gene flow estimates			
$\Gamma_{st}$			0.74
$S_{nn}^*$			0.956
$\chi^2$			42.854 ***
$F_{st}$			0.867
$K_{st}^*$			0.727

The deeper dwelling *P. multidentata* presented four different haplotypes segregated in six polymorphic sites. The North Atlantic populations, from Scotland (East) and Gulf of St. Lawrence

(West), shared a unique haplotype. The Mediterranean populations concentrated the genetic diversity in three haplotypes, one main and two unique haplotypes derived from two different polymorphic sites (Table 3). However, among the nucleotide changes observed in *P. multidentata* none translated into an aminoacid change. The gene flow among the Mediterranean populations was marked with low values of  $F_{st}$  and no significant values of  $S_{nn}$  (0.0000, 0.4286 and 0.6890 respectively) (Table 5).

Table 5: Population genetic variability of *Pasiphaea multidentata* populations across Atlantic-Mediterranean transition. The statistical significance of  $p < 0.001$  is expressed as (\*\*\*),  $p < 0.01$  (\*\*),  $p < 0.05$  (\*) and non-statistical significance (ns).

<i>Pasiphaea multidentata</i>	Within Mediterranean	Within Atlantic	Between Atlantic-Mediterranean
Genetic diversity			
N. of sequences	7	5	12
Sequence size, bp	572	572	572
No. of haplotypes	3	1	4
No. polymorphic sites	2	0	6
Haplotype diversity, $H_d$ ( $\pm$ standard deviation)	0.524 ( $\pm 0.21$ )	0 ( $\pm 0.00$ )	0.697 ( $\pm 0.09$ )
Nucleotide diversity, $\Pi$	0.001	0	0.004
Number of nucleotide differences, $k$	0.571	0	24.546
Neutral tests			
Tajima's D	-1.237 ns	-	0.898 ns
Ramos-Onsins and Rozas, $R^2$	0.226 ns	-	0.187
Gene flow estimates			
$\Gamma_{st}$			0.873
$S_{nn}^*$			1
$\chi^2$			12.000**
$F_{st}$			0.933
$K_{st}^*$			0.762

The genetic differentiation between Atlantic and Mediterranean populations was significant for both *P. sivado* ( $\Gamma_{ST} = 0.741$   $p < 0.0001$ ) and *P. multidentata* ( $\Gamma_{ST} = 0.873$   $p < 0.0001$ ) showing that the Gibraltar strait represents a barrier to the genetic flux in these benthopelagic shrimps (Tables 4 and 5). Moreover, no haplotype frequency cline was observed across the Atlantic-Mediterranean gradient in either pasiphaeid species. The neutrality tests of Tajima and Ramos-Onsins and Rozas showed that the populations of *P. sivado* and *P. multidentata* did not follow a neutral evolving model, indicating the influence of the natural selection in the genetic differentiation of Atlantic and Mediterranean populations.

### 3.3.2. Phylogeny of *Pasiphaea*

The fragment sequence size obtained for the phylogenetic study had a total of 558 bp. The aligned data presented 175 polymorphic sites, from which 170 were parsimony informative. The pasiphaeid species presented unequal base composition (25.99% A, 19.78% C, 18.74% G and 35.49% T ( $\chi^2= 663.9566$   $p< 0.0001$ ). With a substitution rate of 0.4723% of invariable sites (taking just *Glyphocrangon neglecta* as outgroup) the data presented little saturation according to the substitution saturation analysis of Xia *et al.* (2013) (Iss= 0.482, Iss.c= 0.724,  $p<0.0001$ ). Concerning the nucleotide substitution pattern there was no transitions over transversions bias as indicated by the low ts:tv ratio of 0.403. The proportion of total variable sites in the first and second position were almost equal (43.7 and 43.5 respectively) and greater than those in the third position (12.8) where no transition was detected by Xia's D method.

72

Both the Maximum Likelihood and Bayesian Inference methods were straightforward in placing all the sequences of different haplotypes together in their respective species clade. Moreover the estimated trees properly grouped the haplotypes according to pasiphaeid species and the oceanic region sampled. Both methods also showed the same grouping pattern and similar topologies (Figure 3). Two main groups were identified: a first one segregating all *Pasiphaea sivado* haplotypes from the rest of *Pasiphaea* haplotypes. Within this second group, all haplotypes significantly grouped by species, except in the case of *P. tarda* in which two significant subgroups clustered according locality (NE Atlantic vs NE Pacific).




Figure 3: Phylogenetic tree resolved by Bayesian Inference analysis from cytochrome oxidase I (COI) DNA information on *Pasiphaea* species. A similar tree was reached by using Maximum Likelihood analysis. Small numerals show support levels based on 1000 bootstrap replicates/Bayesian posterior probabilities (ML/BI) expressed as percent.

Pairwise comparison among significant haplotype groups identified by the phylogenetic tree analysis (Table 6) showed that the high divergence of *P. sivado* with the rest of *Pasiphaea* species is comparable with the distance between outgroups and all pasiphaeid species, suggesting that, given probable more future information, the genus *Pasiphaea* could be splitted in two or more genera.

The segregation of *P. tarda* haplotypes in two different clades suggests that species can actually comprise two sibling species, as already shown by (Matzen da Silva *et al.* 2011). *P. tarda*, taking into account both the Atlantic and Pacific populations, was the species that presented the highest intraspecific divergence (4.3%), one order of magnitude greater than the following species, *P. pacifica* (0.6%), *P. multidentata* (0.4%) and *P. sivado* (0.3%). The p-distances between *P. tarda* populations from Pacific (0.4%) and Atlantic (0.2%) oceans were comparable to the above intraspecific values remarking the divergent character of the two populations of *P. tarda*.

Table 6: Pairwise comparison in phylogenetic distance (expressed as a proportion) among significant haplotype groups identified by the phylogenetic tree analysis estimated from COI with GTR+G+I model in (A) Pasiphaea species and outgroups, and (B) Pasiphaea (all species), Pasiphaea excluding *P. sivado*, and outgroups.

(A)									
Between	<i>P. multidentata</i>	<i>P. tarda_1</i>	<i>P. hoplocerca</i>	<i>P. tarda_2</i>	<i>P. pacifica</i>	<i>P. sivado</i>	<i>Alvinocaris</i>	<i>Acanthephyra</i>	<i>Glyphocrangon</i>
<i>P. multidentata</i>	-								
<i>P. tarda_1</i>	0.07	-							
<i>P. hoplocerca</i>	0.18	0.19	-						
<i>P. tarda_2</i>	0.08	0.07	0.18	-					
<i>P. pacifica</i>	0.17	0.18	0.19	0.18	-				
<i>P. sivado</i>	0.2	0.21	0.23	0.2	0.23	-			
<i>Alvinocaris</i>	0.25	0.25	0.24	0.24	0.26	0.25	-		
<i>Acanthephyra</i>	0.21	0.21	0.25	0.21	0.25	0.21	0.2	-	
<i>Glyphocrangon</i>	0.22	0.22	0.24	0.22	0.24	0.21	0.22	0.19	-
(B)									
	<i>Pasiphaea sivado</i>	<i>Alvinocaris</i>	<i>Acanthephyra</i>	<i>Glyphocrangon</i>					
<i>Pasiphaea</i> (all)		0.249	0.217	0.22					
<i>Pasiphaea</i> (excluding <i>P. sivado</i> )	0.215	0.249	0.22	0.225					

## 3.4. Discussion

### 3.4.1. Population genetics

The genetic structure of both *P. sivado* and *P. multidentata* showed a marked break characterized by the almost complete separation of haplotypes between Atlantic and Mediterranean populations, with the absence of a common haplotype. This discontinuity was placed in the Gibraltar Strait area, affecting the populations sampled in the Gulf of Cadiz and Western Alboran Sea. This region has already been reported as a genetic barrier for several species of crustaceans, fishes, cephalopods and other invertebrates (Carreras-Carbonell *et al.* 2005, Schunter *et al.* 2011, Perez-Losada *et al.* 2002, Reuschel and Schubart 2006, Fernández *et al.* 2011). Another evidence of the Gibraltar Strait acting as a genetic barrier to Pasiphaeidae concerns the biogeography of the family, with the presence of only

two species in the Mediterranean Sea, when compared with a much richer presence of pasiphaeid shrimps in nearby Atlantic areas. Thus, d'Udekem d'Acoz (1999) reports a minimum of 14 Pasiphaeidae species in NE Atlantic waters, of which 6 belong to the genus *Pasiphaea*. As a geographically close exemple, *Pasiphaea hoplocerca* has been recorded in south Portugal close to the Gulf of Cadiz but the species is absent in the Mediterranean Sea (Dos Santos and Moreira 2003).

Circulation patterns in the Gibraltar Strait area and throughout the Mediterranean are mainly driven by the water deficit in the Mediterranean basin induced by high evaporation levels due to high atmospheric temperatures in summer and strong, dry, cold northerly winds in winter. Evaporation is especially high in the eastern Mediterranean. Water loss by evaporation also implies a strong salinity increase and associated changes in density. These density differences created by the thermal and salinity changes imply that cooler, lower-salinity water from the Atlantic enter the Mediterranean through the Gibraltar Strait through surface layers (shallower than around 250 m depth) to compensate evaporation and Mediterranean denser water outflux through deeper waters. Atlantic waters entering the Mediterranean basin become modified as they travel east through the Alboran Sea, Almeria-Oran Front and along the North African coasts, from where they travel east and north. They sink in the eastern Mediterranean as salinity increases and then circulate westward through deep layers as Levantine Intermediate waters, also mixing with Mediterranean Deep Water, to finaly spill over the Gibraltar Strait sill. This denser Mediterranean Water outflows into the Atlantic while very slowly remixing with Atlantic water masses (Tizperman and Malanotte-Rizzoli 1991, Cacho *et al.* 1999, Millot and Taupier-Letage 2005). However, despite the relatively good knowledge of Mediterranean hydrography, few works have used crustaceans, or particularly pelagic crustaceans, to identify possible oceanographic barriers for the genetic flux and distribution of species (Pannacciulli *et al.* 1997, Zane *et al.* 2000, Reuschel and Schubart 2006, Luttkhuizen *et al.* 2008, Sala Bozano *et al.* 2009, García-Merchán *et al.* 2012).



Morphological differences between Atlantic and Mediterranean populations of pasiphaeid shrimps were already reported by Zariquiey Álvarez (1957) and Omori (1976). When comparing the description of *Pasiphaea japonica* with the very similar *P. sivado*, Omori (1976) detected the occurrence of significant morphological differences between the newly described *P. japonica* and both Atlantic (Gulf of Cadiz and adjacent waters) and western Mediterranean populations of *P. sivado*. In particular, the total number of spinules on the merus of the first and second pereopods was higher in *P. japonica* vs. both Atlantic and Mediterranean *P. sivado*. Also, significant differences between Atlantic and Mediterranean *P. sivado* were detected, with Mediterranean specimens bearing a higher number of spinules than the Atlantic specimens.

In *P. multidentata*, Zariquiey Alvarez (1957) also described some morphological differences between western Mediterranean and Atlantic (North Sea) populations concerning postrostral spine shape (with Mediterranean specimens showing a longer and more pointed rostrum), and, as in *P. sivado*, concerning also the number of spinules on the merus of the first pereopod, as well as and on the basis and merus of the second pereopod. The Mediterranean form of *P. multidentata* was even described as a different species, under the name *Pasiphaea* (Phye) *sicula* Riggio, 1896, later on considered a subspecies of *P. multidentata* (Zariquiey Álvarez 1946). Sivertsen & Holthuis (1956) did not accept the species status of *P. sicula*. Zariquiey Alvarez (1957) reported that his father R. Zariquiey Cenarro had a paper in press detailing the differences between Mediterranean and Atlantic *P. multidentata* populations that was unfortunately lost during an air raid that destroyed the press during the Spanish civil war (1936-1939). He concluded that Mediterranean specimens could be assigned to *P. multidentata*, but clearly remarked the occurrence of these morphological differences that could in the future assign the Mediterranean specimens to a different "race", as then reported by him, or to significantly different populations using present nomenclature. Casanova and Judkins (1976) reported that all the pelagic decapods that occurred in the Mediterranean were also present in the Atlantic. These authors also detected an interruption in the distribution of several deep sea shrimps with pelagic habits, namely *P. multidentata*, *Gennadas elegans* and *Acanthephyra pelagica*, west of the Gibraltar

Strait, thus indicating that geographical isolation could lead to morphological divergence between Atlantic and Mediterranean populations of these species. The absence of *P. multidentata* in the trawlable bottoms of the Cadiz Gulf (López de la Rosa 1997) is in agreement with the distributional gap also reported by Casanova and Judkins (1976).

The occurrence of these phenotype differences already reported by morphologists and taxonomists, together with the occurrence of geographic differences in population biology (see Chapters 1 and 2) are in agreement with the high genetic differences observed in the present study between Atlantic and Mediterranean populations of both *P. sivado* and *P. multidentata*. These morphological and genetic characteristics may probably be adaptive to the different oceanographic conditions found in the two seas, with salinity being markedly higher in the Mediterranean than in the Atlantic (Millot and Taupier-Letage 2005).

In most decapod crustaceans, and particularly in benthic and epibenthic species, connectivity among populations is assumed to occur during the relatively long planktonic larval stage (Pineda *et al.* 2007, Galarza *et al.* 2009, Schunter *et al.* 2011). However, adult pelagic and benthopelagic shrimps can also be involved in long displacements during the pelagic phase of their activity rhythms (Aguzzi *et al.* 2007, Naylor 2010). To date not much is known on the early life stages of pasiphaeid species. *P. sivado* late stage larvae were captured near the sea bottom around the 100 m isobaths by Williams (1960) in the Irish Sea. Nanjo and Konishi (2009) observed that *P. japonica* larvae reached the decapodid stage in 12 days when kept at surface temperatures, but full development was not accomplished at much lower temperatures, such as those inhabited by the adults, suggesting that larvae would have to develop in shallower waters. Considering that juvenile *P. sivado* occur on the upper slope, in much shallower waters than the adults (see Chapter 1 of this thesis) and that they can reach shallow epipelagic waters during the night (see Chapter 3 of this thesis) we can suppose that dispersion of *P. sivado* along the Iberian coast of the Mediterranean Sea would be mainly guided by the oceanographic processes taking place in epipelagic and intermediate waters.

Nanjo and Konishi (2009) also remarked that the larval development of *P. japonica* is lecithotrophic, and that the larvae could go through the entire larval stages utilizing their yolk. These authors also suggested that larvae of *P. sivado* and *P. tarda* could also be able to grow without feeding due to the large amount of yolk in the egg, as reported by Gurney (1942) and Williamson (1960). Ramirez-Llodra *et al.* (2007) also showed that egg size in *P. multidentata* is indicative of lecithotrophic development with a reduction in the number of larval stages. Company *et al.* (2001) also showed that *P. sivado* eggs are remarkably large. This would imply a smaller residence time in the plankton with the consequent much weaker dispersion ability of the larvae and its retention within intermediate and deep waters. Taking into account that larger larvae would tend to feed on larger prey, they would depend less on direct epipelagic sources, while remaining in deeper waters. This would also imply that the genetic flux of the species is governed by the deep Mediterranean circulation, ie. by the displacement of the Levantine Intermediate Water (LIW) from the eastern Mediterranean towards the Gibraltar Strait. This lack of connectivity between Mediterranean and Atlantic basins due to the existence of the Gibraltar sill at around 300 m is probably the cause of the marked population genetic differences between the two basins in both *P. sivado* and *P. multidentata*.

According to Casanova and Judkins (1976), the deep-sea sergestid benthopelagic shrimp *Sergia robusta*, in contrast with *P. multidentata*, is distributed continuously throughout the Mediterranean and neighbouring Atlantic. Two different morphotypes have been reported in this species, one present in Atlantic waters, and another in Mediterranean (Crosnier and Forest 1973, Casanova and Judkins 1976). The distribution of these two morphotypes, based in petasma morphology, in the vicinity of Gibraltar Strait was studied by Casanova and Judkins (1976), reporting that the Mediterranean form occurred throughout the Mediterranean with the westernmost record located in the Gibraltar Strait area slightly west of Tarifa, not entering the Gulf of Cadiz, while the Atlantic form was sampled in the Atlantic not being located east of Gibraltar. These authors also noted that, although the two morphotypes co-occurred in the Gibraltar Strait, no intermediate morphologies were detected, thus excluding interbreeding between these morphs in this zone of contact. This

pattern suggests that the connectivity of *S. robusta*, as that of *P. sivado* and *P. multidentata*, is governed by larval depth distribution in deep water layers in a way that it is not possible for them to enter the Mediterranean through epipelagic waters incoming from the Atlantic, and it is not also possible to exit the Mediterranean through the Gibraltar sill via intermediate waters. Larval distribution in these species should be therefore deeper than the Gibraltar sill.

Present evidences on marked genetic differences between Atlantic and Mediterranean populations of both *P. sivado* and *P. multidentata*, together with assumptions on relatively deep-sea larval location and behaviour, and present patterns of circulation through the Gibraltar Strait, would be suggestive of a limited genetic flux at present. However, hydrographic patterns during Holocene glaciation cycles have necessarily had to change and allow, with much cooler waters in most of the NE Atlantic including the Gibraltar Strait area and NW Africa, a much higher degree of connectivity between the Atlantic and Mediterranean (Cacho *et al.* 1999, Hewitt 2000, Maggs *et al.* 2008, Palero *et al.* 2008). More knowledge on larval ecology and behaviour is necessary to be able to understand the movements and connectivity of these species. The precise location of the larvae in the water column and their daily vertical movements, if any, are a keystone to understand both connectivity and population dynamics of these species.

### 3.4.2 Phylogeny

*Pasiphaeidae* is a worldwide family, with a relatively large number of species: 97 grouped in seven genera. The most representative genus of the family is the *Pasiphaea*, composed by 66 species, while the resting 31 species of the family is distributed in 6 other genera (De Grave & Fransen, 2011). Present results provide evidences on the internal phylogenetic structuring within the genus *Pasiphaea*, with especial emphasis on the northeastern Atlantic and Mediterranean species, although making use of the scarce available information on other species of the family within GenBank database (null concerning other genera of the family). Therefore, while the total

number of species used in the analysis is relatively low, the results have shown significant evidences on the internal structuring of the genus *Pasiphaea*. Two main groups have been recognized, one comprising all *P. sivado* haplotypes, the other grouping the rest of pasiphaeids analysed. Within this second group, all haplotypes significantly clustered by species, except in the case of *P. tarda*, where, as already shown by Matzen da Silva *et al.* (2011), two significant groups were obtained, one corresponding to NE Atlantic haplotypes and the other to NE Pacific haplotypes.

Taxonomists have already reported on the occurrence of several species groups within the genus, the most important one concerning the *P. sivado* species group. Hayashi (1999, 2006) proposed the differentiation of the *Pasiphaea sivado* group, composed by *P. sivado* (Risso 1816) and ten other species from the rest of the *Pasiphaea* genus, based mainly on branchial reduction characteristics. Within this group, two additional subgroups could be identified according to branchial reduction: those species having four pleurobranchiae (*P. debitusae* Hayashi, 1999, *P. fragilis* Hayashi, 1999, *P. gracilis* Hayashi, 1999, *P. laevis* Hayashi, 1999, *P. marisrubri* Iwasaki, 1989, *P. philippinensis* Hayashi, 1999, and *P. mclaughlinae* Hayashi, 2006), and those with five pleurobranchiae, although the last pleurobranchia is small and rudimentary (*P. sivado* (Risso, 1816), *P. propinqua* De Man, 1916, and *P. japonica* Omori, 1976).

Present results, although lacking information on the rest of species of the *P. sivado* group, provided relevant information concerning the significant phylogenetic differences of the species *P. sivado* versus the rest of other available species of the *Pasiphaea* genus. The highly morphologically similar *Pasiphaea japonica* Omori, 1976 was rather recently described as a new species, but remarking the very high similarities with *P. sivado*, with which it was previously confounded, as clarified by Omori (1976). Unfortunately no information on DNA COI haplotypes is yet available for the species, so it could not be used for the phylogenetic analyses. Although these findings may suggest that *P. sivado* could correspond to a distinct genus within the *Pasiphaeidae*, new phylogenetic analysis adding additional pasiphaeid species from the *P. sivado* group, as well as from as many other *Pasiphaea* species is needed to confirm the internal phylogenetic relationships within the genus and family.

*P. multidentata* has shown its closer affinities with the *P. tarda* complex, both with the Pacific and Atlantic probable species, than with the rest of species used, *P. pacifica* and *P. hoplocerca*. Within the species, the three Mediterranean haplotypes clustered together, while the Atlantic haplotype joined this subgroup, although at a non-significant value, however revealing and confirming the occurrence of differences between the two basis, as also detected by morphological and taxonomic studies (Zariquiey Álvarez, 1957, 1968).


The identification of two significant clades within *P. tarda* was already reported by Matzen da Silva *et al.* (2011), as present results confirm. It must be also taken into account that the taxonomic history of *P. tarda* is complex. Sund (1913) considered that *Pasiphaea princeps* Smith 1884, *P. principalis* Sund 1913, and *P. tarda* Kroyer 1845 were three distinct species, while Sivertsen & Holthuis (1956) concluded that these three species were actually synonyms for one species, *P. tarda*. According to Williams (1960) few authors distinguished between *P. multidentata*, *P. tarda* and *P. principalis* and consequently the name *P. tarda* had been generally used to include any of these species. More recently Iwasaki (1990) defended the species status of *P. princeps* based on relevant morphological characteristic. In this scenario a careful taxonomic revision is needed at morphological level together with clarification from the geneticists point of view.

From the phylogenetic point of view, the precise placement of the family *Pasiphaeidae* within the Caridea, as well as the relationships among the genera and species within the family, are not yet fully understood (Christoffersen 1990, Bracken *et al.* 2009, Chan *et al.* 2010, Matzen da Silva *et al.* 2011, Li *et al.* 2011). In the recent work of Li *et al.* (2011), using nuclear genes, the *Pasiphaeidae* were shown to be monophyletic, and they belonged, together with *Rhynchocinetidae*, *Oplophoridae*, *Nematocarinidae*, *Alvinocarididae*, *Campylonotidae*, and *Eugonatonotidae* to one of the two major clades identified within the Caridea. In two also recent molecular studies (Bracken *et al.* 2009, Chan *et al.* 2010) the *Pasiphaeidae* were shown not to be basal lineages within the Caridea, as were sometimes considered based on some morphological characteristics (Thompson 1967, Chace 1992), but rather they were found to be nested within a large caridean clade.

# Chapter

# 4

Vertical and seasonal distribution of pelagic decapod crustaceans over the shelf-break and middle slope in two contrasting zones around Mallorca (western Mediterranean Sea)



# Vertical distribution of pelagic decapod crustaceans over the shelf-break and middle slope in contrasting seasons and zones around Mallorca (western Mediterranean Sea)

## 4.1. Introduction

Pelagic decapod shrimps have a widespread geographic and bathymetric distribution from high latitudes in both hemispheres to intertropical waters (Casanova and Judkins 1977, Fasham and Foxton 1979, Clarke and Holmes 1987, Burghart *et al.* 2007). Provided of high locomotion abilities, some species are able to maintain themselves at particular water masses as well as to migrate to other water masses or layers (Kaartvedt *et al.* 1988, Karuppasamy *et al.* 2006). This displacement is usually cyclic and associated to the photoperiod as shown in some pasiphaeid and sergestid shrimps (Frogliani and Giannini 1982, Cartes *et al.* 1993, Aguzzi *et al.* 2007). The vertical daily migration ability present in the behaviour of these crustaceans (Naylor 2010) provides them with an important role in the transference of matter and energy from the upper primary productive layers of the ocean, where these species tend to feed at night, down to the middle slope epibenthic community, where many of them dwell during day-time hours (Cartes 1993a, Herring and Roe 1988, Naylor 2010). Some of these pelagic shrimps have a relevant ecological importance as well as an exploitation potential as commercial target species (Nanjo and Ohtomi, 2009). Relevant studies faunistics, biology and ecology of pelagic shrimps have however been mainly conducted only in a few regions, such as the Mediterranean Sea (Casanova and Judkins 1977, Orsi-Relini and Relini 1990, Company *et al.* 2001, 2003, Ramirez-Llodra *et al.* 2007), the Japan Sea (Nanjo 2007, Nanjo and Ohtomi 2009), or the Northeast and Southeast Atlantic Ocean (Matthews and Pinnoi 1973, Gibbons *et al.* 1994, Kensley and Schotte 2006).



From the energetics point of view, in oligotrophic areas, such as most of the Mediterranean Sea (Estrada *et al.* 1985, Estrada 1996), the secondary production of deep-sea species depends, in a practically exclusive way, on the role of the vertical migrators throughout the water column (Klages *et al.* 2001). Pelagic shrimps constitute a fundamental food item to fish, other crustaceans and cephalopods with epibenthic habits on the continental slope and deep sea (Dalpadado *et al.* 1998, Garrison and Link 2000, Fanelli and Cartes 2008). Some of them can be considered as benthopelagic since they spend most of the daytime in close contact with the bottom (Cartes 1993a, 2001, Aguzzi *et al.* 2007), and accordingly play, as stated above, a relevant role in the energetics dynamics of the continental slope and bathyal ecosystems.

The present work aims at studying the pelagic decapod crustacean assemblages inhabiting two hydrographically differentiated areas around the island of Mallorca in the western Mediterranean. Faunistic and size composition, differential distribution in relation to oceanographic water masses, geomorphology (shelf break versus deep continental slope), time of the day and seasonality were particularly the specific targets of this study.

## 4.2. Material and Methods

### 4.2.1. Study area and hydrographic setting

Mallorca is the main island of the Balearic archipelago, in the western Mediterranean. It is located east of the Iberian Peninsula and is separated from the continent by a wide hydrogeographical basin with maximum depths of 2200 m (Massutí *et al.* 2008). This archipelago is located in an area showing high hydrographical variability, both in space and time, which is mainly conditioned by the circulation of water masses through their channels (López-Jurado *et al.* 2008). Surface waters are usually formed by lower salinity Atlantic Water entering from the south through the channels between the islands, and between them and the continent. Below 100 m depth more saline Mediterranean water masses predominate. The Western Mediterranean Intermediate Waters

(WIW) usually occupies the 100-300 m depth stratum, the Levantine Intermediate Waters (LIW) are placed between 200-700 m and finally the Western Mediterranean Deep Water (DW) is found on the bottom (Pinot *et al.* 2002).

The geomorphology of the Balearic Basin, coupled to the climatological variability of the region, confers to this Sea a very dynamic hydrographical scenario, which is conditioned mainly by the amount of WIW occupying the submarine channels. During mild winters the Northern current (NC) from the Gulf of Lions flows southward along the continental slope, reaching the Ibiza Channel and bifurcates creating the north-eastern Balearic current (BC) when reaching surface low salinity Atlantic Water, while part of the main current flows southwest along the continental margin. However during severe winters the high amount of WIW occupying the channels block partially the NC current reinforcing the BC, recirculating the waters in the NW basin without a significant transport of Mediterranean waters through the channels into the Algerian Basin (Monserrat *et al.* 2008).

The hydrographical characteristics actually observed during the sampling cruises in the study area (Olivar *et al.* 2012, Torres *et al.* 2013) showed that surface waters were of recent Atlantic origin. During the December cruise, both temperature and salinity were homogeneous within the first 60-80 m of the water column and a highest gradient was located from there to 120 m depth. In the summer the mixing layer was restricted to the first 10 m and from there a gradient zone was placed until 100 m depth. Below the thermocline layer down to 400 m temperature was approximately constant at around 13 °C, while salinity gradually increased up to 38.5 until reaching 300-400 m depth.

### 4.2.2. Sampling

On board the R/V “Sarmiento de Gamboa” two cruises were conducted around the Mallorca Island: one in December 2009 (late autumn) and the other in July 2010 (early summer) (Olivar *et al.* 2012). Two geographic regions were selected according to previous knowledge on water productivity characteristics: one off Cabrera Island (in the SE of Mallorca, Algerian sub-basin) and the other off the town of Sóller, along the western coasts of Mallorca (Balearic sub-basin) (Figure 1). Within each area

samples were taken over (1) the continental shelf-break and (2) the middle slope. Hydrographical data were obtained by means of CTD casts from the surface to the bottom with a CTD SBE911 in two regular grids of stations covering from 100 to 1200 m isobaths.

Aggregations of pelagic organisms were detected by acoustic backscatter using a Simrad EK60 echosounder and then the sampling was performed with midwater trawls at different depths of the water column inside the acoustic layers that presented the strongest and widest echo-reverberation, i.e. the Deep Scattering Layers (DSL) in mid waters (ca. 400 m) and the Benthic Boundary Layer (BBL), ca. 50 m above the bottom.

Nekton and micronekton were sampled by two different pelagic gears: a Pelagic Trawl net (PT) and an Isaaks-Kidd Midwater Trawl (IKMT). Trawl gear geometry and depth were controlled using Scanmar sensors. The Pelagic Trawl net used was a double-warp, modified commercial mid-water trawl, with standard pelagic otter board doors and a codend mesh size of 10 mm; the effective duration of the trawls was of one hour at a speed of 4 knots (nautical miles  $h^{-1}$ ). PT densities were calculated as number of individuals per hour standardized to 100  $m^2$  of net mouth surface. The IKMT net had a length of 3 m, 3  $m^2$  of mouth opening and 3 mm codend mesh size, the trawls had duration of 30 minutes at a speed of 3 knots. IKMT densities were calculated as number of individuals per 104  $m^3$  filtered. More detailed




Figure 1: Map of Mallorca Island (western Mediterranean) showing the study areas (shelf-break (SH) and medium slope (SL) of Sóller and Cabrera).

sampling information is available in Olivar *et al.* (2012). Table 1 shows the number of samples taken within each Sampling Cell (SC), i.e. combination of season (December/July), area (Cabrera/Sóller), shelf/slope and daylight strata by sampling gear (PT and IKMT). A total of 36 valid PT samples were obtained, 12 in Cabrera and 24 in Sóller. In the December cruise all the PT hauls were performed at night time; while in July the shelf of Cabrera could not be sampled with PT. A total of 65 valid IKMT samples were obtained, 37 in Cabrera and 28 in Sóller. Additionally, each sample was assigned to one of the three defined positions in the water column, i.e. Epipelagic (samples taken in near surface waters well above the DSL), DSL (which was usually located around 400 m depth in both study periods) or BBL (samples taken at around 50 m above the bottom).

Pelagic decapod crustaceans were sorted out and identified to species level. Total fresh weight and number of specimens per species and haul were obtained from PT hauls. All individuals, or a representative subsample of a minimum of 50 individuals, were sexed and measured (carapace length (CL), in mm) with a 0.01 mm precision using a digital calliper. The samples from IKMT were fixed in buffered 5% formalin. After sorting out and species identification and quantification, each specimen was sexed and CL measured using a 'Leica MZ12' magnifier provided with ocular micrometer.

Size frequency distributions (SFD) (carapace length, CL, in mm) were generated by pelagic layers (Epipelagic, Deep Scattering Layers - DSL, and Benthic Boundary Layers – BBL) for the most abundant species for both Pelagic Trawl and IKMT samples. Additionally, samples obtained from the epibenthic/demersal survey taking place in parallel with the pelagic survey (Massutí *et al.* 2013) were also used to generate the corresponding SFD for the population fraction of each species sampled in the epibenthic environment during day time. The gears used to collect demersal fauna were a "Huelvano tiranero" trawl with an 16 m horizontal width, opening height of 2.7-3.2 m and 20 mm codend mesh size, and an epibenthic beam trawl (horizontal and vertical openings: 3.5 m and 0.6 m respectively; codend mesh size: 10 mm).

### 4.2.3. Data analysis

Due to the different dimensions and mesh-size of the sampling gears, which affect the efficiency of sampling of particular species or size classes, data from each gear were analysed separately. The abundance values of pelagic hauls (PT) were shown as number of individuals per hour, standardized to an area of 100 m<sup>2</sup> of sampler mouth opening, while the abundance values from the IKMT were standardized to number of individuals per 10.000 m<sup>3</sup> of filtered water. Prior to applying the statistical tests the data were normalised by a double square root transformation. Since samples taken during the day were very scarce concerning decapod crustacean occurrences, only night samples were analyzed for comparative studies.

The faunistic structure of the decapod community sampled by each net was analysed through multivariate two-way clustering and non-metric multidimensional scaling (MDS) analysis. All species matrices were obtained by calculating Bray-Curtis distances similarity. Significance of the cluster branches was tested using the SIMPROF subroutine within the Primer v.6 software using a significance level of 0.05 in the case of the PT samples and of 0.10 in the IKMT samples. The percentage contribution of each species to the dissimilarity between two groups was quantified using SIMPER analysis. All the statistics were performed by using the software PAST v.2.11 (Hammer *et al.* 2001). To specifically test for the occurrence of significant differences between areas (Cabrera/Sóller) and seasons (December/July) an ANOSIM test was performed within epipelagic and DSL layers. Samples without decapod crustaceans were not used in the analyses. Species richness, Shannon-Weaver diversity ( $H'$ ; using  $\log_2$ ) and Pielou's equitability ( $J$ ) indexes were used to analyze the species diversity of the samples.

## 4.3. Results

### 4.3.1. Faunistics

Overall, a total of 14 species were present in the samples taken (both PT and IKMT) (Tables 1 and 2): five species of dendrobranchiate shrimps, eight caridean shrimps and one scyllarid lobster (nisto postlarval stage). Figure 2 shows the overall depth distribution of the different species sampled, taking into account both PT and IKMT samplings, excluding those taxa not identified to species level and the scyllarid nisto stage. *Chlorotocus crassicornis* and *Plesionika heterocarpus* were only sampled in the shelf-break; *Pasiphaea sivado* and *Sergestes vigilax* were sampled throughout the sampled depths, but especially so in the shelf-break; *Sergestes arcticus* was also found throughout, but especially so in the middle slope region; *Gennadas elegans* reached shallow depths, but was especially present on deep samples; *Pasiphaea multidentata* and *Sergia robusta* were only located in deeper waters, never reaching the epipelagic layers.




Figure 2: Depth distribution of the decapod crustacean species sampled with both the Pelagic Trawl and IKMT nets: depth range, 25 and 75 percentiles and median depth and of the overall species occurrences.

Table 1: Mean density (number of individuals h<sup>-1</sup>) of decapod crustacean species sampled by the pelagic trawl gear (PT). The number of samples taken by season (December/July), geographic area (Cabrera/Sóller), bathymetric zone (Shelf/Slope) and time of the day (Day (D)/Night(N)) are shown on the bottom line.

Pelagic samples	December				July											
	Cabrera		Sóller		Cabrera		Sóller									
	Shelf	Slope	Shelf	Slope	Shelf	Slope	Shelf	Slope								
	D N	D N	D N	D N	D N	D N	D N	D N								
<i>Chlorotocus crassicomis</i> (A. Costa 1971)	—	4.1	—	0	—	1.6	—	0	—	—	0	0	0	18	0	0
<i>Gennadas elegans</i> (S.I. Smith 1882)	—	0	—	3	—	0	—	23	—	—	0	5.1	0	0	0	0.7
<i>Pasiphaea multidentata</i> Esmark 1866	—	0	—	1	—	0	—	4	—	—	0	0.9	0	0	0	15
<i>Pasiphaea sivado</i> (Risso 1816)	—	31	—	2	—	16	—	1.4	—	—	0	2.7	0	37	1	9.4
<i>Plesionika heterocarpus</i> (A. Costa 1971)	—	0.6	—	0	—	1	—	0	—	—	0	0	0	2.2	0	0
<i>Plesionika narval</i> (J.C. Fabricius 1787)	—	0	—	0	—	0	—	0	—	—	0	0	0	0	0	0
<i>Plesionika sp.</i>	—	0	—	0	—	0	—	0.1	—	—	0	0	0	0	0	0
<i>Processa canaliculata</i> Leach 1815	—	0	—	0	—	0	—	0	—	—	0	0	0	0	0	0.9
<i>Scyllarus arctus</i> (Linnaeus 1758)	—	0	—	0	—	0	—	0	—	—	0	0	0	0	0	0
<i>Sergestes arachnipodus</i> (Coco 1832)	—	0	—	2	—	0.1	—	8	—	—	0	0	0	0	0	0.9
<i>Sergestes arcticus</i> Krøyer 1855	—	0	—	8	—	0.3	—	34	—	—	0	30	0	0	2	85
<i>Sergestes vigilax</i> Stimpson 1860	—	0	—	0	—	0	—	0	—	—	0	1.8	0	0	0	4.7
<i>Sergia robusta</i> (S.I. Smith 1882)	—	0	—	0	—	0	—	4.9	—	—	0	1.5	0	0	0	6.9

**Table 2:** Mean density of decapod crustaceans sampled by the IKMT net presented in number of individuals per 104 m<sup>3</sup>. The number of samples taken by season (December/July), geographic area (Cabrera/Sóller), bathymetric zone (Shelf/Slope) and photoperiods (Day (D)/Night(N)) are indicated on the bottom of the correspondent column.

KMT samples	December								July							
	Cabrera				Sóller				Cabrera				Sóller			
	Shelf		Slope		Shelf		Slope		Shelf		Slope		Shelf		Slope	
	D	N	D	N	D	N	D	N	D	N	D	N	D	N	D	N
<i>Acanthephyra pelagica</i> (Risso 1816)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Gennadas elegans</i> (S.I. Smith 1882)	0	0	0	6	0	0	0	4.2	0	0	0	11	0	0	30	4
<i>Pasiphaea multidentata</i> Esmark 1866	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0
<i>Pasiphaea sivado</i> (Risso 1816)	0	0	0	0	0	0	0	0.6	0	0	0	1.2	0	1	2	1
<i>Plesionika heterocarpus</i> (A. Costa 1971)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Sergestes arachnipodus</i> (Cocco 1832)	0	0	0	0	1	5	0	0	0	0	0	0	0	0	0	1
<i>Sergestes arcticus</i> Krøyer 1855	0	0	0	4	0	0	15	10	0	0	0	21	0	0	3	4
<i>Sergestes vigilax</i> Stimpson 1860	0	0	0	5	10	0	1.2	0	0	0	1.8	0	0	0	0	1
<i>Sergia robusta</i> (S.I. Smith 1882)	0	0	0	1	0	0	0	0	0	0	0	0	0	0	2	0
<b>No. Samples</b>	3	4	1	6	1	1	1	2	9	8	4	2	8	6	3	6

### 4.3.2. Day/night depth distribution

The depth and time of the day distribution of the occurrences of the main sampled species (Figure 3) clearly showed that all decapod crustacean species sampled avoided the depths shallower than around 400 m during the central hours of the day. A clear U-shaped pattern, with occurrences reaching epipelagic waters during the night, was found in *P. sivado*, *S. arcticus* and *S. arachnipodus*, species which reach 30, 30, and 50 m respectively. However this pattern was not so evident in *G. elegans*; *P. multidentata* that did not reach epipelagic waters found just below 400 m; as well in *P. heterocarpus* and *C. crassicornis* that were only sampled during night-time hours.






Figure 3: Depth and time of the day distribution of the occurrences of the main sampled species. The first figure shows the overall arrangement of the samples taken, including both Pelagic Trawl and IKMT sampling gears. The images of the fresh sampled individuals are presented.

### 4.3.3. Pelagic Trawl samples

A total of 13 decapod crustacean species were present in Pelagic trawls (PT) samples (Table 1). The species showing the highest densities were *Sergestes arcticus* and *Pasiphaea sivado*, followed by *Gennadas elegans*, *Chlorotocus crassicornis* and *Pasiphaea multidentata*.

The species rank plots show that the dominance pattern of pelagic decapod species varied




Figure 4: Dominance plots for the Pelagic Trawls (PT) and IKMT day and night samples. The values represent the percentage of the total abundance of the ranked species within each Sampling Cell, i.e. sampled combination of season (December, D/July, J), area (Sóller, S/Cabrera, C) and shelf (H)/slope (L) factors. Key symbols indicate the combination of factors: black symbols represent December samples; white, July samples: squares, Cabrera-shelf; triangles, Cabrera-slope; circles, Sóller-shelf; diamonds, Sóller-slope.

between sampling cells (Figure 4). The highest dominance was found in the December shelf-break cells in both Sóller and Cabrera, which were dominated by a single species (*P. sivado*) accounting for >85% of the total abundance; only 3-4 species occurred in these SCs. The lowest dominance was found in the December slope cells, where the number of species found was higher, seven in both areas. The summer samples showed an intermediate pattern.

Cluster and MDS analyses (Figures 5 and 6) showed the occurrence of five significant (according to SIMPROF subroutine) groups of samples, arranged according to a depth gradient, clearly evident in the MDS analyses, in addition to one single isolated epipelagic sample characterized by the occurrence of a single species, *Sergestes vigilax*. Table 3 shows the main characteristics of the groups of samples identified by cluster analysis, such as depth of the samples, distance from the bottom, depth of the bottom, percentage of the samples taken in each season, area, depth stratum and day time period. Clusters A and B, comprised the samples taken over the shelf-break; in particular, all BBL shelf-break samples belonged to cluster A, while the epipelagic samples were located in cluster B. Samples in group A were characterized by two species, *Chlorotocus crassicornis* and *Plesionika heterocarpus*, exclusive of the shelf-break clustering, in addition to *Pasiphaea sivado*, a





Figure 5: Two-way clustering analysis of Pelagic Trawl samples and species. The central matrix shows the density values (fourth root transformed) of species in each sample; white squares show values greater or equal to 1, in light grey squares values greater than or equal to 2 and the dark grey square values higher than 3.. The sample codes show the combination of season, area, strata, position in the water column and sample number within each survey (December/July + Cabrera/Sóller + shelf(H)/slope(L) + Epipelagic/Deep Scattering Layer/Benthic Boundary Layer + sample number. Solid lines in the sample cluster represent significant branches, as identified by SIMPROF analysis.

Figure 6: MDS of samples based on species abundances from the Pelagic Trawl. Samples are grouped according to previous results of the cluster analysis. The sample codes show the combination of season, area, strata, position in the water column and sample number within each survey (December/July + Cabrera/Sóller + shelf(H)/slope(L) + Epipelagic/Deep Scattering Layer/Benthic Boundary Layer + sample number.



species widely co-occurring throughout all samples, but that attained its largest densities in the shelf-break BBL. The epipelagic shelf break cluster B was characterized by low densities of only *C. crassicornis* and *P. sivado*. Cluster C was a heterogeneous group of epipelagic July samples located

Table 3: Characteristics of the Pelagic Trawl (PT) and IKMT sample groups identified by cluster analysis.

	Pelagic Trawl clusters					IKMT clusters	
	A	B	C	D	E	A	B
<b>no. Hauls</b>	3	2	2	6	5	8	19
<b>no. Species</b>	5	2	2	7	7	3	8
<b>Depth range of the samples</b>	180 - 245	35 - 55	33	215 - 635	30 - 430	21-414	38-600
<b>Mean sample depth</b>	208	45	33	429	143	104	289
<b>Distance from bottom range</b>	55 - 140	145 - 224	257 - 667	25 - 500	183 - 859	7-963	3-891
<b>Mean distance from the bottom</b>	85	185	462	364	663	396	484
<b>Bottom depth range</b>	239 - 340	200 - 259	262 - 700	655 - 939	613 - 906	156-988	195-995
<b>Bottom mean depth</b>	293	230	481	793	778	500	774
<b>% Winter</b>	66.7	50	100	66.7	60	0	57.9
<b>% Summer</b>	33.3	50	0	33.3	40	100	42.1
<b>% Cabrera</b>	33.3	50	50	33.3	40	25	47.4
<b>% Sóller</b>	66.7	50	50	66.7	60	75	52.6
<b>% Shelf</b>	100	100	50	0	0	50	21.1
<b>% Slope</b>	0	0	50	100	100	50	78.9
<b>% Epipelagic</b>	0	0	100	0	80	12.5	10.5
<b>% DSL</b>	0	0	0	83.3	20	12.5	57.9
<b>% BBL</b>	100	100	0	16.7	0	75	31.6
<b>% Day</b>	0	0	0	0	0	37.5	26.3
<b>% Night</b>	100	100	100	100	100	62.5	73.7

over both the shelf break and middle slope, with just *P. sivado* occurring in the shelf sample, and *P. sivado* and *S. arcticus* in the slope sample. Samples taken over the deep continental slope clustered in groups D and E. In particular, group D encompassed most of the slope DSL, while group E was formed by most of the epipelagic samples. The species that characterized group D were *Gennadas elegans*, *Sergia robusta* and *Pasiphaea multidentata* which did not occur in any of the epipelagic samples. The most characteristic species of the epipelagic assemblage were different sergestid shrimps, namely *Sergestes vigilax*, *Sergestes arachnipodus*, and *Sergestes arcticus*, together with *P. sivado* which, as stated above, was present throughout all the sampling cells. SIMPER analysis (Table 4) quantifies and further clarifies the role of the different species in the differentiation of the identified assemblages. In the MDS graphic (Figure 6) these groups of samples appeared well defined, except for the overlap of the slope epipelagic group (E) with the arrangement of the slope DSL samples. The different sampling areas (Cabrera/Sóller) did not show any significant difference when applying an ANOSIM test to the data, however hints of seasonality, although proved not significant, were present in both the epipelagic and DSL slope assemblages (Figure 6).

Table 4: Species accounting for the main differentiation between groups of Pelagic Trawls and IKMT samples (percentage contribution to dissimilarity), according to SIMPER analysis. The three most relevant species within each pair of groups are shown in bold.

Species	Pelagic Trawls										IKMT
	A x B	A x C	A x D	A x E	B x C	B x D	B x E	C x D	C x E	D x E	A x B
<i>Acantheephyra pelagica</i>	-	-	-	-	-	-	-	-	-	-	0.9
<i>Chlorotocus crassicornis</i>	<b>17.8</b>	<b>37.5</b>	14.1	<b>20.7</b>	<b>47.2</b>	10.4	<b>17.9</b>	0	0	0	-
<i>Gennadas elegans</i>	0	0	<b>15.2</b>	1.9	0.0	<b>19.9</b>	2.8	<b>22.4</b>	3.6	<b>24.0</b>	<b>18.7</b>
<i>Pasiphaea multidentata</i>	0	0	11.9	3.6	0.0	<b>15.7</b>	4.4	17.6	5.6	<b>19.5</b>	0.8
<i>Pasiphaea sivado</i>	<b>42.0</b>	<b>22.6</b>	<b>13.9</b>	<b>20.3</b>	<b>36.7</b>	4.3	9.1	8.2	<b>16.2</b>	6.9	15.7
<i>Plesionika heterocarpus</i>	<b>28.1</b>	<b>25.9</b>	9.8	14.3	0	0	0	0	0	0	1.6
<i>Processa canaliculata</i>	0	0	0	2.0	0	0	2.4	0	3.0	2.5	-
<i>Sergestes arachnipodus</i>	5.4	4.9	6.8	10.3	0	9.6	<b>18.3</b>	10.8	<b>23.0</b>	10.7	15.0
<i>Sergestes arcticus</i>	6.7	9.1	<b>14.9</b>	<b>22.7</b>	<b>16.1</b>	<b>22.6</b>	<b>39.3</b>	<b>21.2</b>	<b>41.4</b>	9.5	<b>20.1</b>
<i>Sergestes vigilax</i>	0	0	1.3	4.2	0	1.7	5.8	1.9	7.3	5.8	<b>24.1</b>
<i>Sergia robusta</i>	0	0	12.2	0	0	<b>16.0</b>	0	<b>18.0</b>	0	<b>21.0</b>	3.2

The main diversity indexes of the Pelagic Trawl samples within each significant group of samples identified by cluster analysis are shown in Table 5, namely number of species (S), Shannon-Wiener  $\log_2$  diversity index (H) and Equitability (J). Concerning the number of species present, the most diverse assemblages were those sampled above the middle slope bottoms, both in epipelagic (D) and DSL (E) samples, with a total of eight species within each group; the lowest number of species

was found in the epipelagic shelf assemblages, B and C, with just two species in each. The Shannon-Wiener diversity index ranged between 0.09 in assemblage C, which was dominated by *P. sivado*, to 1.24 in the DSL slope (D). Equitability was high in assemblage B (shelf epipelagic) with just two co-occurring species with similar relative densities, as well as in the most diverse slope DSL (group D) in which no high dominance of any particular species was found.

Table 5: Diversity indexes of the Pelagic Trawl and IKMT samples within each significant group of samples identified by cluster analysis: Number of species (S), Shannon diversity (H) and Equitability (J).

		Pelagic Trawls					IKMT	
		A	B	C	D	E	A	B
		SH Epi.	SH BBL	SH	SL DSL	SL Epi.	SH Epi.	SL DSL
No. species	Mín S	3	2	1	5	2	1	1
	Máx S	5	2	2	6	6	2	6
	Median S	3.7	2	1.5	5.7	3.6	1.3	2.6
	Total S	5	2	2	8	8	3	8
Diversity	Mín H	0.43	0.64	0	0.9	0.24	0	0
	Máx H	0.77	0.64	0.2	1.44	1.17	0.69	1.72
	Median H	0.62	0.64	0.1	1.24	0.6	0.17	0.81
Equitability	Mín J	0.39	0.92	0	0.5	0.32	0	0
	Máx J	0.7	0.92	0.3	0.81	0.85	1	1
	Median J	0.5	0.92	0.1	0.72	0.48	0.25	0.77
Number of Samples		7	6	5	11	1	8	19

#### 4.3.4. IKMT samples

Overall, a total of nine decapod crustacean species were present in the IKMT samples (Table 2). The species showing the highest densities were *G. elegans* and *Sergestes arcticus*, followed by *S. vigilax* and *S. arachnipedus*.

The relatively higher number of samples taken with the IKMT gear allowed the analysis of day and night samples separately. The species rank plots show that the dominance pattern of pelagic decapod species varied between sampling cells (Figure 4). Most day samples showed a higher dominance of a single species which accounted for over 75% of the abundance, with respect to the rest of species, while night samples were more evenly distributed.

Cluster and MDS analyses (Figure 7 and 8) showed the occurrence of two significant assemblages, one encompassing most shelf-break samples (A), and the other (B) most slope samples.




Figure 7: Two-way clustering analysis of IKMT samples and species. The central matrix shows the density values (fourth root transformed) of species in each sample; white squares show values greater or equal to 1, in light grey squares values greater than or equal to 2 and the dark grey square values higher than 3. The sample codes show the combination of season, area, strata, position in the water column and sample number within each survey (December/July + Cabrera/Sóller + shelf(H)/slope(L) + Epipelagic/Deep Scattering Layer/Benthic Boundary Layer + sample number. Solid lines in the sample cluster represent significant branches, as identified by SIMPROF analysis.

Assemblage A was characterized by the occurrence of exclusively *P. sivado* in most samples, although *S. vigilax* was also present in some of them, and one single occurrence of *P. heterocarpus*; all samples of this group were taken in July. Within assemblage B, a subcluster, although identified as non-significant by the SIMPROF subroutine, included most of the DSL slope samples, while another subgroup contained most of the epipelagic slope samples. The DSL subcluster was characterized by the co-occurrence of *S. vigilax*, *S. arcticus* and *G. elegans*, while in the epipelagic group *G.*




Figure 8: MDS of samples based on species abundances from the IKMT net. Samples are grouped according to previous results of the cluster analysis. The sample codes show the combination of season, area, strata, position in the water column and sample number within each survey (December/July + Cabrera/Sóller + shelf(H)/slope(L) + Epipelagic/Deep Scattering Layer/Benthic Boundary Layer + sample number.

*elegans* was absent. *P. sivado* showed very few occurrences in assemblage B and *S. robusta* was only present in the DSL subgroup. No clearer differential distribution patterns could be assessed concerning the MDS results.

### 4.3.5. Population size structures

For the most abundant species size frequency distributions were generated by pelagic layers using both Pelagic Trawl and IKMT samples (Figure 9). Additionally, samples from the demersal and epibenthic surveys were also used to generate SFD for the population fraction present in the epibenthic environment during the day. Species-specific patterns were clearly shown for most species. Although the codend mesh size of each sampling gear was different (according to the gear size, structure and organisms size spectra targets), all of them were able to sample both the juvenile and adult fraction of the populations of most species, except perhaps the GOC demersal trawl for the smallest species, such as *Gennadas elegans* and *Sergestes vigilax*.

In the case of the species occurring only over the shelf-break, *Chlorotocus crassicornis* showed a similar size structure in both the epibenthic and BBL layers. In *Plesionika heterocarpus* two cohorts were sampled in the epibenthos, while most of the adult fraction did not appear to show pelagic habits; it is also worth mentioning that *P. heterocarpus* was not found in any epipelagic sample. In the case of *P. sivado*, who was present throughout the pelagic and demersal environments, both adult and juvenile individuals were present in the water column.

Concerning the bathyal benthopelagic species, practically all size classes of *Genadas elegans* were present in the water column, although just a few adults were collected with the larger mesh-size demersal gear, a fact most probably due to the small size of the species coupled to the codend mesh size of the gear. The case of *Pasiphaea multidentata* is interesting because both the juvenile and adult fractions of the population showed pelagic habits; it must also be remarked that the vertical movements of the species did not reach the epipelagic layer, but rather reached the deeper DSL.




Figure 9: Size frequency distribution (in percentage of the total number of individuals sampled within each cell) of the main decapod crustacean pelagic species according to the strata sampled: Epipelagic, DSL, BBL and Epibenthic. Bars represent Pelagic and Demersal Trawl samples, while lines represent IKMT and epibenthic samples. The number of individuals sampled within each cell as well as the mean sampled depth are also shown. The depth of the epipelagic stratum was presented as mean water column depth sampled, the DSL depth was fixed around 400 m, the depth of BBL stratum is presented as distance from the bottom and the epibenthic stratum as bottom depth. PELG: pelagic samples; IKMT: IKMT samples; DEM: demersal samples; EPI: epibenthic samples.






Figure 9: Continuation.




Figure 9: Continuation.

## 4.4. Discussion

### 4.4.1. Study area

The pelagic decapod crustacean fauna in the Mediterranean Sea has not been as thoroughly studied as the epibenthic/demersal one (Casanova 1977, Orsi-Relini and Relini 1990, Koukouras *et al.* 2000). Although specific approaches have focused on some life history aspects and on the ecological role of benthopelagic species (Cartes 2001, Company *et al.* 2001, Papiol *et al.* 2012), information on the distribution, dynamics and role of pelagic shrimps within the Mediterranean pelagic environment is scarce (Koukouras 2000, Froggia and Giannini 1982). There is certainly information on patterns of

occurrence, vertical range, etc, especially so concerning their association with the bottom layers on the continental slope, where most of the species with pelagic habits occur during the day and can be sampled using demersal trawl gears used in bottom fisheries and research cruises (Abelló *et al.* 2002, Fanelli *et al.* 2007, Cartes *et al.* 2009). Present results contribute to assess the role of pelagic and benthopelagic species over the continental edges of the western Mediterranean Sea. Most pelagic decapod shrimps have been reported to be preyed upon by many pelagic and epibenthic predators, including fish, crabs, and shrimps (Abelló 1989, Mauchline and Gordon 1991, Cartes 1993a,b, Valls *et al.* 2011).

The sampling schedule allowed the differential identification of the relevant species occurring on the Balearic shelf edge and slope, their night/day vertical distribution, and provided hints on their seasonality characteristics. The two geographic areas studied, off Sóller and Cabrera in the Balearic Islands, were selected based on previous information on their different oligotrophy degrees and hydrographic dynamics (Massutí *et al.* 2008, Moranta *et al.* 2008, Balbín *et al.* 2012). Overall they did not show significant differences concerning their decapod crustacean faunistic composition. Neither was seasonality in faunistic composition and structure found to vary significantly. In the present study most species showed species-specific patterns of distribution occurrence, especially so concerning the two geomorphologic areas sampled (over the shelf break and over the middle slope) as well as concerning depth in the water column. On the shelf break, species diversity was low, with just two to five decapod crustacean species occurring in the samples; a high dominance of just one or two species was evident in that area. The slope community was more diverse, with up to six species per sample and a total of eight species in the area, followed by a higher overall equitability. In the colder December samples the variation of the dominance pattern between shelf break and slope was more evident than in summer. The species composition, diversity and dominance between the geographic regions, Cabrera and Sóller, was similar in both seasons, suggesting that the dynamics of the pelagic environment (López-Jurado *et al.* 2008, Torres *et al.* 2011, Balbín *et al.* 2012) would tend to uniformize the ecological dynamics of the system.

Differences in pelagic species communities have been reported at different oceanographic scales in many oceanic regions, such as in the Gulf of Mexico, off southern California, western North Pacific,

central-eastern Atlantic, Norwegian Sea, etc, and have been mainly attributed to large scale oceanographic patterns and differences between coastal/neritic and oceanic waters, especially so concerning island environments (e.g. Burghart *et al.* 2007, Gibbons *et al.* 1994, Dalpadado *et al.* 1998, Suntsov and Domokos, 2013). Pelagic decapod crustaceans include individuals belonging to several families, but most importantly to Sergestidae, Penaeidae and Aristeidae, among the Penaeoidea, and Pasiphaeidae, Oplophoridae and Pandalidae among the Caridea (Hopkins *et al.* 1994, Karuppasamy *et al.* 2006). Seasonal differences in the pelagic community composition have also been reported and attributed to seasonal differences in hydrological features, such as occurrence and disruption of thermoclines, upwelling strength, etc. (e.g. Franqueville 1971, Williams and Koslow 1997).

In the Balearic Sea no pelagic decapod species were detected in epipelagic waters shallower than 400 m depth during the day light hours, in a similar way as that reported by Fasham and Foxton (1979) in the Atlantic Ocean. Differences in depth distribution limits between seas can be attributed to the oligotrophic characteristics of the study area in the Mediterranean Sea which allows light spectra to penetrate deeper than in more productive areas (Aguzzi *et al.* 2004, 2007). The avoidance of displacement to illuminated waters by small pelagic species is suggested as a way to avoid the pressure by visual predators (De Robertis 2002, Kronfeld-Schor and Dayan 2003).

At the shelf break, species present in the pelagic samples occur also on the bottom during the day (Abelló *et al.* 1988, Moranta *et al.* 2008). Concerning the species sampled over the middle slope, most of them were also sampled during the day in the DSL and BBL layers, and all of them can also be sampled in close contact with the bottom with both demersal gears (Abelló *et al.* 2002, Moranta *et al.* 2008) and epibenthic sledges (Ramón *et al.* 2013), becoming in this way important actors in their role as prey of epibenthic and demersal predators (Orsi-Relini and Relini 1990, Cartes *et al.* 2009).

Present results also showed the occurrence, in all pelagic decapods, of a rapid vertical displacement during the crepuscular hours (Franqueville 1971). This remarkable cyclic migration pattern has been well studied in many works and interpreted as of high adaptive value to reduce predation in this period (Reid *et al.* 1991, Naylor 2010, Olivar *et al.* 2012).

### 4.4.2. Species and faunistic assemblages

The distribution of the pandalid shrimps *Chlorotocus crassicornis* and *Plesionika heterocarpus* in the present samples was restricted to the continental shelf-break. These species, which typically inhabit the deep continental shelf and the upper slope (Company and Sardá, 1997, Carbonell and Abelló, 1998) have been usually considered to be epibenthic/demersal (Colloca *et al.* 2004, Muñoz *et al.* 2012), but this study has clearly shown their ability to perform pelagic daily movements into the water column. The presence of both species in the diet of *Merluccius merluccius*, by Cartes *et al.* (2004) can corroborate the pelagic character of these species. The presence of *Chlorotocus crassicornis* in the stomach content of fishes sampled in depths above the upper limit of the bottom distribution of this prey together with their absence in the diet of fishes captured on their preferential depth suggests that *M. merluccius* is preying specially on the pelagic fraction of *C. crassicornis*. Other pandalid species widely occurring in the western Mediterranean Sea, particularly in the study area, such as *Plesionika antigai*, *P. edwardsii*, *P. narval*, *P. martia*, or *P. acanthonotus* (Abelló *et al.* 1988, 2002, Moranta *et al.* 2008), were not found to show incursions into the pelagic environment. *Plesionika narval* and an unidentified juvenile *Plesionika* specimen were only identified in one pelagic trawl sample. The processid shrimp *Processa canaliculata* was also only found once.

104

In the present study the pasiphaeid shrimp *Pasiphaea sivado* was found to be the commonest species in the study area, occurring both over the shelf break in neritic waters, and over the middle slope, as well as through all sampled depth strata, from epipelagic waters through DSL and BBL to the epibenthic layers. During the day time the species is known to occur near the bottom, preferentially between 350-500 m depth where it can be sampled by bottom gears along the continental slope (Abelló *et al.* 2002, Aguzzi *et al.* 2007, Simão *et al.* submitted). In the Mediterranean Sea, *P. sivado* shows a preferential occurrence depth shallower than its congeneric species *Pasiphaea multidentata* (Simão *et al.* submitted). On the contrary, the distribution of *P. multidentata* was restricted to the middle slope, and never reached epipelagic waters.

The family Sergestidae was that with the highest taxonomic richness in the samples, with up to four different species occurring commonly in the samples. The most abundant species was

*Sergestes arcticus* and was clearly a dominant species in many samples. Its distribution was however confined to the middle slope, although it reached epipelagic waters during the night (Franqueville 1971, Frogliá and Giannini 1982). The patterns of occurrence and abundance of *S. arcticus* establish it as a key species that characterizes the epipelagic samples for both shelf-break and middle slope areas. The characterization of *S. arcticus* as the dominant species in the epipelagic community is well documented in the Mediterranean Sea and Northeast Atlantic (Frogliá and Giannini 1982, Vestheim and Kaartvedt 2009), often being the second most abundant crustacean after the euphausiid *Meganyctiphanes norvegica* (Franqueville 1971). Larvae of *S. arcticus* are commonly and abundantly found in the study area in both summer and late autumn, being included among the three more abundant decapod crustacean larvae (Torres *et al.* 2013). *S. arcticus* co-occurred with the larger *Sergestes arachnipedus* in many samples, but the former was dominant in abundance in all sampling cells. *S. arachnipedus* was also restricted to the middle slope and had a wide distribution along the water column from epipelagic waters to the epibenthic environment. *Sergestes vigilax* has a smaller mean size than the before mentioned species and occurred more often in IKMT trawls, probably due the smaller mesh size of this gear. In these samples the species was more common than *S. arachnipedus* but never as abundant as *S. arcticus*. The preferential distribution of *S. vigilax* was similar to that of *S. arcticus* but showed a much shallower median depth of occurrence, located at around 200 m. This shallower mean distribution depth, compared with the 400 m depth for *S. arcticus*, could have adaptive value to minimise competition by other sergestids. Also, food targets and feeding behaviour are probably different given the large differences occurring in feeding structures, such as the third maxillipeds, much larger in *S. vigilax* than in *S. arcticus* or *S. arachnipedus*.

*Sergia robusta* is the largest sergestid shrimp found in the area, and its distribution was clearly restricted to the middle slope, never reaching epipelagic waters. Its number of occurrences was not very high but clearly characterized the middle slope environment (Frogliá and Giannini 1982).

The small aristeid shrimp *Gennadas elegans* was practically only sampled in the DSL, BBL and epibenthic layers of the middle slope, not on the shelf break, and rarely in epipelagic waters (Andersen *et al.* 1998).

### 4.4.3. Size structure

The size structure of pelagic decapods in the water column is poorly studied in the literature (Omori and Gluck 1979, Orsi Relini and Relini 1990, Vestheim and Kaartvedt 2009, Suntsov and Domokos 2013). The Sergestidae, except *Sergestes vigilax*, in which no large sized individuals were captured with the epibenthic samplers, showed a quite homogeneous size structure in the water column, with both juveniles and adults occurring in the pelagic environment, in a similar pattern as that shown by Vestheim and Kaartvedt (2009). Size structure was however slightly displaced towards the largest size classes in the epibenthic environment, although small sized individuals were also present in the epibenthic samples.

*Pasiphaea sivado* individuals of all size classes were present in the epibenthic environment during the daytime, but in the water column the population showed a size structure of increasing predominance of large individuals towards the deeper strata, with the smallest sized individuals occurring predominantly in the epipelagic environment at night.

The huge size range of *Pasiphaea multidentata* sampled in the water column showed that all size classes do perform vertical movements in the water column, although the predominance of small size classes in the DSL can suggest a higher independence of the seabed in juvenile individuals. In the present study adult *P. multidentata* in both the DSL and BBL revealed that adults are able to usually perform vertical migrations. The occurrence of pelagic habits in the adult population had only been clearly documented once in this species (Koukouras *et al.* 2000). The adults of the species had been previously thought to perform daily bathymetric movements along the middle slope (Cartes *et al.* 1993, Aguzzi *et al.* 2007). Present results clearly showed that adults of this species may also perform vertical migrations into the water column, however restricted to non-epipelagic layers (up to around 300-400 m depth), in agreement with the low ratio between size and wet weight characteristic of pelagic species (Cartes *et al.* 1993).

#### 4.4.4. Colouration

Pelagic species body colour has been related to depth distribution (Herring and Roe 1988). Transparency is common in pelagic species that dwell in both the euphotic and aphotic zones primarily because it is the only camouflage that is successful from all points of view and at all depths (Johnsen 2001). At the aphotic environment (i.e. below 1000 m) transparent species are rare, being generally replaced by species red or black pigmented. At these depths the visual predation is based on bioluminescent light. The bioluminescent wavelength is almost totally absorbed by red or black pigmentation conferring a more effective crypsis in species with all red body pigmentation than transparency, due the characteristic surface reflections of transparent objects. Despite the red is the first spectral colour that becomes extinct in the first few meters of the water column, the red pigmentation confers opacity to the body at low light intensity waters in a way that the animal will present the same colour of the background (Herring and Roe 1988, Aguzzi *et al.* 2007).

In the study area, the DSL environment was mainly characterized by species showing intense red pigmentation such as *Gennadas elegans*, *Sergia robusta* and *Pasiphaea multidentata*. The depth distribution of these species did not, or rarely did, reach the epipelagic environment. The transparent shrimps of the Sergestidae family presented a huge vertical distribution encompassing all the depths sampled. *Sergestes arcticus* concentrates intense red pigments due the presence of numerous large chromatophores in a restricted area of the cephalothorax, the gut and the extremities of maxillary appendixes, the rest of the body being transparent/translucent. This coloration pattern probably allows *S. arcticus* to be visually identified by other organisms as a much smaller sized mesozooplankton species, which could be of adaptive value. The same mimicry can also be suggested for *Sergestes arachnipedus*, which additionally presents a dorsal dispersion of small red spots and a larger size than *S. arcticus*. Indubitably, to assume these speculations more studies about translucence and light reflection of these species as well as related to the visual sensibility of small pelagic species are needed (Herring and Roe 1988, Johnson *et al.* 2000).

The ontogeny of *P. multidentata* is a remarkable evidence of the influence of preferential depth distribution in body pigmentation pattern of pelagic decapods, since juveniles, which have a



much shallower habitat than adults, present an all transparent body (which often leads to identification confusions with *P. sivado*), while along growth the species accumulates red pigments, becoming redder with age, with larger individuals inhabiting deeper waters (Herring and Roe 1988, Company *et al.* 2001). Differential distribution between juvenile and adult components of a population may be of adaptive value to decrease competition between different age groups.

# Conclusions



## Conclusions

- In both *Pasiphaea sivado* and *Pasiphaea multidentata* the bathymetric distribution, density, maturity size, and population size structure found in the Alboran Sea were clearly different from those in the Balearic basin. The populations in the intermediate Alacant sector showed more affinities with the Alboran Sea populations, especially so in *P. multidentata*, than with the northern populations, in agreement with their common location in the Algerian basin.
- Densities of *P. sivado* showed a heterogeneous distribution with two main nuclei, one in the Alboran Sea (especially so in the western sector) and another along the North Catalan coast. Additionally the bathymetric distribution in the western Alboran sector reached shallower depths than in the rest of sectors, in agreement with the occurrence of coastal upwellings along the northwesternmost region of the Alboran Sea, around Malaga.
- The relationship between occurrence and density patterns by depth allowed delimitating an optimal depth range of 300-500 m in *P. sivado*, and of 500-800 m (the deepest sampled depth) in *P. multidentata*.
- The size structure analysis showed that these species have a marked size increase with depth. *P. sivado* juveniles were present in the upper slope strata of all sectors. However, *P. multidentata* juveniles were restricted to the Balearic basin, implying that seasonality of reproduction significantly differs between basins.
- Juveniles of both species showed a wider occurrence thermohaline window, suggesting their ability to cope with a wider variability in temperature and salinity conditions.
- The occurrence of sexual dimorphism in the shape of the pleopod protopod of *P. sivado* was evidenced by both the traditional method of biometric ratios and by applying geometric morphometrics methodologies: Overall, in all five pleopod pairs, adult female protopods were clearly much thinner than those in adult males. Additionally, the anterior edge of the male protopods was clearly much more convex than in females, while the posterior edge was straight in both males and females.
- The studied populations of Mediterranean Sea *P. sivado* showed the occurrence of sexual dimorphism in size, with females being slightly smaller than males in all geographic areas and depth strata, suggesting that this pattern is intrinsic to the population biology of the species.

- In all geographic sectors females were dominant at sizes smaller than around 18-19 mm CL; after that size, males were dominant. The precise size at which males became dominant decreased progressively from the Alboran Sea (21 mm CL) to the Catalan Sea (19 mm CL) suggesting the occurrence of environmental and ecological processes affecting growth rates.
- The genetic chapter of this thesis is a first attempt to investigate the influence of the Gibraltar Strait on the population genetic structure of pasiphaeid shrimps, namely *P. sivado* and *P. multidentata*. Additionally, the chapter also presents a first approximation to the phylogenetic tree of the *Pasiphaeidae* using the sequences obtained during this study as well as all the COI sequences of pasiphaeid species available in the Genbank database.
- The Gibraltar Strait was clearly identified as a barrier to the genetic flux in both *P. sivado* and *P. multidentata* due the marked genetic structure characterized by the presence of two different main haplotypes in each species, one Atlantic and another Mediterranean. The absence of a genotype gradient and also the absence of a shared haplotype between samples from the different seas indicated that the Gibraltar Strait is a strong and unique geographical barrier to the genetic flux of both pasiphaeid species at the geographical scale analysed.
  - The phylogenetic reconstruction obtained showed that *P. sivado* formed a well-differentiated group from the rest of *Pasiphaea* species studied, in agreement with morphological studies performed within the family. Within the second group, all haplotypes significantly grouped by species, except in the case of *P. tarda* in which, as already shown by other researchers, two well differentiated subgroups clustered according to their Atlantic or Pacific origin. *P. multidentata* has shown its close affinities with *P. tarda* complex.
- The study of the pelagic decapod crustacean fauna of two different zones around Mallorca (Sóller and Cabrera), characterized by different hydrographic dynamics and oligotrophy levels, evidenced the occurrence of several distinct faunistic assemblages related to the bathymetry of the area sampled (shelf-break and middle slope), as well as to the sampling depth (near-surface epipelagic waters and Deep Scattering Layers).
- The two studied zones showed a similar faunistic diversity, with the DSL showing the highest species diversity. No significant differences concerning sampling area or seasonality were found. Some species were restricted to the continental shelf-break:

## Conclusions

*Chlorotocus crassicornis* and *Plesionika heterocarpus*. Others were exclusive of deeper waters over the middle slope region, such as *Gennadas elegans*, *Pasiphaea multidentata*, and *Sergia robusta*. *Pasiphaea sivado* was common in all pelagic strata. The virtually absence of differences in the species composition, diversity and dominance between the geographic regions (Cabrera and Sóller) and seasons (early winter and summer), suggested that the dynamics of the pelagic environment would tend to uniformize the ecological dynamics of the system.

- Size analysis showed the occurrence of species-specific patterns concerning the size/age movements into the water column throughout the day-night cycle.
- The present study clearly confirms the presence of adult *P. multidentata* in the water column, evidencing that the adult fraction of the population has also the ability to perform vertical migrations.
- The importance of several species, such as *Sergestes arcticus*, *Pasiphaea sivado*, *Gennadas elegans* or *Sergia robusta*, due to their abundance and vertical migration patterns, must be taken into account when trying to explain transfers of matter and energy in the pelagic environment coupled to the epibenthic ecology.
- Overall, the size structure of both pasiphaeids and sergestids was displaced towards the largest size classes in the epibenthic environment. This pattern was more marked in *P. sivado*.

# Bibliography

- Abad E, Preciado I, Serrano A, Baro J (2007) Demersal and epibenthic assemblages of trawlable grounds in the northern Alboran Sea (western Mediterranean). *Sci Mar* 71:513-524.
- Abelló P (1989) Feeding habits of *Macropipus tuberculatus* (Brachyura, Portunidae) off the Catalan coast (NW Mediterranean). *Misc Zool* 13:45-50.
- Abelló P, Aagaard A, Warman CG, Depledge MH (1997) Spatial variability in the population structure of the shore crab *Carcinus maenas* (Crustacea: Brachyura) in a shallow-water, weakly tidal fjord. *Mar Ecol-Prog Ser* 147:97-103
- Abelló P, Carbonell A, Torres P (2002) Biogeography of epibenthic crustaceans on the shelf and upper slope off the Iberian Peninsula Mediterranean coasts: implications for the establishment of natural management areas. *Sci Mar* 66S2:183-198
- Abelló P, Cartes JE (1992) Population characteristics of the deep-sea lobsters *Polycheles typhlops* and *Stereomastis sculpta* (Decapoda: Polychelidae) in a bathyal mud community of the Mediterranean Sea. *Mar Biol* 114:109-117
- Abelló P, Macpherson E (1992) Epibiosis and rhizocephalan infestation patterns in relation to the reproductive biology of *Lithodes ferox* (Filhol, 1885) (Anomura: Lithodidae). *J Crust Biol* 12:561-570
- Abelló P, Pertierra JP, Reid DG (1990) Sexual size dimorphism, relative growth and handedness in *Liocarcinus depurator* and *Macropipus tuberculatus* (Brachyura: Portunidae). *Sci Mar* 54:195-202
- Abelló P, Valladares FJ, Castellón A (1988) Analysis of the structure of decapod crustacean assemblages off the Catalan coast (North-West Mediterranean). *Mar Biol* 89:39-49
- Aguzzi J, Company JB, Abelló P (2004) Locomotor activity rhythms of continental slope *Nephrops norvegicus* (Decapoda: Nephropidae). *J Crust Biol* 24:282-290

- Aguzzi J, Company JB, Abelló P, García JA (2007) Ontogenetic changes in vertical migratory rhythms of benthopelagic shrimps *Pasiphaea multidentata* and *P. sivado*. *Mar Ecol Prog Ser* 335:167-174
- Aguzzi J, Costa C, Antonucci F, Company JB, Menesatti P, Sardá F (2009) Influence of diel behaviour in the morphology of decapod natantia. *Biol J Linn Soc* 96:517-532
- Andersen V, François F, Sardou J, Picheral M, Scotto M, Nival P (1998) Vertical distributions of macroplankton and micronekton in the Ligurian and Tyrrhenian Seas (northwestern Mediterranean). *Oceanol Acta* 21:655-676
- Apollonio S (1969) Breeding and fecundity of glass shrimp, *Pasiphaea multidentata* (Decapoda, Caridea) in Gulf of Maine. *J. Fish. Res. Bd. Canada*, 26, 1969-1983.
- Balbín R, Flexas MM, López-Jurado JL, Peña M, Amores A, Alemany F (2012) Vertical velocities and biological consequences at a front detected at the Balearic Sea. *Cont Shelf Res* 47:28-41
- Battaglia P, Andaloro F, Consoli P, Esposito V, Malara D, Musolino S, Peda C, Romeo T (2013) Feeding habits of the Atlantic bluefin tuna, *Thunnus thynnus* (L. 1758), in the central Mediterranean Sea (Strait of Messina). *Helgol Mar Res* 67:97–107
- Ben Mariem S (2004) Première approche de la croissance de *Penaeus kerathurus* (Decapoda Penaeidae) dans le golfe de Gabes, Tunisie. *Crustaceana* 68: 583-596
- Bertrand J, Gil de Sola L, Papaconstantinou C, Relini G, Souplet A (2002) The general specifications of the MEDITS surveys. *Sci Mar* 66S2:9-17
- Bertrand JA, Gil de Sola L, Papaconstantinou C, Relini G, Souplet A (2000) An international bottom trawl survey in the Mediterranean: the MEDITS programme. *Actes de Colloques IFREMER* 26:76-93



- Biagi F, Sartor P, Ardizzone GD, Belcari P, Belluscio A, Serena F (2002) Analysis of demersal assemblages off the Tuscany and Latium coasts (north-western Mediterranean). *Sci Mar* 66S2:233-242
- Bracken HD, De Grave S, Felder DL (2009) Phylogeny of the infraorder Caridea based on mitochondrial and nuclear genes (Crustacea: Decapoda). In: Martin JW, Crandall KA, Felder DL (Eds.). *Decapod Crustacean Phylogenetics*. Crustacean Issues 18. CRC Press, Boca Raton, FL, USA, 274–300
- Burghart SE, Hopkins TL, Torres J (2007) The bathypelagic Decapoda, Lophogastrida, and Mysida of the eastern Gulf of Mexico. *Mar Biol* 152:315–327
- Cacho I, Grimalt JO, Pelejero C, Canals M, Sierro FJ, Flores JA, Shackleton NJ (1999) Dansgaard-Oeschger and Heinrich event imprints in Alboran Sea temperatures. *Paleoceanography*, 14, 698–705.
- Carbonell A, Abelló P (1998) Distribution characteristics of pandalid shrimps (Decapoda: Caridea: Pandalidae) along the Western Mediterranean Sea. *J Nat Hist* 32: 1463–1474
- Carbonell A, Palmer M, Abelló P, Torres P, Alemany R, and Gil de Sola L (2003) Mesoscale geographical patterns in the distribution of pandalid shrimps *Plesionika* spp. in the Western Mediterranean. *Mar Ecol Prog Ser* Vol. 247: 151–158
- Carr SD, Tankersley RA, Hench JL, Forward RB, Luettich RA (2004) Movement patterns and trajectories of ovigerous blue crabs *Callinectes sapidus* during the spawning migration. *Estuar Coast Shelf S* 60: 567-579.
- Carreras C, Pascual M, Cardona L, Marco A, Bellido JJ, Castillo JJ, Tomas J, Raga JA, Sanfelix M, Fernandez, G, Aguilar A (2011) Living together but remaining apart: Atlantic and Mediterranean Loggerhead Sea Turtles (*Caretta caretta*) in shared feeding grounds. *Journal of Heredity*, 102, 666-677.

- Carreras-Carbonell J, Macpherson E, Pascual M (2005) Rapid radiation and cryptic speciation in Mediterranean triplefin blennies (Pisces : Tripterygiidae) combining multiple genes. *Molecular Phylogenetics and Evolution*, 37, 751-761.
- Cartes JE (1993a) Deep sea decapoda fauna of the Western Mediterranean: Bathymetric distribution and biogeographic aspects. *Crustaceana* 65(1):29-40
- Cartes JE (1993b) Diets of deep-water pandalid shrimps on the western Mediterranean slope. *Mar Ecol Prog Ser* 96: 49-61
- Cartes JE (1993c) Day-night feeding by decapod crustaceans in a deep-water bottom community in the western Mediterranean. *J Mar Biol Assoc UK* 73:795-811
- Cartes JE (1993d) Feeding habits of pasiphaeid shrimps close to the bottom on the western Mediterranean slope. *Mar Biol* 117:459-468
- Cartes JE (1994) Influence of depth and season on the diet of the deep-water aristeid *Aristeus antennatus* along the continental slope (400 to 2300 m) in the Catalan Sea (western Mediterranean) *Mar Biol* 120:639-648
- Cartes JE (1998) Dynamics of the bathyal Benthic Boundary Layer in the northwestern Mediterranean: depth and temporal variations in macrofaunal-megafaunal communities and their possible connections within deep-sea trophic webs. *Prog Oceanog* 41:111-139.
- Cartes JE (2001) Trophic structure of a bathyal benthopelagic boundary layer community south of the Balearic Islands (southwestern Mediterranean). *Mar Ecol Prog Ser* 215:23-35
- Cartes JE, Abelló P, Lloris D, Carbonell A, Torres P, Maynou F, Gil de Sola L (2002a) Feeding guilds of western Mediterranean demersal fish and crustaceans: an analysis based on a spring survey. *Scia Mar* 66S2:209-220

- Cartes JE, Abelló P, Lloris D, Carbonell A, Torres P, Maynou F, Gil de Sola L (2002) Feeding guilds of western Mediterranean demersal fish and crustaceans: an analysis based on a spring survey. *Sci Mar* 66(S2):209-220
- Cartes JE, Brey T, Sorbe JC, Maynou F (2002) Comparing production–biomass ratios of benthos and suprabenthos in macrofaunal marine crustaceans. *Can J Fish Aquat Sci* 59:1616–1625
- Cartes JE, Company JB, Maynou F (1994) Deep-water decapod crustacean communities in the Northwestern Mediterranean: influence of submarine canyons and season. *Mar Biol* 120:221-229
- Cartes JE, Huguet C, Parra S, Sánchez F (2007a) Trophic relationships in deep-water decapods of Le Danois bank (Cantabrian Sea, NE Atlantic): Trends related with depth and seasonal changes in food quality and availability. *Deep Sea Res I* (54):1091-1110
- Cartes JE, Maynou F (1998) Food consumption by bathyal decapod crustacean assemblages in the western Mediterranean: predatory impact of megafauna and the food consumption–food supply balance in a deep-water food web. *Mar Ecol Prog Ser* 171:233-246
- Cartes JE, Maynou F, Abelló P, Emelyanov M, Gil de sola L, Solé M (2011) Long-term changes in the abundance and deepening of the deep-sea shrimp *Aristaeomorpha foliacea* in the Balearic Basin: relationships with hydrographic changes at the Levantine Intermediate Water. *J Marine Syst* 88:516-525
- Cartes JE, Maynou F, Fanelli E, Papiol V, Lloris D (2009) Long-term changes in the composition and diversity of deep-slope megabenthos and trophic webs off Catalonia (western Mediterranean): Are trends related to climatic oscillations? *Prog Oceanog* 82: 32-46
- Cartes JE, Papiol V, Guijarro B (2008) The feeding and diet of the deep-sea shrimp *Aristeus antennatus* off the Balearic Islands (Western Mediterranean): Influence of environmental factors and relationship with the biological cycle. *Prog Oceanog* 79:37-54

- Cartes JE, Rey J, Lloris D, Gil de Sola L (2004) Influence of environmental variables on the feeding and diet of European hake (*Merluccius merluccius*) on the Mediterranean Iberian coasts. *J Mar Biol Ass UK* 84:831-835
- Cartes JE, Sardá F, Abelló P (1993) Decapod crustaceans collected by deep-water trawls (between 1000 and 2200 m) in the Catalan area (North-western Mediterranean). *Bios, Thessaloniki* 1: 207-213
- Cartes JE, Serrano A, Velasco F, Parra S, Sánchez F (2007b) Community structure and dynamics of deep-water decapod assemblages from Le Danois Bank (Cantabrian Sea, NE Atlantic): Influence of environmental variables and food availability. *Prog Oceanog* 75:797-816
- Cartes, J. E. & Maynou, F. 1998. Food consumption by bathyal decapod crustacean assemblages in the western Mediterranean: predatory impact of megafauna and the food consumption food supply balance in a deep-water food web. *Marine Ecology - Progress Series*, 171, 233-246.
- Cartes, J. E., Maynou, F., Abelló, P., Emelyanov, M., Gil de sola, L. & Solé, M. 2011. Long-term changes in the abundance and deepening of the deep-sea shrimp *Aristaeomorpha foliacea* in the Balearic Basin: relationships with hydrographic changes at the Levantine Intermediate Water. *Journal of Marine Systems*, 88, 516-525.
- Cartes, J. E., Maynou, F., Morales-Nin, B., Massutí, E. & Moranta, J. 2001. Trophic structure of a bathyal benthopelagic boundary layer community south of the Balearic Islands (southwestern Mediterranean). *Marine Ecology - Progress Series*, 215, 23-35.
- Casanova JP (1976) Crustacés décapodes pélagiques de la province Atlanto-Méditerranéenne - Notes taxonomiques et biogéographiques. *Rapp Comm int Mer Médit* 23(9):63-64
- Casanova JP (1977) La faune pélagique profonde (zooplancton et micronecton) de la province atlanto-méditerranéenne. Aspect taxonomique, biologique et zoogéographique. pp. 461. Marseille: Université de Provence.

- Casanova JP, Judkins DC (1976) Les crustacés décapodes pélagiques de part et d'autre de Gibraltar. Relations entre les faunes Atlantique et Méditerranéenne. CIEM CM 16:1-6
- Casanova JP, Judkins DC (1977) Les décapodes pélagiques en Méditerranée. Répartition et secteurs faunistiques. Rapp comm int Mer Médit 24(10):125-127
- Chace Jr FA (1992) On the classification of the Caridea (Decapoda). Crustaceana 63:70–80
- Chan TY, Lei HC, Li CP, Chu KH (2010) Phylogenetic analysis using rDNA reveals polyphyly of Oplophoridae (Decapoda Caridea). Invert Syst 24:172–181
- Childress JJ, Nygaard M (1974) Chemical composition on buoyance of midwater crustaceans as function of depth of occurrence off Southern California. Mar Biol 27:225-238
- Christoffersen ML (1990) A new superfamily classification of the Caridea (Decapoda: Pleocyemata) based on phylogenetic pattern. Z zool Syst Evolut -forsch 28:94-106
- Clarke A, Holmes LJ (1987) Notes on the biology and distribution of *Pasiphaea* species from the southern ocean. Brit Antarct Surv B 74:17-30
- Colloca F, Carpentieri P, Balestri E, Ardizzone GD (2004) A critical habitat for Mediterranean fish resources: shelf-break areas with *Leptometra phalangium* (Echinodermata: Crinoidea). Mar Biol 145:1129-1142
- Company JB, Maiorano P, Tselepides A, Politou CY, Plaity W, Rotllant G, Sardà F (2004) Deep-sea decapod crustaceans in the western and central Mediterranean Sea: preliminary aspects of species distribution, biomass and population structure. Sci Mar 68(S3):73-86
- Company JB, Sará F, Puig P, Cartes JE, Palanques A (2003) Duration and timing of reproduction in decapod crustaceans of the NW Mediterranean continental margin: is there a general pattern? Mar Ecol-Prog Ser 2621:201-216

- Company JB, Sardá F (1997) Reproductive patterns and population characteristics in five deep-water pandalid shrimps in the Western Mediterranean along a depth gradient (150-1100 m). *Mar Ecol-Prog Ser* 148:49-58
- Company JB, Sardá F (2000) Growth parameters of deep-water decapod crustaceans in the Northwestern Mediterranean Sea: a comparative approach. *Mar Biol* 136:79-90
- Company JB, Cartes JE, Sardá F (2001) Biological patterns and near-bottom population characteristics of two pasiphaeid decapod crustacean species, *Pasiphaea sivado* and *P. multidentata*, in the north-western Mediterranean Sea. *Mar Biol* 139(1):61-73
- Correa C, Thiel M (2003) Mating systems in caridean shrimp (decapoda caridea) and their evolutionary consequences for sexual dimorphism and reproductive biology. *Rev Chil Hist Nat* 76:187-203
- Crosnier A, Forest J (1973) Les crevettes profondes de l'Atlantique oriental tropical. *Faune Trop* 19:1-409
- Cusson M, Bourget E (2005) Global patterns of macroinvertebrate production in marine benthic habitats. *Mar Ecol Prog Ser* 297:1–14
- Dalpadado P, Ellertsen B, Melle W, Skjoldal HR (1998) Summer distribution patterns and biomass estimates of macrozooplankton and micronekton in the Nordic seas. *Sarsia* 83:103-116
- De Grave, Fransen (2011) Carideorum catalogus (Crustacea: Decapoda). *Zool Med Leiden* 85, 588 p
- Delgado M, Rueda JL, Gil J, Burgos C, Sobrino I (2013) Spatial characterization of megabenthic epifauna of soft bottoms around mud volcanoes in the Gulf of Cádiz. *J Nat Hist* 47:1803-1831
- Doi S (1975) Notes on *Pasiphaea* in Toyama Bay. *Japan Sea Reg Fish Res News* 285:1-6

- D'Onghia G, Tursi A, Maiorano P, Matarrese A, Panza M (1998) Demersal fish assemblages from the bathyal grounds of the Ionian Sea (middle-eastern Mediterranean), *Ital J Zool* 65(S1):287-292
- d'Udekem d'Acoz C (1999) Inventaire et distribution des crustacés décapodes de l'Atlantique nord-oriental, de la Méditerranée et des eaux continentales adjacentes au nord de 25°N. *Patrimoines naturels (MNHN/SPN)* 40:1-383
- Eppley RW, Peterson BJ (1979) Particulate organic-matter flux and planktonic new production in the deep ocean. *Nature*, 282, 677-680.
- Estoup A, Largiadèr CR, Perrot E, Chourrout D (1996) Rapid one tube DNA 659extraction for reliable PCR detection of fish polymorphic marker and 660transgenes. *Mol Mar Biol Biotechnol* 5:295–298
- Estrada M (1996) Primary production in the northwestern Mediterranean. *Sci Mar* 60(S2):55-64
- Estrada M, Vives F, Alcaraz M (1985) Life and the productivity of the open ocean, in: Margalef, R. (Ed.), *Key Environments: Western Mediterranean*, Pergamon Press, Oxford, 148-197
- Fanelli E, Cartes JE (2004) Feeding habits of pandalid shrimps in the Alboran Sea (SW Mediterranean): influence of biological and environmental factors. *Mar Ecol-Prog Ser* 280:227-238
- Fanelli E, Cartes JE (2008) Spatio-temporal changes in gut contents and stable isotopes in two deep Mediterranean pandalids: influence on the reproductive cycle. *Mar Ecol-Prog Ser* 355:219–23
- Fanelli E, Colloca F, Ardizzone G (2007). Decapod crustacean assemblages off the West coast of central Italy (western Mediterranean). *Sci Mar* 71:19-28

- Fantucci MZ, Biagi R, Mantelatto FL (2009) Use of pleopod morphology to determine sexual dimorphism and maturity in hermit crabs: *Isocheles sawayai* as a model. *Helgol Mar Res* 63:169–175
- Fariña AC, Freire J, González Gurriarán E (1997) Megabenthic decapod crustacean assemblages on the Galician continental shelf and upper slope (north-west Spain). *Mar Biol* 127:419-434.
- Fasham MJR, Foxton P (1979) Zonal distribution of pelagic Decapoda (Crustacea) in the eastern north Atlantic and its relation to the physical oceanography. *J Exp Mar Biol Ecol* 37:225-253
- Fernández MV, Heras S, Maltagliati F, Turco A, Roldán MI (2011) Genetic structure in the blue and red shrimp *Aristeus antennatus* and the role played by hydrographical and oceanographical barriers. *421:163–171*
- Fiorentini L, Dremière PY, Leonori I, Sala A, Palumbo V (1999) Efficiency of the bottom trawl used for the Mediterranean international trawl survey (MEDITS). *Aquat Living Resour* 12(3):187-205
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R., 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3(5):294-299
- Frank TM (2000) Temporal resolution in mesopelagic crustaceans. *Phil Trans R Soc Lond B* 355: 1195-1198
- Franqueville C (1971) Macroplancton profond (Invertébrés) de la Méditerranée nord-occidentale. *Téthys* 3:11-56
- Froggia C, Giannini S (1982) Osservazioni sugli spostamenti verticali nictemerali si *Sergestes arcticus* Kroyer e *Sergia robusta* (Smith) (Crustacea, Decapoda, Sergestidae) nel



Mediterraneo Occidentale. Atti del Convegno delle Unità Operative afferenti ai sottoprogetti Risorse Biologiche e Inquinamento Marino 1:311-319

Galarza JA, Carreras-Carbonell J, Macpherson E, Pascual M, Roques S, Turner GF, Rico C (2009) The influence of oceanographic fronts and early-life-history traits on connectivity among littoral fish species. PNAS 106:1473-1478

García-Merchán VH, Robainas-Barcia A, Abelló P, Macpherson E, Palero F, García-Rodríguez M, Gil de Sola L, Pascual M (2012). Phylogeographic patterns of decapod crustaceans at the Atlantic–Mediterranean transition. Molecular Phylogenetics and Evolution 62:664-672

García-Rodríguez M, Abelló P, Fernández A, Esteban A (2011) Demersal assemblages on the soft bottoms off the Catalan-Levante coast of the Spanish Mediterranean. Journal of Marine Biology 2011:1-16

Garrison LP, Link JS (2000) Dietary guild structure of the fish community in the Northeast United States continental shelf ecosystem. Mar Ecol Prog Ser 202:231–240

Gartner JV, Sulak KJ, Ross SW, Necaie AM (2008) Persistent near-bottom aggregations of mesopelagic animals along the North Carolina and Virginia continental slopes. Mar Biol 153:825-841

Gayanilo FC Jr, Sparre P, Pauly D (2005) FAO-ICLARM Stock Assessment Tools II (FiSAT II). User's guide. FAO Computerized Information Series (Fisheries). No. 8, Revised version. Rome, FAO 168 p

Gibbons MJ, Macpherson E, Barangé M (1994) Some observations on the pelagic decapod *Pasiphaea semispinosa* Holthuis 1951 in the Benguela upwelling system. S Afr J Marine Sci 14:59-67

- González-Gurriarán E, Olaso I (1987) Cambios espaciales y temporales de los crustáceos decápodos de la plataforma continental de Galicia (NW de España). *Invest Pesq* 51:323-341
- Guerao G, Pertez-Baquera J, Ribera C (1994) Growth and reproductive biology of *Palaemon xiphias* Risso, 1816 (Decapoda: Caridea: Palaemonidae). *J Crustacean Biol*, 14: 280-288
- Guijarro B, Fanelli E, Moranta J, Cartes JE, Massutí E (2012) Small-scale differences in the distribution and population dynamics of pandalid shrimps in the western Mediterranean in relation to environmental factors. *Fish Res* 119-120:33-47
- Guijarro B, Massutí E, Moranta J, Cartes JE (2009) Short spatio-temporal variations in the population dynamics and biology of the deep-water rose shrimp *Parapenaeus longirostris* (Decapoda: Crustacea) in the western Mediterranean. *Sci Mar* 73:183-197
- Guijarro B, Massutí E, Moranta J, Díaz P (2008) Population dynamics of the red shrimp *Aristeus antennatus* in the Balearic Islands (western Mediterranean): Short spatio-temporal differences and influence of environmental factors. *J Marine Syst* 71:385-402
- Guijarro B, Tserpes G, Moranta J, Massutí E (2011) Assessment of the deep water trawl fishery off the Balearic Islands (western Mediterranean): from single to multi-species approach. *Hydrobiologia* 670:67-85
- Gurney R (1942) *Larvae of Decapod Crustacea*. London: Ray Society
- Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucl Acids Symp Ser* 41:95-98
- Hammer O, Harper D, Ryan P (2001) PAST: paleontological statistics software for education and data analysis. *Paleontología Electrónica* 4: 1-9
- Hassani S, Antoine L, Ridoux V (1997) Diets of Albacore, *Thunnus alalunga*, and Dolphins, *Delphinus delphis* and *Stenella coeruleoalba*, Caught in the Northeast Atlantic Albacore Drift-net Fishery: A Progress Report *J Northw Atl Fish Sci* 22:119–123

- Hayashi KI (1999) Crustacea Decapoda: Revision of *Pasiphaea sivado* (Risso, 1816) and related species, with descriptions of one new genus and five new species (Pasiphaeidae). In: A. Crosiner (ed.) Résultats des Campagnes MURSORSTOM Volume 20. Mémoires du Muséum national d'Histoire naturelle, Paris 180:267-302
- Hayashi KI (2006) A new species of the *Pasiphaea sivado* species group from Taiwan (Decapoda, Caridea, Pasiphaeidae). *Zoosystema* 28:341-346
- Herring PJ, Roe HSJ (1988) The photoecology of pelagic oceanic decapods. *Symp zool Soc Lond* 59:263-290
- Hewitt G, 2000. The genetic legacy of the Quaternary ice ages. *Nature* 405: 907-913
- Hicks, J. W. 1985. The breeding behaviour and migrations of the terrestrial crab *Gecarcoidea natalis* (Decapoda, Brachyura). *Australian Journal of Zoology* 33:127-142
- Hopkins TL, Flock ME, Gartner JV, Torres JJ (1994) Structure and trophic ecology of a low latitude midwater decapod and mysid assemblage. *Mar Ecol Prog Ser* 109:143-156
- Hopkins TS (1985) Physics of the sea. In: R. Margalef (ed.) *Key environments: Western Mediterranean*. Pergamon Press, New York, 100-125
- Hudson RR (2000) A new statistic for detecting genetic differentiation. *Genetics* 155:2011–2014
- Iwasaki N (1990) Pasiphaeid shrimps from the eastern atlantic and the Caribbean sea, with the description of a new species of *Pasiphaea* (crustacea: decapoda: pasiphaea). *Zoologische Mededelingen* 63(15):187-203
- Johnsen S (2001) Hidden in plain sight: The ecology and physiology of organismal transparency. *Biol Bull* 201:301-318
- Johnsen S (2005) The red and the black: Bioluminescence and the color of animals in the deep sea. *Integr Comp Biol* 45:234-246

- Johnson ML, Shelton PMJ, Gaten E, Herring PJ (2000) Relationship of dorsoventral eyeshine distributions to habitat depth and animal size in mesopelagic decapods. *Biol Bull* 199:6-13
- Judkins DC, Kensley B (2008) New genera in the family Sergestidae (Crustacea: Decapoda: Penaeidea). *Proc Biol Soc Wash* 121:72-84
- Kaartvedt S, Aksnes AL, Aadnesen A (1988) Winter distribution of macroplankton and micronekton in Masfjorden, western Norway. *Mar Ecol-Prog Ser* 45:45–55
- Karuppasamy PK, Menon NG, Nair KKC, Achuthankutty CT (2006) Distribution and abundance of pelagic shrimps from the deep scattering layer of the eastern Arabian Sea. *J Shellfish Res* 25:1013-1019
- Kensley B, Schotte M (2006) Pelagic shrimp (Crustacea: Decapoda) from shelf and oceanic waters in the southeastern Atlantic Ocean off South Africa. *Proc Biol Soc Wash* 119:384-394
- Kevrekidis K, Thessalou-Legaki M (2006) Catch rates, size structure and sex ratio of *Melicertus kerathurus* (Decapoda: Penaeidae) from an Aegean Sea trawl fishery. *Fish Res* 80:270–279
- Kim S (2005) Population structure, growth, mortality and size at sexual maturity of *Palaemon gravieri* (Decapoda: Caridea: Palaemonidae). *J Crust Biol* 25(2):226–232
- Klages M, Vopel K, Bluhm H (2001) Deep-sea food falls: first observation of a natural event in the Arctic Ocean *Polar Biol* 24:292-295
- Klingenberg CP (2011) MorphoJ: an integrated software package for geometric morphometrics. *Mol Ecol Resour* 11:353-357
- Klingenberg CP, McIntyre GS (1998) Geometric morphometrics of developmental instability: analyzing patterns of fluctuating asymmetry with procrustes methods. *Evolution* 52(5):1363-1375

- Komai T, Lin CW, Chan TY (2012) Bathypelagic shrimps of the genus *Pasiphaea* (Decapoda: Caridea: Pasiphaeidae) from waters around Taiwan, with descriptions of four new species. *Journal of Crustacean Biology* 32(2):295-325
- Koukouras A (2000) The pelagic shrimps (Decapoda, Natantia) of the Aegean Sea, with an account of the Mediterranean species. *Crustaceana* 73(7):801-814
- Koukouras A, Doulgeraki S, Kitsos MS (2000) Notes on the vertical distribution of pelagic shrimps (Decapoda, Natantia) in the Aegean Sea. *Crustaceana* 73(8):979-993
- Kronfeld-Schor N, Dayan T (2003) Partitioning of time as an ecological resource. *Annu Rev Ecol Evol Syst* 34:153-181
- Lagardère JP (1972) Recherches sur l'alimentation des crevettes de la pente continentale marocaine. *Téthys* 3:655-675
- Lagardère JP (1977) Recherches sur la distribution et sur l'alimentation des crustacés décapodes benthiques de la pente continentale du Golfe de Gascogne. Analyse des groupements carcinologiques. *Bull. Cent. Étud. Rech. Sci., Biarritz*, 11, 367-440.
- Li CP, Grave AD, Chan TY, Lei HC and Chu KH (2011) Molecular systematics of caridean shrimps based on five nuclear genes: Implications for superfamily classification. *Zoologischer Anzeiger* 250:270-279
- Li X (2006) Additional pandaloid shrimps from the south China Sea (Crustacea: Decapoda: Caridea), with description of one new species. *Raffles B Zool* 54(2):361-372
- Librado P, Rozas J (2009) DnaSP 5: A software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* 25:451-452
- Lleonart J, Salat J (1992) VIT, programa de análisis de pesquerías. *Informes Técnicos de Scientia Marina*, 168-169:1-116

- López RI (1997) Crustáceos decápodos capturados durante las campañas del IEO ARSA 0393 y ARSA 1093 en el golfo de Cádiz: distribución batimétrica. Publ Espec Inst Esp Oceanogr 23:199-206
- López-Jurado JL, Marcos M, Monserrat S (2008) Hydrographic conditions affecting two fishing grounds of Mallorca island (Western Mediterranean): during the IDEA Project (2003–2004). J Mar Syst 71:303-315
- Lorentsen SH, Klages N and Røv N (1998) Diet and prey consumption of Antarctic petrels *Thalassoica antarctica* at Svarthamaren, Dronning Maud Land, and at sea outside the colony. Polar Biology 19: 414-420
- Luttikhuisen P, Campos J, van Bleijswijk J, Peijnenburg K, van der Veer H (2008) Phylogeography of the common shrimp, *Crangon crangon* (L.) across its distribution range. Mol Phylogenet Evol 46:1015–1030
- Maggs CA, Castillo R, Foltz D, Henzler C, Jolly MT, Kelly J, Olsen J, Perez KE, Stam W, Väinölä R, Viard F and Wares J (2008) Evaluating signatures of glacial refugia for north Atlantic benthic marine taxa. Ecology 89(11):S108-S122
- Massutí E, Monserrat S, Oliver P, Moranta J, López-Jurado JL, Marcos M, Hidalgo M, Guijarro B, Carbonell A, Pereda P (2008) The influence of oceanographic scenarios on the population dynamics of demersal resources in the western Mediterranean: Hypothesis for hake and red shrimp off Balearic Islands. J Mar Syst 71:421–438
- Massutí E, Olivar MP, Monserrat S, Rueda L, Oliver P (2013) Towards the understanding of the influence of the environmental conditions on the demersal resources and ecosystems in the western Mediterranean: Motivations, aims and methods of the IDEADOS project. J Mar Syst (in press).

- Matthews JBL, Pinnoi S (1973) Ecological studies on the deep-water pelagic community of Korsfjorden, western Norway. The species of *Pasiphaea* and *Sergestes* (Crustacea Decapoda) recorded in 1968 and 1969. *Sarsia* 52:123-144
- Matzen da Silva J, Creer S, dos Santos A, Costa AC, Cunha MR, Costa FO, Carvalho GR (2011) Systematic and evolutionary insights derived from mtDNA COI barcode diversity in the Decapoda (Crustacea: Malacostraca). *PLoS ONE*, 6, e19449:1-15
- Mauchline J, Gordon JDM (1991) Oceanic pelagic prey of benthopelagic fish in the benthic boundary layer of a marginal oceanic region. *Mar Ecol Prog Ser* 74:109-115
- Maurin C (1961) Répartition des crevettes profondes sur les côtes sud du bassin occidental de la Méditerranée et dans la région atlantique ibéro-marocaine. *Rapp. Comm. int Mer Médit.* 16:529532
- Maynou F, Cartes JE (1998) Daily ration estimates and comparative study of food consumption in nine species of deep-water decapod crustaceans of the NW Mediterranean. *Mar Ecol Prog Ser* 171:221-231
- Maynou F, Cartes JE (1998) Daily ration estimates and comparative study of food consumption in nine species of deep-water decapod crustaceans of the NW Mediterranean. *Mar Ecol Prog Ser* 171:221-231
- Maynou F, Conan GY, Cartes JE, Company JB, Sardà F (1996) Spatial structure and seasonality of decapod crustacean populations on the northwestern Mediterranean slope. *Limnol Oceanogr* 41(1):113-125
- McGaw IJ, Naylor E (1992) Salinity preference of the shore crab *Carcinus maenas* in relation to coloration during intermoult and to prior acclimation. *J Exp Mar Biol Ecol* 155:145-159

- McManus MA, Woodson CB (2012) Plankton distribution and ocean dispersal. *J Exp Biol* 215:1008-1016
- Millot C (2005) Circulation in the Mediterranean Sea: evidences, debates and unanswered questions. *Sci Mar* 69S1: 5-21
- Millot Pannacciulli C, Taupier-Letage I (2005) Circulation in the Mediterranean Sea. *Hdb Env Chem*, 5(K): 29-66
- Montserrat S, López-Jurado JL, Marcos M (2008) A mesoscale index to describe the regional circulation around the Balearic Islands. *J Mar Syst* 71:413-420
- Monteiro P, Araújo A, Erzini K, Castro M (2001) Discards of the Algarve (southern Portugal) crustacean trawl fishery. *Hydrobiologia* 449:267-277
- Morales-Nin B, Maynou F, Sardà F, Cartes JE, Moranta J, Massutí E, Company JB, Rotllant G, Bozzano A, Stefanescu C (2003) Size influence in zonation patterns in fishes and crustaceans from deep-water communities of the western Mediterranean. *J Northw Atl Fish Sci* 31:413-430
- Moranta J, Quetglas A, Massutí E, Guijarro B, Hidalgo M, Díaz P (2008) Spatio-temporal variations in deep-sea demersal communities off the Balearic Islands (western Mediterranean). *J Mar Syst* 71:346-366
- Muñoz I, García-Isarch E, Sobrino I, Burgos C, Funny R, González-Porto M (2012) Distribution, abundance and assemblages of decapod crustaceans in waters off Guinea-Bissau (north-west Africa). *J Mar Biol Ass UK* 92:475-494
- Nanjo N (2007) Feeding habits of the glass shrimp *Pasiphaea japonica* in Toyama Bay of the Sea of Japan. *Crust Res* 36:45-51
- Nanjo N and Ohtomi J (2009) Reproductive biology of *Pasiphaea japonica* females in Toyama Bay, central Japan. *Fisheries Sci* 75:1189-1195



- Nanjo N, Konishi K (2009) Complete larval development of the japonise glass shrimp *Pasiphaea japonica* Omori, 1976 (Decapoda: Pasiphaeidae) undr laboratory conditions. Crustacean research 38:77-89
- Naylor E (2010) Chronobiology of marine organisms. Cambridge University Press, Cambridge
- Oh CW, Hartnoll RG and Nash RDM (1999) Population dynamics of the common shrimp, *Crangon crangon* (L.), in Port Erin Bay, Isle of Man, Irish Sea. ICES Journal of Marine Science 56:718–733
- Oh CW, Kim JY, Jeong IJ, Suh HL, Cho YK (2006) Reproduction and population dynamics of *Leptochela gracilis* (Decapoda : Pasiphaeidae) on the western coast of Korea, Yellow Sea. J Mar Biol Ass UK 86:113-120
- Oh CW, Suh HL, Park KY, Ma CW, Lim HS (2002) Growth and reproductive biology of the fresh water shrimp *Exopalaemon modestus* (Decapoda: Palaemonidae) in a lake of Korea. J Crustacean Biol 22:357-366
- Olivar MP, Bernal A, Molí B, Peña M, Balbín R, Castellón A, Miquel J, Massutí E (2012) Vertical distribution, diversity and assemblages of mesopelagic fishes in the western Mediterranean. Deep-Sea Res I 62:53–69
- Omori M (1976) The glass shrimp, *Pasiphaea japonica* sp. nov. (caridea, pasiphaeidae), a sibling species of *Pasiphaea sivado*, with notes on its biology and fishery in Toyama Bay, Japan. Bull Natn Sci Mus Ser A (zool) 2 (4):249-266
- Omori M, Gluck D (1979) Life history and vertical migration of the pelagic shrimp *Sergestes similis* off the southern California coast. Fish Bull 77:183-198
- Orsi Relini L, Pinca S (1990) Reproductive patterns of *Pasiphaea sivado* in the Ligurian Sea. Rapp Comm Int Mer Médit 32:223

- Orsi-Relini L, Relini G (1990) The glass shrimp *Pasiphaea sivado* in the food chains of the Ligurian Sea. In: Barnes M, Gibson RN (eds) Trophic relationships in the marine environment (Proc. 24th European Marine Biology Symposium). Aberdeen University Press, Aberdeen, pp 334-346
- Palero F, Abelló P, Macpherson E, Beaumont M, Pascual M (2011) Effect of oceanographic barriers and overfishing on the population genetic structure of the European spiny lobster (*Palinurus elephas*). *Biol J Linn Soc* 104:407-418
- Palero F, Abelló P, Macpherson E, Gristina M, Pascual M (2008) Phylogeography of the European spiny lobster (*Palinurus elephas*): Influence of current oceanographical features and historical processes. *Mol Phylogenet Evol* 48:708-717
- Palero F, Lopes J, Abelló P, Macpherson E, Pascual M, Beaumont MA (2009) Rapid radiation in spiny lobsters (*Palinurus* spp) as revealed by classic and ABC methods using mtDNA and microsatellite data. *BMC Evol Biol* 9:1-13
- Palumbi SR, Benzie J (1991) Large mitochondrial DNA differences between morphologically similar Penaeid shrimp. *Molec Mar Biol Biotech* 1: 27-34
- Pannacciulli FG, Bishop JDD, Hawkins SJ (1997) Genetic structure of populations of two species of *Chthamalus* (Crustacea: Cirripedia) in the north-east Atlantic and Mediterranean. *Mar Biol* 128:73–82
- Papiol V, Cartes JE, Fanelli E, Maynou F (2012) Influence of environmental variables on the spatio-temporal dynamics of benthic-pelagic assemblages in the middle slope of the Balearic Basin (NW Mediterranean). *Deep Sea Res I* 61:84-99
- Patarnello T, Volckaert F, Castilho R (2007) Pillars of Hercules: is the Atlantic–Mediterranean transition a phylogeographic break? *Mol Ecol* 16:4426– 4444

- Perez-Losada M, Guerra A, Carvalho GR, Sanjuan A, Shaw PW (2002) Extensive population subdivision of the cuttlefish *Sepia officinalis* (Mollusca: Cephalopoda) around the Iberian Peninsula indicated by microsatellite DNA variation. *Heredity* 89:417–424
- Pineda J, Hare J A and Sponaugle S, 2007. Larval transport and dispersal in the coastal ocean and consequences for population connectivity. *Oceanography* 20:22-39
- Pinot JM, López-Jurado JL, Riera M (2002) The CANALES experiment (1996–1998). Interannual, seasonal, and mesoscale variability of the circulation in the Balearic Channels. *Prog Oceanogr* 55:335–370
- Politou CY, Karkani M, Dokos J (2000) Distribution of decapods caught during MEDITS surveys in Greek waters. *Actes de Colloques IFREMER* 26:196-207
- Polunin NVC, Morales-Nin B, Pawsey WE, Cartes JE, Pinnegar JK, Moranta J (2001) Feeding relationships in Mediterranean bathyal assemblages elucidated by stable nitrogen and carbon isotope data. *Mar Ecol Prog-Ser* 220:13-23
- Posada D (2008) jModelTest: phylogenetic model averaging. *Mol Biol Evol* 25:1253-1256
- Queiroga H (1993) An analysis of the size structure of *Carcinus maenas* (L.) in Canal de Mira (Ria de Aveiro, Portugal) using the probability paper method. *Bios (Thessaloniki)* 1:89-106
- Radulovici AE, Sainte-Marie B and Dufresne F (2009) DNA barcoding of marine crustaceans from the Estuary and Gulf of St Lawrence: a regional-scale approach. *Mol Ecol Resour* 9(S1):181-187
- Ramirez-Llodra E, Company JB, Camps M, Rotllant G (2007) Spatio-temporal variations in reproductive patterns and population structure of *Pasiphaea multidentata* (Decapoda: Caridea) in the Blanes canyon and adjacent margin, Northwestern Mediterranean Sea. *Mar Ecol Evol Perspect* 28:470-479

- Ramón M, Abelló P, Ordinas F, Massutí E (2013) Epibenthic communities in two contrasting areas of the Balearic Islands. *J Mar Syst* (in press).
- Reid DG, Abelló P, Kaiser MJ, Warman CG (1997) Carapace colour, intermoult duration and the behavioural and physiological ecology of the shore crab *Carcinus maenas*. *Estuar Coast Shelf S* 44:203-211
- Reid DG, Abelló P, Warman CG, Naylor E (1994) Size-related mating success in the shore crab *Carcinus maenas* (Crustacea: Brachyura). *J Zool Lond* 232:397-407
- Reid SB, Hirota J, Young RE, Hallacher LE (1991) Mesopelagic-boundary community in Hawaii - micronekton at the interface between neritic and oceanic ecosystems. *Mar Biol* 109:427-440
- Reuschel S, Schubart CD (2006) Phylogeny and geographic differentiation of Atlanto–Mediterranean species of the genus *Xantho* (Crustacea: Brachyura: Xanthidae) based on genetic and morphometric analyses. *Mar Biol* 148: 853–866
- Robertis A (2002) Small-scale spatial distribution of the euphausiid *Euphausia pacifica* and overlap with planktivorous fishes. *J Plankton Res* 24:1207-1220
- Rohlf FJ (2010a) TpsDig, digitize landmarks and outlines, version v2.16. Department of Ecology and Evolution, State University of New York at Stony Brook
- Rohlf FJ (2010b) TpsRelw, relative warps analysis, version 1.49. Department of Ecology and Evolution, State University of New York at Stony Brook
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Syst Biol*. 61(3):539-42
- Rosecchi E, Tracey DM, Webber WR (1988) Diet of orange roughy, *Hoplostethus atlanticus* (Pisces: Trachichthyidae) on the Challenger Plateau, New Zealand *Mar Biol* 99:293-306

- Rufino MM, Abelló P, Yule AB (2006) Geographic and gender shape differences in the carapace of *Liocarcinus depurator* (Brachyura: Portunidae) using geometric morphometrics and the influence of a digitising method. *J Zool* 269: 458-465
- Rufino MM, Abelló P, Yule AB, Torres P (2005) Geographic, bathymetric and inter-annual variability in the distribution of *Liocarcinus depurator* (Brachyura: Portunidae) along the Mediterranean coast of the Iberian Peninsula. *Sci Mar* 69:503-518
- Sabatés A, Olivar MP, Salat J, Palomera I, Alemany F (2007) Physical and biological processes controlling the distribution of fish larvae in the NW Mediterranean. *Prog Oceanogr* 74:355-376
- Sala-Bozano M, Ketmaier V, Mariani S (2009) Contrasting signals from multiple 800markers illuminate population connectivity in a marine fish. *Mol Ecol* 18(801)4811–4826
- Salat J (1995) The interaction between the Catalan and Balearic currents in the southern Catalan Sea. *Oceanol Acta* 18:227-234
- Sampedro MP, González-Gurriarán E (2004) Aggregating behaviour of the spider crab *Maja squinado* in shallow waters. *J Crustacean Biol* 24:168-177
- Santos A, Moreia A (2003) On the occurrence of *Sergia grandis* and *Pasiphaea hoplocerca* (Sund, 1920) and *Pasiphaea hoplocerca* Chace, 1940 (Decapoda) off the southern coast of Portugal. *Crustaceana* 75(9):1061-1065
- Sardà F, Compay JB, Costa C (2005) A morphological approach for relating decapod crustacean cephalothorax shape with distribution in the water column. *Mar Biol* 147:611–61
- Schunter C, Carreras-Carbonell J, Planes S, Sala E, Ballesteros E, Zabala M, Harmelin JG, Harmelin-Vivien M, Macpherson E, Pascual M (2011) Genetic connectivity patterns in an endangered species: The dusky grouper (*Epinephelus marginatus*). *J Exp Mar Biol Ecol* 401(1):126-133

- Serejo CS, Young PS, Cardoso IC, Tavares C, Rodrigues C, Almeida TC (2007) Abundância, diversidade e zonation dos crustáceos no talude da costa central do Brasil (11°-22° S) coletados pelo programa REVIZEE/ Score central: Prospecção pesqueira. In: Costa PA, Olavo G, Martins AS (Eds.) Biodiversidade da fauna marinha profunda na costa central brasileira. Rio de Janeiro, Museo Nacional, (Série Livros 24)133–162
- Serrano A, Velasco F, Olaso I and Sánchez F (2003) Macrobenthic crustaceans in the diet of demersal fish in the Bay of Biscay in relation to abundance in the environment. *Sarsia*, 88: 36-48
- Silva L, Vila Y, Torres MA, Sobrino I, Acosta JJ (2011) Cephalopod assemblages, abundance and species distribution in the Gulf of Cadiz (SW Spain). *Aquatic Living Resources* 24:13-26
- Simão DS, Carbonnel A, Abelló P (submitted). Pasiphaeid shrimp populations in the western Mediterranean: geographical variability in distribution and population patterns.
- Sivertsen E, Holthuis LB (1956) Crustaca decapoda (The peneidea and Stenopodidea excepted). *Rep Sci Res "Michael Sars" North Atlant. Deep-sea Exped. 1910.* 5(12):1-54
- Skjoldal HR, Bamstedt U (1977) Ecobiochemical studies on deep-water pelagic community of Korsfjorden, western Norway - adenine-nucleotides in zooplankton. *Mar Biol* 42:197-211
- Sokal RR, Rohlf FJ (1981) *Biometry*, 2nd edition. W.H. Freeman, Co., New York. 937p
- Stephensen K (1923) *Decapoda Macrura, excl. Sergestidae*. Report on the Danish oceanographic expeditions 1908-10 to the Mediterranean and adjacent seas 2:1-85
- Sumida PYG, Pires-Vanin AMS (1997) Benthic Associations of the Shelfbreak and Upper Slope off Ubatuba-SP, South-eastern Brazil. *Estuar Coast Shelf S* 44(6):779-784

- Sund O (1913) The glass shrimp (*Pasiphaea*) in the northern waters. Bergens Mus Aarb 1912(6):1-18
- Suntsov A, Domokos R (2013) Vertically migrating micronekton and macrozooplankton communities around Guam and the Northern Mariana Islands. Deep-Sea Research Part I-Oceanographic Research Papers 71,:113-129
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, and Kumar S (2011) MEGA5: Molecular Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods. Mol Biol Evol 28:2731-2739
- Tavares, CR and Cardoso IA (2006) Deep sea Pasiphaeidae (crustacean: Decapoda: Caridea) from off the Brazilian central between 11° and 22° S, collected by the Revizee Program. Zootaxa 1174:23-39
- Tecchio S, Coll M, Christensen V, Company JB, Ramírez-Llodra E and Sardà F (2013) Food web structure and vulnerability of a deep-sea ecosystem in the NW Mediterranean Sea. Deep-Sea Res 1(75):1–15
- Tecchio S, Ramirez-Llodrà E, Sardá F, Company JB (2011) Biodiversity of deep-sea demersal megafauna in western and central Mediterranean basins. Sci Mar 75(2):341-350
- Thompson JR (1967) Comments on phylogeny of section Caridea (Decapoda Natantia) and the phylogenetic importance of the Oplophoroidea. In: Proceedings of the Symposium on Crustacea, Marine Biological Association of India 1:314–326
- Tintoré J, La Violette PE, Blade I, Cruzado A (1988) A study of an intense density front in the eastern Alboran Sea: the Almeria-Oran front. J Phys Oceanogr 18:1384-1397
- Tizperman E, Malanotte-Rizzoli P (1991) The climatological seasonal circulation of the Mediterranean Sea. Journal of Marine Research 49: 411-434

- Torres AP, Dos Santos A, Balbín R, Alemany F, Massutí E, Reglero P (2013) Decapod crustacean larval communities in the western Mediterranean: Composition and vertical distribution patterns. *J Marine Syst* (in press).
- Torres AP, Reglero P, Balbín R, Urtizberea A, Alemany F (2011) Coexistence of larvae of tuna species and other fish in the surface mixed layer in the NW Mediterranean. *J Plankton Res* 33:1793-1812
- Ungaro N, Marano C A, Ceriola L, Martino M (2005) Distribution of demersal crustaceans in the southern Adriatic Sea. *Acta Adriatica* 46:27-40
- Vafidis D, Leontarakis PK, Dailianis A, Kallianiotis A (2008) Population characteristics of four deep-water pandalid shrimps (Decapoda: Caridea) in the northern Aegean Sea (NE Mediterranean). *J Nat Hist* 42:2079-2093
- Vafidis D, Leontarakis PK, Dailianis A, Kallianiotis A (2004) Growth parameters of the most abundant pandalid shrimps (Decapoda: Caridea) from the northern Aegean Sea. *Rapp Comm int Mer Médit* 37:453
- Vafidis D, Politou CY, Carbonell A, Company JB (2005) A review of the biology and fisheries of the genus *Plesionika* Bate, 1888 (Decapoda, Caridea, Pandalidae) in European waters. *Crustaceana* 78:335-352
- Valls M, Quetglas A, Ordines F, Moranta J (2011) Feeding ecology of demersal elasmobranchs from the shelf and slope off the Balearic Sea (western Mediterranean). *Sci Mar* 75:633-639
- Vargas-Yáñez M, Sabatés A (2007) Mesoscale high-frequency variability in the Alboran Sea and its influence on fish larvae distributions. *J Marine Syst* 68:421-438
- Vereshchaka AL (1994) North Atlantic and Caribbean species of *Sergia* (Crustacea, Decapoda, Sergestidae) and their horizontal and vertical distribution. *Steenstrupia* 20:73-95



- Vereshchaka AL (1995) Distribution of the benthopelagic shrimps over the continental slopes and seamounts of the western Indian ocean. *Oceanology* 35(4):528-531
- Vestheim H, Kaartvedt S (2009) Vertical migration, feeding and colouration in the mesopelagic shrimp *Sergestes arcticus*. *J Plankton Res* 31:1427-1435
- Viegas I, Martinho F, Neto J, Pardal M (2007) Population dynamics, distribution and secondary production of the brown shrimp *Cragon cragon* (L.) in southern European estuary. Latitudinal variations. *Scientia Marina* 71(3):451-460
- Viscosi V, Cardini A (2011) Leaf Morphology, Taxonomy and Geometric Morphometrics: A Simplified Protocol for Beginners. *PLoS ONE* 6(10):e25630
- Wenner AM (1972) Sex ratio as a function of size in marine Crustacea. *Am Nat* 106:321-350
- Williams A, Koslow JA (1997) Species composition, biomass and vertical distribution of micronekton over the mid-slope region off southern Tasmania, Australia. *Mar Biol* 130:259-276
- Williamson DI, 1960. Larval stages of *Pasiphaea sivado* and some other Pasiphaeidae (Decapoda). *Crustaceana* 1:331-341
- Würtz M (2010) Mediterranean Pelagic Habitat: Oceanographic and Biological Processes, An Overview.
- Xia X (2013) DAMBE5: A comprehensive software package for data analysis in molecular biology and evolution. *Mol Biol Evol* 30(7):1720-1728
- Xia X, Xie Z, Salemi M, Chen L, Wang Y (2003) An index of substitution saturation and its application. *Mol Phylogenet Evol* 26:1–7
- Yamasaki M (1988) The ecological study of mantis shrimp *Oratosquilla oratoria* (De Haan) with reference to its bio-production processes. *Bull Seikai Reg Fish Res Lab* 66:69-100

## Bibliography

- Zane L, Ostellari A, Maccatrozzo, Bargelloni L, Cuzin-Roudly J, Buchholz F, Patarnello T (2000) Genetic differentiation in pelagic crustacean (*Meganyciophanes norvegica*: Euphausiacea) from the east Atlantic and the Mediterranean Sea. *Mar Biol* 136:191-199
- Zariquiey Álvarez R (1946) *Crustáceos Decápodos Mediterráneos*. Barcelona: Instituto Español de Estudios Mediterráneos. 217 p
- Zariquiey Álvarez R (1957) Decápodos españoles XIII - Las Pasiphaeas del Mediterráneo occidental. *Trabajos del Museo de Zoología Nueva Serie Zoológica* 2:1-41
- Zariquiey Álvarez, R (1968) Crustáceos decápodos ibéricos. *Invest Pesq* 32:1-510
- Zeng C, Naylor E (1997) Rhythms of larval release in the shore crab *Carcinus maenas* (Decapoda: Brachyura) *J Mar Biol Ass UK* 77:451-461