

UNIVERSITAT DE BARCELONA  
FACULTAT DE FARMÀCIA  
DEPARTAMENT DE FARMÀCIA I TECNOLOGIA FARMACÈUTICA

ESTUDI DE LA FORMACIÓ DE NANO-EMULSIONS  
DE FASE EXTERNA AQUOSA I SOLUBILITZACIÓ DE  
FÀRMACS LIPÒFELS

Núria Sadurní Gràcia, 2006

## A

Agache, P., Girardot, I., Bernengo, J.C. Optical properties of the skin, in *Cutaneous Investigation in Health and Disease: Noninvasive Methods and Instrumentation*. Marcel Dekker, 241–274, 1989.

Alberti, I., Kalia, Y.N., Naik, A., Bonny, J.D., Guy, R.H., In vivo assessment of enhanced topical delivery of terbinafine to human stratum corneum, *J. Controlled Release*, v. 71(3), 319–327, 2001.

Albery, W.J., Guy, R.H., Percutaneous absorption: transport in the dermis. *International Journal of Pharmaceutics*, v. 15, 125–148, 1983.

Albery, W.J., Hadgraft, J. Percutaneous absorption: theoretical description. *J. Pharm. Pharmacol.*, v. 31, 129–139, 1979.

Alexandridis, P., Olsson, U., Lindman, B., A record nine different phases (four cubic, two hexagonal and one lamellar lyotropic liquid crystalline and two micellar solutions) in a ternary isothermal system of an amphiphilic block copolymer and selective solvents (water and oil), *Langmuir*, v. 14(10), 2627–2638, 1998.

Amselem, S., Friedman, D., *Submicron emulsions as drug carriers for topical administration*, in: S. Benita (Ed.), *Submicron emulsions in drug targeting and delivery*, Harwood Academic Publishers, London, 153–173, 1998.

Anderberg, E.K., Nystrom, C., Artursson, P., *Proc. Int. Symp. Controlled Release Bioact. Mater.*, v. 17, 345, 1990.

Andersen, K.E., Nielsen, R., Lipistick dermatitis related to castor oil, *Contact Dermatitis*, v. 11(4), 253–254, 1984.

Antonietti, M., Landfester, K., Polyreactions in miniemulsions, *Prog. Polym. Sci.*, v. 27, 689–757, 2002.

Asua, J.M., Miniemulsion polymerization, *Prog. Polym. Sci.*, v. 27, 1283–1346, 2002.

Attwood, D., Ktistis, G., A light scattering study on oil-in-water microemulsions. *Int. J. Pharm.* 52, 165–171, 1989.

## B

- Bhalani, V.T., Satishchandra, S.P. US *Patent* 5858401 A, 1999.
- Baluom, M., Friedman, D.I., Rubinstein, A., Absorption enhancement of calcitonin in the rat intestine by carbopol-containing submicron emulsions, *Int. J. Pharm.*, v. 154, 235–243, 1997.
- Bar-Ilan, A., Baru, H., Beilin, M., Friedman, D., Amselem, S., Neumann, R. *Reg. Immunol.*, vol. 6., 166–168, 1994.
- Barnette, D.T., Schork, F.J., Continuous Miniemulsion Polymerization, *Chem. Eng. Prog.*, v. 83, 25–30, 1987.
- Barry, B.W., *Dermatological Formulations*, Marcel Dekker, New York, NY, 1–14, 1983.
- BASFa. Butlletí tècnic del Cremophor® EL. 2005.
- BASFb. Butlletí tècnic del Solutol® HS 15. 2005.
- Becher, P., Emulsiones, Teoría y Práctica, Ed. Blume, 1972.
- Bech, H., Bracher, M., Faller, C., Hofer, H. *Prediction of Percutaneous Penetration: Methods, Measurements and Modelling*, London: IBC Technical Services, v. 2, 441–450, 1991.
- Beckman, M., Eckhardt, B., Kaiser, B., Goering, S. *Ger Offen DE 196118809 C1*, 1997.
- Bedri, E., Sudol, E.D., Dimonie, V.L., El-Aasser, M.S., Encapsulation of Inorganic Particles via Miniemulsion Polymerization, *Macromol. Symp.*, 155, 181–198, 2000.
- Beilin, M., Bar-Ilan, A., Amselem, S., Schwarz, J., Yogev, A. Neumann, R., Ocular retention time of submicron emulsion (SME) and the miotic response To pilocarpine delivered in SME, *Invest. Ophthalmol. Vis. Sci.*, v. 36, S166, 1999.
- Benavides, T., Martínez, V., Mitjans, M., Infante, M.R., Moran, C., Clapes, P., Clothier, R., Vinardell, M.P., Assessment of primary eye and skin irritants by in vitro cytotoxicity and phototoxicity models: an in vitro approach of new arginine-based surfactant-induced irritation, *Toxicology*, v. 201, 87–93, 2004.
- Benita, S. *Submicron emulsions in drug targeting and delivery*. Benita, S., ed. 9, 1–19. 1998.
- Benita, S., Friedman, D., Weinstock, M., Physostigmine emulsion: a new injectable controlled release delivery system, *Int. J. Pharm.* V. 30, 47–55, 1986.

Benita, S., Levy M. J., Submicron emulsions ad colloidal drug carriers for intravenous administration: comprehensive physicochemical characterization. *J. Pharm. Sci.*, v. 82, 1069–1079, 1993.

Benita, S., Magalhaes, N.S., Cave, G., Seiller, M., The stability and in vitro release kinetics of a clofibríde emulsion. *Int. J. Pharm.*, v. 76, 225–237, 1991.

Bhalla, H.L., Menon, M.R., Gopal, N.G.S., Radiation Sterilization of Polyethylene glycols, *Int. J. Pharmaceutics*, 351–355, 1983.

Bivas–Benita, M., Oudshoorn, M., Romeijn, S., van Meijgaarden, K., Koerten, H., Van der Meulen, H., Gregory, L., Ottenhoff, T., Benita, S., Junginger, H., Borchard, G., Cationic submicron emulsions for pulmonary DNA immunization, *J. Control. Rel.*, v. 100, 145–155, 2004.

Blume, A., Jansen, M., Ghyczy, M., Gareiss, J., Interaction of phospholipid liposomes with lipid model mixtures for stratum corneum lipids. *Int. J. Pharmac.*, v. 99, 219–228, 1993.

Boelsma, E., Tanojo, H., Boddé, H.E., Ponec, M., Assessment of the potential irritancy of oleic acid on human skin: evaluation *in vitro* and *in vivo*, *Toxicology in Vitro*, v. 10, 729–742, 1996.

Boinpally, R.R., Zhou, S.–L., Poondru, S., Devraj, G., Jasti, B.R., Lecithin vesicles for topical delivery of diclofenac, *European Journal of Pharmaceutics and Biopharmaceutics*, v.56 (3), 389–392, 2003.

Bourrel, M., Schechter, R.S. *Microemulsion and Related System*, Surfactant Science Series, Marcel Dekker, New York, v. 30, 1988.

Brain, K.R., Walters, K.A., Watkinson, A.C., Methods for studying percutaneous absorption, *Dermatological and Transdermal Formulations*, v. 119, 197–270, 2002.

Bru, M.N., Guillon, X., Breton, P., Couvreur, P., Lescure, F., *French Patent 2755136*, 1998.

Buszello, K., Harnisch, S., Müller, R.H., Müller, B.W., The influence of alkali fatty acids on the properties and the stability of parenteral O/W emulsions modified with Solutol HS 15, *Eur. J. Pharm. Biopharm.*, v. 49, 143–149, 2000.

## C

Catálogo de Especialidades Farmacéuticas, Colección Consejo Plus, 2005.

Chung, H., Kim, T.W., Kwon, M., et al., Oil components modulate physical characteristics and function of the natural oil emulsions as drug or gene delivery system, *Journal of Controlled release*, v. 71, 339–350, 2001.

Clarys, P., Barel, A.O., Gabard, B. *Prediction of Percutaneous Penetration*. (Brain, KR, James, VJ, Walters, KA, eds.), v.3b, 502–509, 1993.

Clarys, P., Buchet, E., Barel, A.O., *Prediction of Percutaneous Penetration*, (Scott, R.C., Guy, RH, Hadgraft, J, Boddé, H.E., eds.), v. 2, 46–59, 1991.

Clemens, T., Boehm, A., Kielhorn–Bayer, S., Nanocolor–ants: Pigments with Dyestuff Properties. Polymer Preprints, *Polym. Prep. (Am Chem. Soc. Div. Polym. Chem.)* v. 41(1), 24–25, 2000.

Clowes, H.M., Smith, F.M., Scott, R.C. *Preparation of intact epidermal membranes from whole skin: an “in-depth” assessment of the dermatome*. (Brain, K.R., James, V.J., Walters, K.A., eds.) Predictions of percutaneous penetration, STS Publishing, v. 3b., 123–127, 1993.

Code of Federal Regulations, Title 49 Part 173.210, 1975, and Title 16 Part 1500.41, 1976.

Constantinides, P.P., Lambert, K.J., Tustian, A.K., Schneider, B., Lalji, S., Ma, W., Wentzel, B., Kessler, D., Worah, D., Quay, C., Formulation development and antitumor activity of a filter-sterilizable emulsion of paclitaxel, *Pharmaceutical Research*, v. 17, 2, 175–182, 2000.

Coquette, A., Berna, N., Vandenbosch, A., Rosdy, M., De Weber, B., Poumay, Y., Analysis of interleukin-1 $\alpha$  (IL-1 $\alpha$ ) and interleukin-8 (IL-8) expression and release in in vitro reconstructed human epidermis for the prediction of in vivo skin irritation and/or sensitization, *Toxicol. In Vitro*, v. 17, 311–321, 2003.

Council Directive (76/768/EEC), *Official Journal of the European Communities*, L 262/169, 27th July, 1976.

Curdy, C., Naik, A., Kalia, Y.N., Alberti, I., Guy, R.H., Non-invasive assessment of the effect of formulation excipients on stratum corneum barrier function in vivo, *Int. J. Pharmaceutics*, v. 271, 251–256, 2004.

## D

Dahl, G.B., Svensson, L., Kinnander, N.J.G., Stability of vitamins in soybean oil fat emulsion under conditions simulating intravenous feeding of neonates and children, *J. Parenter. Enteral Nutr.*, v. 18(3), 2234–2239, 1994.

Danielsson I., Lindman B., The definition of microemulsion. *Colloids and Surfaces*, v.3 (4), 391–392, 1981.

Dardel, O.V., Mebius, C., Mossberg, T., Svenson, B., Fat emulsion as a vehicle for diazepam. A study of 9492 patients, *J. Anaesth.*, v. 55, 41–47, 1983.

Davis, A.F., Hadgraft, J. Effect of supersaturation on membrane transport: 1. Hydrocortisone acetate. *Int. J. Pharm.*, vol. 76– 1–8. 1991.

Degussa BioActives. Butlletí tècnic de l'Epikuron 200. 2002.

Delgado, I., Solubilización de anestésicos locales en microemulsiones, *Tesi doctoral*, Universitat de Barcelona, 1998.

Delgado–Charro, M.B., Iglesias–Vilas, G., Blanco–Méndez, J., López–Quintela, A., Marty, J.–P., Guy, R.H., Delivery of a hydrophilic solute through the skin from novel microemulsion systems, *Eur. J. of Pharm. and Biopharm.*, v.43, 37–42, 1997.

Deminiere, B., Colin, A., Leal Calderon, F., Bibette, J., Lifetime and destruction of concentrated emulsions undergoing coalescence. (Binks, B.P. eds.), *Modern Aspects of Emulsion Science*. The Royal Society of Chemistry, Cambridge, 261–291, 1998.

Demorest, L.J., Hamilton, J.G., *Pharmaceutical dosage forms: parenteral medications*, Avis, K.E., Lieberman, H.A., Lachman, L., Eds., Marcel Dekker, v. 1, 251–282, 1992.

Dick, I.P., Scott, R.C., Pig ear skin as an in-vitro model for human skin permeability. *J. Pharm. Pharmacol.*, v. 44, 640–645, 1992.

Díez, O. *Manual de Cosmetología*. Videocinco, ed. 7–117, 2001.

Do Amaral, M., Asua, J.M., Synthesis of large, high–solid–content latexes by miniemulsion polymerization, *J. Polym. Sci. A. Polym. Chem.*, v. 42, 4222–4227, 2004.

Douin V., Cazin, B., Decoster, S. L'oreal S.A., *Fr. Patent*. Jpn. Kokai Tokkyo Koho, JKXXAF JP 2001214081 A2 2001214081. 19pp, Aug 2001.

Douin, V., Cazin, B., Simonnet, J.T., Aubrun, O. L'oreal S.A., *Fr. Eur. Pat. Apply*. EP 1120102 A2 1, 25pp, Aug 2001.

Draize, J.H., Woodward, G., Calvery, H.O., Methods for the study of irritation and toxicity of substances applied topically to the skin and mucous membranes, *J. Pharmacol. Exp. Therap.*, v. 82, 377, 1944.

Dykes, P.J., Pearse, A.D., in *Dermatological and Transdermal Formulations*. Drugs and the Pharmaceutical Sciences. Walters, K.A., v. 119, 511–547, 2002.

## E

Eisenhauer, E.A., Bokkel Huinink, W.W., Swenerton, K.D., European–Canadian randomized trial of paclitaxel in relapsed ovarian cancer: high-dose versus low-dose and long versus short infusion, *J. Clin. Oncol.*, v. 12, 2654–2666, 1994.

El-Aasser, M.S., Lack, C.D., Choi, Y.T., Min, T.I., Vanderhoff, J.W., Fowkes, F.M., Interfacial aspects of miniemulsions and miniemulsion polymers. *Colloids and Surfaces.*, v. 12, 79–97, 1984.

El-Aasser, M.S., Miller, C.M., Preparation of latexes using miniemulsions. (Asua, J.M. (Eds.), *Polymeric Dispersions: Principles and Applications*. Kluwer Academic Publishers, Dordrecht, pp 109–126, 1997.

Elias, P.M., Epidermal barrier function: intercellular lamellar lipid structures, origin, composition and metabolism. *J. Control. Rel.*, v. 15, 199–208, 1991.

Elias, P.M., Epidermal lipids, barrier function, and desquamation, *J. Invest. Dermatol.*, v. 80, 44–49, 1982.

Estebe, J.P., Malledant, Y., Fat embolism after lipid emulsion infusion, *Lancet.*, v. 337, 673, 1991.

Evans D. F., Wennerström H. *The Colloidal Domain*. VCH Publishers Inc. New York, 1994.

## F

Fartasch, M., Bassukas, I.D., Dipegen, T.L., Structural relationship between epidermal lipid lamellae, lamellar bodies and desmosomes in human epidermis: an ultrastructural study, *J. Dermatol.*, v. 128, 1–9, 1993.

Fernandez, C., Marti-Mestres, G., Ramos, J., Maillols, H., LC analysis of benzophenone-3: II application to determination of 'in vitro' and 'in vivo' skin penetration from solvents, coarse and submicron emulsions, *J. Pharm. and Biom. Anal.*, v. 24, 155-165, 2000.

Fevrier, F., Bobin, M.F., Lafforgue, C., Martini, M.C., In vitro cutaneous penetration of proline with microemulsion formulation compared to emulsion, *STP Pharma Sci.*, v. 1, 60-63, 1991.

Finsky, R., Particle sizing by quasi-elastic light scattering, R., *Adv. Colloid Interface Sci.*, v. 52, 79-143, 1994.

Fisher, L.B., Berman, B., Contact allergy to sulfonated castor oil, *Contact Dermatitis*, v. 7(6), 339-340, 1981.

Flynn, G.L., Smith, R.W. Membrana diffusion III: Influence of solvent composition and permeant solubility on membrane transport. *J. Pharm. Sci.*, v. 61, 61-66, 1972.

Forgiarini, A., Esquena, J., González, C., Solans, C., Formation of nano-emulsions by low-energy emulsification methods at constant temperature. *Langmuir*, v. 17, 2076-2083, 2001.

Förster T., Von Rybinski, Wadle, A., Influence of microemulsion phases on the preparation of fine-disperse emulsions, *Adv. Colloid Interface Sci.*, v. 58, 119-149, 1995.

Förster T, Rybinski WV. *Modern Aspects of Emulsion Science*. BP Binks, ed. Cambridge: The Royal Society of Chemistry, 395-426, 1998.

Förster T. *Surfactants in Cosmetics*, (Rieger M, Rhein LD, eds.) New York, Marcel Dekker, 105-125, 1997.

Freel, R.W., Hatch, M., Earnest, D.C., Goldner, A.M., Dihydroxy bile salt-induced alterations in NaCl transport across the rabbit colon, *Am. J. Physiol.*, v. 245, G808-15, 1983.

Friberg, S.E., Buraczewska I., Microemulsions in water- potassium oleate-benzene system, *Progr. Colloid Polym. Sci.*, v. 63, 1-9, 1978.

Friedman, D. Schwarz, J. In *Methods to overcome Biological Barriers in Drug Delivery*, Symposium Proceedings, 75, 1993.

Friedman, D.I., Schwarz, J.S., Weisspapir, M., Submicron emulsion vehicle for enhanced transdermal delivery of steroidal and nonsteroidal anti-inflammatory drugs, *J. Pharm. Sci.*, v. 84, 324-329, 1995.



Frisbee, S.E., McGinity, J.W., Influence of Nonionic Surfactants on the Physical and Chemical Properties of a Biodegradable Pseudolatex, *Eur. J. Pharm. Biopharm.*, v. 40(6), 355–363, 1994.

Fruijtier-Pölloth, C., Safety assessment on polyethylene glycols (PEGs) and their derivatives as used in cosmetic products, *Toxicology*, v. 214 (1-2), 1-38, 2005.

Fuchs, J., Groth, N., Herrling, T., In vitro and in vivo assessment of the irritation potential of different spin straps in human skin, *Toxicology*, v. 151, 55–63, 2000.

## G

Gajewska, M., Sznitowska, M., Janicki, S., Diazepam submicron emulsions containing soya-bean oil and intended for oral or rectal delivery, *Pharmazie*, v. 56(3), 220–222, 2001.

Galindo-Rodriquez, S., Allémann, E., Fessi, H., Doelker, E., Physicochemical parameters associated with nanoparticle formation in the salting-out emulsification-diffusion, and nanoprecipitation methods, *Pharm. Res.*, v. 21 (8), 1428–1438, 2004.

García-Celma, M.J., in Solubilization of drugs in microemulsions, (Solans, C., Kunieda, H., Eds), *Industrial applications of microemulsions*. Marcel Dekker, New York, v. 66, 1–19, 1997.

Gattefossé. Butlletí tècnic del Transcutol P. 2001.

Gelderblom, H., Verweij, J., Nooter, K., Sparreboom, A., Cremophor EL: the drawbacks and advantages of vehicle selection for drug formulation, *Eur. J. of Cancer*, v. 37, 1590–1598, 2001.

Gennes, P.G., Polymers at an interface: a simplified view, *Adv Colloid Interface Sci*, v. 27, 189–209, 1987.

Gennes, P.G., *C R Acad Sci*, Paris, v. 300, 839–843, 1985.

George, L., Tadlock, C.C., Tsolis, P.J., Balian, R.J., *Patent*. PCT Int. Appl. PIXXD2 WO 2002080864 A1 20021017, 15pp, 2002.

Goodman, M., Barry, B.W., Lipid-protein-partitioning (LPP) theory of skin enhancer activity: finite dose technique. *Int. J. Pharm*, v. 57, 29–40, 1989.

Gopal, K., Bhogi B., S., A novel self emulsifying parenteral drug delivery system. *J. Pharm. Sci. Technol.*, v. 53(4), 168–176, 1999.

Gray, G.M., Yardley, H.J., Lipid compositions of cells isolated from pig, human, and rat epidermis. *J. Lipid Res.*, v. 16, 434–440, 1975.

Gröningsson K., Lindgren J.-E., Lundberg E., Sandberg R., Wahlen A., Lidocaine base and hydrochloride. (Florey, K. Eds.), *Analytical profiles of drug substances*. Academic Press, New Jersey, 14, pp 207–235, 1985.

Guy, R.H. and Hadgraft, J., Selection of drug candidates for transdermal drug delivery. (Hadgraft, J. and Guy, R.H. eds.), *Transdermal Drug Delivery*, Dekker, New York, 75–76, 1989.

Guy, R.H., Hadgraft, J., *Mathematical models of percutaneous absorption*. (Bronaugh RL, Maibach HI, eds). *Percutaneous absorption. Mechanisms – Methodology – Drug delivery*. New York: Marcel Dekker, Inc., 13–26, 1989.

## H

Hadgraft, J, Guy, R.H., in Feasibility assessment in topical and transdermal delivery: mathematical models and in vitro studies, *Transdermal Drug Delivery*, v. 123, 1–24, 2003.

Hadgraft, J., Skin, the final frontier, *Int. J. Pharm.*, v. 224, 1–18, 2001.

Handbook of Pharmaceutical Excipients. (Rowe R.C., Sheskey P.J., Weller P.J. Eds.) Pharmaceutical Press and American Pharmaceutical Association, 2005.

Hanson, W.A., *Theoretical concepts. Handbook of Dissolution Testing*, Pharmaceutical Technology Publications, Oregon, p. 4, 1989.

Harrison, J.E., Watkinson, A.C., Green, D.M., Hadgraft, J, Brain, K. The relative effect of azone and transcutol on permeant diffusivity and solubility in human stratum corneum, *Pharm. Res.*, v. 13, 542–546, 1996.

Haskell, R.J., Shifflett, J.R., Elzinga, P.A., *Particle-sizing technologies for submicron emulsions*, in: S. Benita (Ed.), *Submicron emulsions in drug targeting and delivery*, Harwood Academic Publishers, London, 21–98, 1998.

Higuchi, T., Physical chemical analysis of percutaneous absorption process, *J. Soc. Cosmet. Chem.* v. 11, 85–97, 1960.

Hilgers, L.A.Th., Blom, A. G., *Vaccine, in press*, 2005.

Hirai, S., Yashiki, T., Mima, H. Effect of surfactants on the nasal absorption in rats, *Int. J. Pharm.*, v, 9, 165, 1981.

Hiyama, D.T., Griggs, B., Mittman, R.J., Hypersensitivity following lipid emulsion infusion in an adult patient, *J. Parenter. Enteral Nutr.*, v. 13, 318–320, 1989.

Hope, M.J., Kitson, N. Liposomes. A perspective for dermatologists. *Dermatologic clinics*, v. 11, 143–154, 1993.

I

ICI. Butlletí tècnic del Synperonic F68. 2002.

Ilan, E., Amselem, S., Weisspapir, M., Schwarz, J., Yogev, A., Zawoznik, E., Friedman, D., Improved oral delivery of desmopressin via a novel vehicle: mucoadhesive submicron emulsion, *Pharm. Res.*, v. 13, 1083–1087, 1996.

Illel, B., Schaefer, H., Wepierre, J., Doucet, O., Follicles play an important role in percutaneous absorption. *J. Pharm. Sci.*, v. 80, 424–427, 1991.

Irwin, R., NTP Technical report on the toxicity studies of castor oil, *Toxic. Rep. Ser.*, v. 12, 1–B5, 1982.

Israelachvili, J.N., *Intermolecular and Surface Forces*. San Diego: Academic Press, 1991.

Izquierdo, P., Esquena, J., Tadros, Th. F., Dederen, C., García, M.J., Azemar, N., Solans, C., Formation and stability of nano-emulsions prepared using the phase inversion temperature method. *Langmuir*, v.18, 26–30, 2002.

Izquierdo, P., Studies on nano-emulsion formation and stability, *Tesi doctoral*, 2002.

## J

Jakes, J., Testing of the constrained regularization method of inverting Laplace transform on simulated very wide quasielastic light scattering autocorrelation functions, *Czech, J. Phys.*, 38, 1305–1316, 1988.

Jeong Su, K., Ji Su, K., Young Dae, K., Geun, P., Patent. KRXXA7 KR 2003090240 A 20031128, Repub. Korean, 2003.

Johncock, W., Formulation of sunscreens. Favorable and unfavorable interactions, *Cosmet. Technol.* (Milano), 3, 5, 25–31, 2000.

Johnsen, R.M., en *Light Scattering in Biochemistry*, (S.E. Harding, D.B. Sttelle, V.A. Bloomfield, Eds.), The Royal Society of Chemistry, Cambridge, 77–91, 1992.

Jönsson, B., Lindman, B., Holmberg, K., Kronberg, B., in *Surfactant and polymers in aqueous solution*, (John Wiley & Sons ed.), 61–89, 1989.

Jumaa, M., Kleinebudde, P., Müller, B.W., Physicochemical properties and hemolytic effect of different lipid emulsion formulations using a mixture of emulsifiers, *Pharmaceutica Acta Helvetiae*, v. 73 (6), 293–301, 1999.

Jumaa, M., Müller, B.W., Influence of the non-ionic surfactant PEG-660-12-hydroxy stearate on the surface properties of phospholipids monolayers and their effects on lipid emulsion stability, *Colloid Polym. Sci.*, v. 227, 347–353, 1999.

Jumaa, M., Müller, B.W., Development of a novel parenteral formulation for tetrazepam using a lipid emulsion, *Drug Dev. Ind. Pharm.*, 27(10), 1115–1121, 2001.

Jumaa, M., Müller, B.W., Parenteral emulsions stabilized with a mixture of phospholipids and PEG-660-12-hydroxy-stearate: evaluation of accelerated and long-term stability, *Eur. J. Pharm. Biopharm.*, v. 54, 207–212, 2002.

Jumaa, M., Müller, B.W., The effect of oil component and homogenization conditions on the physicochemical properties and stability of parenteral fat emulsions, *Int. J. Pharm.*, v. 163, 81–89, 1998a.

Jumaa, M., Müller, B.W., The stabilization of parenteral fat emulsion using non-ionic ABA copolymer surfactant, *Int. J. Pharm.*, v. 174, 29–37, 1998b.

## K

Kalia, Y.N., Alberti, I., Naik, A., Guy, R.H., *Assessment of topical bioavailability in vivo: the importance of stratum corneum thickness. Skin Pharmacology and Applied Skin Physiology*, v. 14, 82–86, 2001.

Kan, P., Chen, Z.-B., Kung, R.-Y., Lee, C.-J., Chu, I.-M., Study on the formulation of O/W emulsion as carriers for lipophilic drugs, *Colloids and Surfaces B: Biointerfaces*, v. 15, 117–125, 1999.

Kim, J.Y., Ku, Y.S., Enhanced absorption of indomethacin after oral or rectal administration of a self-emulsifying system containing indomethacin to rats, *Int. J. Pharm.*, v. 194, 81–89, 2000.

Kimura, T., Takeda, K., Kageyn, A., Toda, M., Kurpsaki, Y., Nakayama, T., Intestinal absorption of dolichol from emulsions and liposomes in rats. *Chem. Pharm. Bull.*, v. 37, 463–466, 1989.

King, J.W., Determination of the solubility parameter of soybean oil by inverse gas chromatography, *Lebensm.-Wiss. u.-Technol.*, v. 28, 190–195, 1995.

Klang, S., Benita, S., *Submicron emulsions in drug targeting and delivery*, (Simon Benita Eds.), 119–152, 1998.

Ko, K.T., Needham, T.E., Zia, H., Emulsion formulations of testosterone for nasal administration, *J. Macroencapsulation*, v. 15, 197–205, 1998.

Kompaore, F., Dupont, C., Marty, J.P., In vivo evaluation in man by two noninvasive methods of the stratum corneum barrier function after physical and chemical modifications, *Int. J. Cosmet. Sci.*, v. 13, 293–302, 1991.

Konica Minolta, *Precise Color Communication*, 2004.

Koppel, D., Analysis of macromolecular polydispersity in intensity correlation spectroscopy: the method of cumulants, *J. Chem. Phys.*, 57, 4814, 1972.

Kranke, B., Komericki, P., Aberer, W., Olive oil contact sensitizer or irritant?, *Contact Dermatitis*, v. 36 (1), 5–10, 1997.

Krishna, G., Sheth, B.B., *J. Pharm. Sci.*, A novel self emulsifying parenteral drug delivery system, *Technol.*, v. 53(4), 168–176, 1999.

Kronevi, T., Ljungberg, S., Sequelae following intra-arterially injected diazepam formulations, *Acta Pharm. Suecica*, v. 20, 389–396, 1983.

Kunieda, H., Friberg, S., Critical phenomena in a surfactant/water/oil system. Basic study on the correlation between solubilization, microemulsions and ultralow interfacial tensions, *Bull. Chem. Soc. Jpn.*, v. 54 (4), 1010–1014, 1981.

Kunieda, H., Shigeta, K., Suzuki, M., Phase behavior and formation of reverse vesicles in long-polyoxyethylene-chain nonionic surfactant systems. *Langmuir*, v. 15, 3118–3122, 1999.

## L

Landfester, K., Willert, M., Antonietti, Preparation of polymer particles in non-aqueous direct and inverse miniemulsions, *Macromolecules*, v. 33 (7), 2370–2376, 2000.

Larsson, K., Crystal and liquid crystal structures of lipids. (Friberg Eds.), *Food emulsions*. Marcel Dekker, New York, 36–66, 1976.

Lahti, A., Kopola, H., Harila, A., Mullylä, A., Hannuksela, M., Assessment of skin erythema by eye, laser Doppler flowmeter, spectroradiometer, two-channel erythema meter and Minolta chroma meter, *Arch. Dermatol. Res.*, v. 285, 278–282, 1993.

Lee, G.W.J., Tadros, Formation and stability of emulsions produced by dilution of emulsifiable concentrates. Part I. An investigation of the dispersion on dilution of emulsifiable concentrates containing cationic and non-ionic surfactants, Th. F., *Colloids Surf.*, v. 5, 105–115, 1982.

Lehmann, L., Fluhr, J.W., Clemens, M., Gloor, M., Efficacy of different tretinoin formulations under occlusion: TEWL and erythema, *Journal of the European Academy of Dermatology and Venereology*, v.11(2), S274, 1998.

Levy, M.Y., Benita, S., Short-and long-term stability assessment of a new injectable diazepam submicron emulsion, *J. Parenteral Sci. Technol.*, v. 45(2), 101–117, 1991.

Levy, M.Y., Schutze, W., Fuhrer, C., Benita, S., *J. Microencapsulation*, v. 11, 79–92, 1994.

Li, P., Ghosh, A., Wagner, R.F., Krill, S., Joshi, Y.M., Serajuddin, A.T.M., Effect of combined use of nonionic surfactant on formation of oil-in-water microemulsions, *Int. J. Pharm.*, v. 288, 27–34, 2005.

Lidgate, D.M., Fu, R.C., Byars, N.E., Foster, L.C., Fleitman, J.S., Formulation of Vaccine Adjuvant Muramyl dipeptides. 3. Processing Optimization, Characterization, and Bioactivity of an Emulsion Vehicle, *Pharm. Res.*, v. 6, 748–752, 1989.

Lifshitz, I.M., Slyozov, V.V., The kinetics of precipitation from supersaturated solid solutions. *J. Phys. Chem. Solids*, v. 19, 35–50, 1961.

Liu, X., Guan, Y., Ma, Z., Liu, H., Surface modification and characterization of magnetic polymer nanospheres prepared by miniemulsion polymerization, *Langmuir*, v.20, 10278–10282, 2004.

Lotte, C., Rougier, A., Wilson, D.T., Maibach, H.I. In vivo relationship between transepidermal water loss and percutaneous absorption of some organic compounds in man: effect of anatomic site. *Arch Dermatol Res*, v. 279, 351–356, 1987.

Lovell, W.W, Sanders, D.J., Phototoxicity testing in guinea-pigs, *Food Chem. Toxicol.*, v. 30 (2), 155–160, 1992.

Lucks, J.S., Müller, B., *Pharmaceutical emulsion and suspensions*, (Nielloud, F., Marti-Mestres, ed.), Marcel Dekker, 235–239, 2000.

Lundberg B.B., A submicron lipid emulsion coated with amphipathic polyethylene glycol for parenteral administration of paclitaxel (Taxol), *J. Pharm. Pharmacol.*, v. 49, 16–21, 1997.

Lundberg, B., Preparation of drug-carrier emulsions stabilized with phosphatidylcholine-surfactant mixtures, *J. Pharm. Sciences*, v. 83 (1), 72–75, 1994.

Lundberg, B.B., Mortimer, B.-C., Redgrave, T.G., Submicron lipid emulsions containing amphipathic polyethylene glycol for use as drug-carriers with prolonged circulation time, *Int. J. Pharm.*, v. 134, 119–127, 1996.

## M

Macek, T.J., Preparation of parenteral dispersions. Method for solubilization of selected substances. *J. Pharm. Sci.*, v. 52, 694–699, 1963.

Magnusson, B., Identification of contact sensitizers by animal assay, *Contact Dermatitis*, v. 6, 46, 1980.

Magnusson, B., Kligman, A.M., *Allergic Contact Dermatitis in the Guinea-pig. Identification of Contact Allergens*, Charles C., Thomas, 1970.

Malcolmson, C., Lawrence, M.J., A comparison of the incorporation of model steroids into non-ionic micellar and microemulsion systems, *J. Pharm. Pharmacol.*, v. 45, 141–143, 1993.

Marszall, L., *Nonionic Surfactants*, (Shick, M. J. ed.), Surfactant sciences series, Marcel Dekker, New York, v. 23, 493–547, 1987.

Martindale, Guía Completa de Consulta Farmacoterapéutica, dirigida por S.C. Sweetman, Pharma Editores, 2003.

Mbela, T.K.M., Deharo, E., Haemers, A., Ludwig, A., Submicron oil-in-water emulsion formulations for mefloquine and halofantrine: effect of electric-charge inducers on antimalarial activity in mice, *J. Pharm. Pharmacol.*, v. 50, 1221–1225, 1998.

McNiff, B.L., Clinical use of 10% soybean oil emulsion, *Am. J. Hosp. Pharm.*, v. 34, 1080–1086, 1977.

Mead, R., Curnow, R. N., Hasted, A.M., in *Statistical Methods in agriculture and experimental biology*, Chapman and Hall/CRC, 3rd Edition, 245–251, 2003.

Medina, J., Salvadó, A., del Pozo, A., Use of ultrasound to prepare lipid emulsions of lorazepam for intravenous injection, *International Journal of Pharmaceutics*, v. 216, 1–8, 2001.

MERCK. Butlletí tècnic del Tween 80. 2000.

Mezei, M., in *Drug Permeation Enhancement*, (Hsieh D.S., eds.), Drugs and the Pharmaceutical Sciences, v. 62, 171–199, 1994.

Miller, K.A., Balakrishnan, G., Eichbauer, G., Betley, K., *1% Lidocaine injection, EMLA cream, or 'Numby Stuff' for topical analgesia associated with peripheral intravenous cannulation*, AANA J. 69, 185–187, 2001.

Miralles, M.J., McGinity, J.W., Martin, A., A combined water-soluble carriers for coprecipitates of tolbutamide, *J. Pharm. Sci.*, v. 71, 302–304, 1982.

Morales D., Gutierrez J.M., Garcia-Celma M.J., Solans C., A study of the relation between bicontinuous microemulsions and oil/water nano-emulsion formation. *Langmuir*, v. 19, 7196–7200, 2003.

Moser, K., Kriwet, K., Naik, A., Kalia, Y.N., Guy, R.H., Passive skin penetration enhancement and its quantification in vitro. *Eur. J. Pharm. Biopharm.* v. 52, 103–112, 2001.

## N

Nakajima H., Tomomasa S., Kochi M. *J. Soc. Cosmet. Chem. Jpn.* v. 34, 335, 1983.



Nakajima, H., Microemulsions in cosmetics, (Solans, C., Kunieda, H. Eds.), *Industrial applications of microemulsions*. Marcel Dekker, New York, v. 66, 175–197, 1997.

Nakajima, H.; Tomomasa, S.; Okabe, M., Preparation of nano-emulsions. Recueil des resumes. Premier congres mondial de l'emulsion, Paris, v. 1, 63, 1993.

Nankevis, R., Davis, S.S., Day, N.H., Studies on the intravenous pharmacokinetics of three retinoids in the rat, *Int. J. Pharm.*, v. 101, 249–256, 1994.

Nasirideen, S., Kas, H.S., Öner, F., Alpart, R., Hincal, A.A., Naproxen incorporated lipid emulsions. I. Formulation and stability studies, *Journal of Clinical Pharmacy and Therapeutics*, Vol. 23, 57–65, 1998.

Nicolaos, G., Crauste-Manciet, S., Farinotti, R., Brossard, D., Improvement of cefpodoxime proxetil oral absorption in rats by an oil-in-water submicron emulsion, *Int. J. Pharm.*, v. 263, 165–171, 2003.

Nilsson, G.E., Measurement of water exchange through skin, *Med. Biol. Eng. Comput.*, v. 15, 209–218, 1977.

## O

Overbeek, J.Th.G., The first rideal lecture. Microemulsions, a field at the border between lyophobic and lyophilic colloids, *Faraday Disc. Chem. Soc.*, v. 65, 7, 1978.

## P

Pan, G., Shower, S., Lu, D.R., In vitro gene transfection in human glioma cells using a novel and less cytotoxic artificial lipoprotein delivery system, *Pharm. Res.*, v. 20, 738–744, 2003.

Park, J.W., Park, E.-S., Chi, S.-C., Kil, H. Y., Lee, K.-H, The effect of lidocaine on the globule size distribution of propofol emulsions. *Anesthesia & Analgesia*, v. 97(3), 769–771, 2003.

Pendlington, R.U., Minter, H.J., Sanders, D.J., Howes, D. *Comparative in vitro percutaneous absorption of 2-hydroxyacids*, *Perspect. Percutan. Penetrat.*, 6A, 107, 1998.

Pich, A., Datta, S., Musyanovych Adler, H.-J.P., Engelbrecht, L, Polymeric particles prepared with fluorinated surfmer, *Polymer*, v. 46, 1323–1330, 2005.

Pinnagoda, J., Tupker, R.A. *Measurement of the TEWL*. In: Serup J, Jemec GBE, eds. *Non-invasive Methods and the Skin*. Boca Raton, FL: CRC Press, v. 173, 1995.

Pinnagoda, J., Tupker, R.A., Coenraads, P.J., Nater, J.P., Prediction of susceptibility to on irritant response by transepidermal water loss, *Contact Dermatitis*, v. 20, 241–246, 1989.  
Porrás, M., Solans, C., González, C., Martínez, A., Guinart, A., Gutiérrez, J.M., Studies of formation of W/O nano-emulsions, *Colloids Surf., A. Physicochem. Eng., Asp.*, v.249, 115–118, 2004.

Potts, R.O., Francoeur, M.L., *Lipid biophysics of water loss through the skin*. Proc. Natl. Acad. Sci., v. 87, 3871–3873, 1990.

Potts, R.O., Guy, R.H. Predicting skin permeability, *Pharm. Res.*, v. 9, 663–669, 1992.

Priborsky, J., Muhlbachova, E. Evaluation of in vitro percutaneous absorption across human skin and animal models. *J. Pharm. Pharmacol.*, v. 42, 468–472, 1990.

Provencher, S.W., A constrained regularization method for inverting data represented by linear algebraic equations, *Comput. Phys. Commun*, v. 27, 213–217, 1982a.

Provencher, S.W., A general purpose constrained regularization method for inverting data represented by linear algebraic equations, *Comput. Phys. Commun*, v. 27, 229–241, 1982b.

## R

Real Farmacopea Española, Tercera edición, 2005.

Roberts, M.E., Mueller, K.R., Comparisons of in vitro nitroglycerin (TNG) flux across Yucatan pig, hairless mouse, and human skins. *Pharm. Res.*, v. 7, 673–676, 1990.

Roberts, M.S., Walters, K.A., The relationship between structure and barrier function of skin. (Roberts M.S., Walters, K.A., eds.) *Dermal absorption and toxicity assessment*. New York: Marcel Dekker, Inc., 1–42, 1998.

Rosano, H.L., Lan, T., Weiss A., Whittam, J.H., Gerbacia, W.E.F., Unstable microemulsions. *J. Phys. Chem.*, v. 85, 468–473, 1981.

Rowinsky, E.K., Eisenhauer, E.A., Chaudhry, V., Arbuck, S.G., Clinical toxicities encountered with paclitaxel (Taxol), *Semin. Oncol.*, v. 20, 1–15, 1993.

## S

Sadurní, N., Solans, C., Azemar, N., García-Celma, M.J., Studies on the formation of O/W nano-emulsions, by low-energy emulsification methods, suitable for pharmaceutical applications. *Eur. J. Pharm. Sc.*, v. 26, 438–445, 2005.

Sagitani, H. En, *Organized solutions*, Eds. Friberg S.E., Lindman B. New York, Marcel Dekker, 259–271, 1992.

Sagitani, H., Nabeta, K., Nagai, M., *Yukagaku*, v. 40 (11), 988–94, 1991.

Santos-Magalhaes, N.S., Pontes, A., Pereira, V.M.W., Caetano, M.N.P., Colloidal carriers for benzathine penicillin G: nanoemulsions and nanocapsules, *Int. J. Pharm.*, v. 208, 71–80, 2000.

Schaefer, H., Redelmeier, T.E., Benech-Kieffer, F., in *Novel Cosmetic Delivery Systems*. (Magdassi, S., Touitou, E. eds.), Cosmetic Science and Technology Series, v. 19, 9–49, 1999.

Scheuplein, R.J., Blank, I.H., Permeability of the skin, *Physiol. Rev.*, v. 51, 702–747, 1971.

Schork, F.J., Poehlein, G.W., Wang, S., Reimers, J., Rodrigues, J., Samer, C., Miniemulsion polymerization, *Colloids Surf. A*, 153, 39–45, 1999.

Schreier, H., Bouwstra, J. Liposomes and niosomes as topical drug carriers: dermal and transdermal drug delivery. *J. Control. Rel.*, v. 30, 1–15, 1994.

Schwarz, J.S., Weisspapier, M.R., Friedman, D.I., Enhanced transdermal delivery of diazepam by submicron emulsion (SME) creams, *Pharmac. Res.*, v. 12, 687–692, 1995.

Scott, R.C. *Biological monitoring for pesticide exposure: measurement, estimation and risk reduction*. In: Wang RGM, Franklin, CA, Honeycutt RC, Reinert JC, eds. ACS Symposium Series 382. Washington, DC: American Chemical Society, 158–168, 1989.

Seijo, B., Fattal, E., Treupel, L. R., Couvreur, P., Design of nanoparticles of less than 50nm diameter: preparation, characterization. and loading, *Int. J. Pharm*, v. 62, 1–7, 1990.

Seung Gi, H., Sang Jin, K., Cheong Taek, K., Ho Jeong, K., Jae Chan, Y., Yeong Chang, Y. *Patent*. KRXXA7 KR 2003037828 A 20030516. Repub. Korean, 2003.

Shinoda, K., Kunieda, H. En *Encyclopedia of Emulsion Technology*, (Becher, P., ed.), Marcel Dekker, New York, v. 1, 337–367, 1983.

Shinoda, K., Saito, H. *J. Colloid Interface Sci.*, v. 30, 258, 1969.

Shinoda, K., Saito, H. *J. Colloid Interface Sci.*, v. 26, 70, 1968.

Simmonet, J.-T., Legret, S., Sonnevile, O. L'Oreal, *Fr. Eur. Pat. Appl.* EPXXDW EP 1013339 A1 20000628. 9pp, 2000b.

Simmonet, J.-T., Sonnevile, O., Legret, S. L'Oreal, *Fr. Eur. Pat. Appl.* EPXXDW EP 1010413 A1 20000621. 12pp, 2000a.

Simmonet, J.-T., Sonnevile, O., Legret, S. L'Oreal, *Fr. Eur. Pat. Appl.* EP 1018363 A1. 2000c.  
Singh, J., Maibach, H.I., in *Dermatological and Transdermal Formulations*, v. 119, 529–547, 2002.

Singh, M.A., Ghosh, S.S., Shannon R. F., A direct method of beam–height correction in small-angle X-ray scattering, *J. Appl. Cryst.*, v. 26, 787–794, 1993.

Sintov, A.C., Shapiro, L., New microemulsion vehicle facilitates percutaneous penetration in vitro and cutaneous drug bioavailability in vivo, *J. Controlled Release*, v. 95, 173–183, 2004.

Smidt, P.C., Campanero, M.A, Troconiz, I.F., Intestinal absorption of penclomedine from lipid vehicles in the conscious rat: contribution of emulsification versus digestibility, *Int. J. Pharm.*, v. 270 (1–2), 109–118, 2004.

Solans, C., Esquena, J., Forgiarini, A., Uson, N., Morales, D., Izquierdo, P., Azemar, N., García-Celma, M.J., 2003. Nano-emulsions: Formation, properties, and applications. (Mittal K.L., Shah, Dinesh O. Eds.), *Adsorption and Aggregation of Surfactants in Solution*. Marcel Dekker, New York, v. 109, 525–554, 2003.

Solans, C., Izquierdo, P., Nolla, J., Azemar, N., Garcia-Celma, M.J., Nano-emulsions, *Current Opinion in Colloid and Interface Science*, v. 10, 102–110, 2005.

Solans, C., Pons, R., Kunieda, H., Overview of basic aspects of microemulsions. (Solans, C., Kunieda, H., Eds), *Industrial applications of microemulsions*. Marcel Dekker, New York, v. 66, 1–19, 1997.

Sonneville, O., Simmonet, J.T., Legreg, S. *Eur. Pat. Appl.* EP 1010414 A1, 2000.

Sonneville–Aubrun, O., Simonnet, J.T., L’Alloret, F, Nano–emulsions a new vehicle for skincare products, *Adv. Colloid Interface Sci.*, v. 108, 145–149, 2004.

Souto, E.B., Wissing, S.A., Barbosa, C.M., Müller, R.H., Evaluation of the physical stability of SLN and NLC before and after incorporation into hydrogel formulations, *Eur. J. Pharm. Biopharm.*, v. 58, 83–90, 2004.

Surber, C., Davis, A.F., in *Dermatological and Transdermal Formulations. Drugs and the Pharmaceutical Sciences.* v. 119, 401–498, 2002.

Sznitowska, M.; Gajewska, M., Janicki, S.; Radwanska, A., Lukowski, G., Bioavailability of diazepam from aqueous–organic solution, submicron emulsion and solid lipid nanoparticles after rectal administration in rabbits, *Eur. J. Pharm. Biopharm.*, v. 52, 159–163, 2001.

Sznitowska, M.; Janicki, S.; Dabrowska, E.; Zurowska–Pryczkowska K., Submicron emulsions as drug carriers. Studies on destabilization potential of various drugs. *Eur. J. Pharm. Sci.*, v. 12, 175–179, 2001.

## T

Tabibi, E. En *Specialized Drug Delivery Systems*, (Tyle P., ed.), Drug and the Pharmaceutical Sciences Series, Marcel Dekker, New York, v. 41, 317–331, 1990.

Tabosa do Egito, E.S., Fessi, H., Appel, M., Puisieux, F., Bolard, J., Devissaguet, J.P., New techniques for preparing submicronic emulsions: application to amphotericin B, *S.T.P. Pharma Sciences*, v.4 (2), 155–162, 1994.

Tadros, Th. F., Vincent, B., en: *Encyclopedia of Emulsion Technology* (Ed. Binks P.), Marcel Dekker, v. 1, 1–56, 1983.

Taha, E.I., Al–Saidan, S., Samy, A.M., Khan, M.A., Preparation and in vitro characterization of self–nanoemulsified drug delivery system (SNEDDS) of all–trans–retinol acetate, *Int. J. Pharm.*, v. 285. 109–119, 2004.

Tamilvanan, S., Oil–in–water lipid emulsions: implications for parenteral and ocular delivering systems, *Prog. Lipid Res.* v. 43, 489–533, 2004.

Tanaka, S., Takashima, Y., Murayama, H., Tsuchiya, S. *Int. J. Pharm.*, v. 27, 29–38, 1985.

Toguchi, H., Pgawa, Y., Shimamoto, T., Effects of the physicochemical properties of the emulsion formulation on the bioavailability of ethyl 2-chloro-3-[4-(2-methyl-2-phenylpropyloxy)phenyl]propionate in rats, *Chem. Pharm. Bull.*, v. 38, 2797–2800, 1990.

Treffel, P., Gabard, B., Bieli, E., *Prediction of Percutaneous Penetration*. (Brain, KR, James, VJ, Walters, KA, eds.) V.3b, 520–529, 1993.

Trotta, M., Gasco, M.R., Pattarino, F., *Proceedings of the 10th Pharmaceutical Technology Conference*, 388–404, 1991.

Trotta, M., Pattarino, F., Ignoni, T., Stability of drug-carrier emulsions containing phosphatidylcholine mixtures, *Eur. J. Pharm. Biopharm.*, v. 53, 203–208, 2002.

## U

Ugelstad, J.; El-Aasser, M.S.; Vanderhoff, J.W., Emulsion polymerization: initiation of polymerization in monomer droplets. *J. Polym. Sci., Polym.*, v. 11, 503–513, 1973.

Uson, N., Garcia M.J., Solans C., Formation of water-in-oil (W/O) nano-emulsions in a water/mixed non-ionic surfactant/oil systems prepared by a low-energy emulsification method. *Colloids and surfaces A*, 250, 415–421, 2004.

Uson, N. Tesi doctoral: *Estudi de la formació de nano-emulsions de tipus aigua en oli*, Universitat de Barcelona, 2003.

## V

Van Joost, T., Smitt, J.H., Van Ketel, W.G., Sensitization to olive oil (*olea europeae*), *Contact Dermatitis*, v. 7(6), 309–310, 1981.

Verma, D.D., Fahr, A., Synergistic penetration enhancement effect of ethanol and phospholipids on the topical delivery of cyclosporina, *Journal of Controlled Release*, v. 97 (1), 55–66, 2004.

Verwey, E.S.W., Overweek, J.Th.G., *Theory of the Stability of Lyophobic Colloids*, Elsevier, Amsterdam, 1948.

Vohr, H.W., Blumerl, J., Blotz, A., Homey, B., Ahr, H.J., An intra-laboratory validation of the integrated model for the differentiation of skin reactions (IMDS): discrimination between

(photo) allergic and (photo) irritant skin reactions in mice, *Arch Toxicol.*, v. 73, 501–509, 2000.

## W

Wadle, A., Förster, T., von Rybinski, W., Influence of the microemulsion phase-structure on the phase inversion temperature emulsification of polar oils, *Colloid Surfaces A: Physicochem. Eng. Aspects*, v. 76, 51–57, 1993.

Wadle, A., Tesmann, H., Leonard, M., Förster, T. In *Surfactants in Cosmetics*, Rieger, M. M., Rhein, L. D., eds.) Surfactant Sciences Series, v. 68, 207–224, 1997.

Wagner, C., Theorie der Alterung von Niederschlägen durch Umlosen (Ostwald-Reifung). *Zeitschrift fuer Elektrochemie*, v. 65, 581, 1961.

Walstra, P., Formation of emulsions. (Becher, P. Eds), *Encyclopedia of Emulsion Technology*. Marcel Dekker, New York, v. 1, 57–127, 1983.

Walstra, P. In *Encyclopedia of Emulsion Technology*, (Becher, P., ed.), Marcel Dekker, New York, v. 4, 1–62, 1996.

Walstra, P., Smulders PEA. In *Modern Aspects of Emulsion Science*. (BP Binks, ed.), Cambridge: The Royal Society of Chemistry, 56–59, 1998.

Walters, K., Brain, K.R., in *Dermatological and Transdermal Formulations*, Walters, K.A., eds., *Drugs and the Pharmaceutical Sciences*, v. 119, 319–400, 2002.

Watkinson, A.C., Hadgraft, J., Walters, K.A., Brain, K.R., Measurement of diffusional parameters in membranes using ATR-FTIR spectroscopy. *Int. J. Cosmet. Sci.*, v. 16, 199–210, 1994.

Weatherall I.L., Coombs, B.D., Skin color measurements in terms of CIELAB colour space values, *J. Invest. Dermatol.*, v. 99, 468–473, 1992.

Webb, N.E., Method for solubilization of selected drug substances, *Bull Parenter. Drug Assoc.*, v. 30, 180–186, 1976.

Weingarten, C., Magalhaes, N.S.S., Baszkin, A., Benita, S., Seiller, M., Interactions of a nonionic ABA copolymer surfactant with phospholipid: monolayers-possible relevance to emulsion stabilization, *Int. J. Pharm.*, v. 75, 171–179, 1991.

Weiss, R.B., Donehower, R.C., Wiernik, P.H., Hypersensitivity reactions from taxol, *J. Clin. Oncol.*, v. 8, 1263–1268, 1990.

Wertz, P.W., Downing, D.T., *Stratum corneum: biological and biochemical considerations*, in: J. Hadgraft, R.H. Guy (Eds.), *Transdermal Drug Delivery: Developmental Issues and Research Initiatives*, Marcel Dekker, New York, 1–22, 1989.

Wheeler, D.A., Moyler, D.A., Thirkettle, J.T., Instrumental colour assessment: some practical experience, *J. Soc. Cosmet. Chem.*, v. 27, 15–43, 1976.

Wilhelm, K.-P., Surber, C., Maibach, H.I., Quantification of sodium lauryl sulphate dermatitis in man: comparison of four techniques: skin color reflectances, transepidermal water loss, laser Doppler flow measurement and visual scores, *Arch. Dermatol. Res.*, v. 281, 293–295, 1989.

Wilson, D.R., Maibach, H.I., *TEWL: a review*, (Leveque J.L., ed). *Cutaneous investigation in Health and Disease: Noninvasive Methods and Instrumentation*. New York: Marcel Dekker, 113, 1989.

Winsor, P.A., Hydrotropy, solubilisation and related emulsification processes. *Trans. Faraday Soc.*, v. 44, 376–398, 1948.

Wretling, A.J., The pharmacological basis for the use of fat emulsions in intravenous nutrition, *Acta Chir. Scand.*, v. 325, 31–42, 1964.

Wu, H., Ramachandran, C., Bielinska, A.U., Kingzett, K., Sun, R., Weiner, N.D., Roessler, B.J., Topical transfection using plasmid DNA in a water-in-oil nano-emulsion, *Int. J. Pharm.*, v. 221, 23–34, 2001.

Wu, H., Ramachandran, C., Weiner, N.D., Roessler, B.J., Topical transport of hydrophilic compounds using water-in-oil nano-emulsions, *Int. J. Pharm.*, v. 220, 63–75, 2001.

## Y

Yalin, M., Öner, F., Öner, L., Hincal, A.A., Preparation and properties of a stable intravenous lorazepam emulsion, *Journal of Clinical and Therapeutics*, v. 22, 39–44, 1997.

Youenang Piemi, M.P., Korner, D., Benita, S., Marty, J.P. Positively and negatively charged submicron emulsions for enhanced topical delivery of antifungal drugs. *J. Controlled Release*, v. 58, 177–187, 1999.



## Z

Zhang, S.-W., Zhou, S.-X., Weng, Y.M., Weng, Wu, L.-M., Synthesis of SiO<sub>2</sub>/polystyrene nanocomposite particles via miniemulsion polymerization, *Langmuir*, v. 21, 2124–2128, 2004.