

Departament de Geografia Física i Anàlisi Geogràfica Regional  
Facultat de Geografia i Història  
Universitat de Barcelona



**Tesi doctoral**

***Reconstrucció paleoambiental holocena de Sierra Nevada  
a partir de registres sedimentaris***

Memòria presentada per en  
**Marc Oliva i Franganillo**  
Per optar al títol de Doctor en Geografia

Programa de Doctorat  
*Gestió ambiental, Paisatge i Geografia*  
Bienni 2004-2006

Els directors de tesi

Dr. Antonio Gómez Ortiz

Dr. Lothar Schulte

## CAPÍTOL 9

## **BIBLIOGRAFIA**



## **9. Bibliografia**

- Abrantes, F.; Baas, J.; Hafliðason, H.; Rasmussen, T.; Klitgaard, D.; Lončarić, N. & Gaspar, L. (1998).** Sediment fluxes along the northeastern European Margin: inferring hydrological changes between 20 and 8 kyr. *Marine Geology*, 152: 7-23.
- Abreu L.; Shackleton N. J.; Schonfeld J.; Hall M. & Chapman M. (2003)** Millennial-scale oceanic climate variability of the Western Iberian margin during the last two glacial periods. *Marine Geology*, 196: 1–20.
- AG Bodenkundliche Kartieranleitung, (2005).** 5. verbesserte und erweiterte Auflage. Bundesanstalt für Geowissenschaften und Rohstoffe, AG Boden, Hannover.
- Akerman J.H. (1993).** Solifluction and creep rates 1972-1991, Kapp Linne, West Spitsbergen. In: Frenzel, B., Matthews, J.A., Gläser, B. (eds). *Solifluction and Climatic Variation in the Holocene*. Paläoklimaforschung, Vol. 11: 225-249.
- Al-Alwani, G. (1997).** *Hidrología e hidrogeología de la cuenca del río Guadalfeo*. Tesi doctoral. Universitat de Granada. 315 pp.
- Alcoforado, M.J., Nunes, M.F., Garcia, J.C. & Taborda, J.P. (2000).** Temperature and precipitation reconstruction in Southern Portugal during the late Maunder Minimum (AD 1675-1715). *The Holocene*, 10 (3): 333-340.
- Alexander, C.S. & Price, L.W. (1980).** Radiocarbon dating of the rate of movement of two solifluction lobes in the Ruby Range, Yukon Territory. *Quaternary Research*, 13: 365-379.
- Allen, J.R.M.; Watts, W.A.; McGee, E. & Huntley, B. (2002).** Holocene environmental variability-the record from Lago Grande di Monticchio, Italy. *Quaternary International*, 88: 69-80.
- Alley, R.B. & Clark P.U. (1999).** The deglaciation of the Northern Hemisphere: a global perspective. *Annual Review Earth Planetary Science*, 27: 149-182.
- Alley, R.B. (2000).** The Younger Dryas cold interval as viewed from central Greenland. *Quaternary Science Reviews*, 19: 213-226.
- Álvarez, M.C.; Flores, J.A.; Sierro, F.J.; Diz, P.; France, G.; Pelejero, C. & Grimalt, J. (2005).** Millennial surface water dynamics in the Ría de Vigo during the last 3000 years as revealed by coccoliths and molecular biomarkers. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 218: 1–13.
- Alverson, K.D.; Bradley, R.S. & Pedersen, T.F. (2002).** *Paleoclimate, global change and the future (PAGES)*. Springer, Berlin, 220 pp.
- Andersson, J.G. (1906).** Solifluction, a component of subaerial denudation. *Journal of Geology*, 14: 91-112.
- Andren, A.W. & Strand, J.W. (1981).** Atmospheric deposition of particulate organic carbon and polyaromatic hydrocarbons to Lake Michigan. In: Eisenreich, S.J., Editor, , 1981. *Am. Chem. Soc., Div. Environ. Chem., Prepr.*. Vol. 19(2): 216-218.
- Andrews, J. T.; H. Erlenkeuser, K.; Tedesco, A. E.; Aksu, & Jull, T. (1994).** Late Quaternary (stage 2 and 3) meltwater and Heinrich events, northwest Labrador Sea, *Quaternary Research*, 41:26.

- Arduino, E., E. Barberis, F. Carraro, and M. G. Forno, 1984.** Estimating relative ages from iron-oxide/total-iron ratios of soils in the western Po Valley, Italy. *Geoderma*, 33: 39-52
- Austin, W.E.N. & Kroon, D. (1996).** The Lateglacial palaeoceanographic evolution of the Hebridean Continental Shelf, N.W. Scotland. *Geological Society Special Publication*, 111: 187-214.
- Ballantyne C.K. & Harris, C. (1994).** *The periglaciation of Great Britain*. University of Cambridge, Cambridge.
- Barriendos, M. (1994).** *El clima histórico de Catalunya. Aproximación a sus características generales (Siglos XVII-XIX)*. Universitat de Barcelona.
- Barriendos, M. (1997).** Climatic variations in the Iberian Peninsula during the late Maunder Minimum (AD 1675-1715): an analysis of data from rogation ceremonies. *The Holocene*, 7 (1): 105-111.
- Barriendos, M. & Martín-Vide, J. (1998).** Secular climatic oscillations as indicated by catastrophic floods in the Spanish Mediterranean coastal area (14th-19th Centuries). *Climatic Change*, 38: 473-491.
- Barriendos, M. & Llasat, M.C. (2003).** The case of the 'Maldá' anomaly in the Western Mediterranean basin (AD 1760–1800): an example of a strong climatic variability. *Climatic Change*, 61: 191-216.
- Bäumler, R. & Zech, W. (1994).** Soils of the high mountain region of Eastern Nepal: classification, distribution and soil forming processes. *Catena*, 22: 85–103.
- Bech, J.; Rustullet, J.; Garrigo, J.; Tobias, F.J. & Martinez, R. (1997).** The iron content of some red Mediterranean soils from northeast Spain and its pedogenic significance. *Catena* 28: 211–229.
- Beer, J.; Mende, W. & Stellmacher, R. (2000).** The role of the sun in climate forcing. *Quaternary Science Reviews*, 19: 403-415.
- Bell, M. & Walker, M.J. (2005).** *Late Quaternary Environmental Change. Physical and Human perspectives*. PearsonPrentice Hall, Glasgow. 273 pp.
- Benedict, J. B. (1966).** Radiocarbon Dates from a Stone-Banked Terrace in the Colorado Rocky Mountains, USA. *Geografiska Annaler. Series A, Physical Geography*, 48 (1): 24-31.
- Benedict, J.B. (1970).** Downslope soil movement in a Colorado alpine region: rates, processes and climatic significance. *Arctic and Alpine Research*, 2: 165-226.
- Benedict, J.B. (1976).** Frost creep and gelifluction features: a review. *Quaternary Research*, 6: 55-76.
- Bennet, K.D. & Willis K.J. (2001).** Pollen. In: Last, W.M. & Smol, J.P (eds). *Tracking Environmental Changes Using Lake Sediments*. Kluwer Academic Publishers. Països Baixos. Vol. 3: 5-32.
- Berglund, B. (1986).** *Handbook of Holocene Palaeoecology and Palaeohydrology*. Wiley, Chichester, 869 pp.
- Birkeland, P.W. (1999).** *Soils and geomorphology*. Oxford University Press, New York.
- Birks, H. H. & Lotter, A. F. (2000).** *NORPEC sampling protocol for AMS radiocarbon dating*.
- Björk, S.; Walker, M.J.C.; Cwynar L.C.; Knudsen, K.L.; Lowe, J.J. & Wohlfarth, B. (1998).** An event stratigraphy for the Last Termination in the North Atlantic region based on the Greenland ice core record: a proposal by the INTIMATE group. *Journal of Quaternary Science*, 13: 283-292.
- Blake, W. J. (1988).** Geological survey of Canada radiocarbon dates XXVII. *Geological Survey of Canada*, 87(7): 1–100.

- Blott, S. & Pye, K. (2006).** Particle size distribution analysis of sand-sized particles by laser diffraction: an experimental investigation of instrument sensitivity and the effects of particle shape. *Sedimentology*, 53 (3): 671-685.
- Blunier, T. & Brook E. (2001).** Timing of millennial-scale climate change in Antarctica and Greenland during the Last Glacial period. *Science* 291: 109.
- Boelhouwers, J.; Holness, S. & Summer, P. (2003).** The maritime Subantarctic: a distinct periglacial environment. *Geomorphology*, 52: 39-55.
- Boissier, C.E. (1839).** Voyage botanique dans le midi de l'Espagne pendant l'année 1837. Gide et Cie. París. A BOISSIER, C.E.: *Viaje botánico al sur de España durante el año 1837*. Colección Sierra Nevada y la Alpujarra. Fundación Caja de Granada- Universidad de Málaga. Granada, 1995.
- Bolós, M. de (1978).** La vegetación en España. A: Terán, M.; Solé Sabarís, L.. & Vilà Valentí, J. *Geografía general de España*. Editorial Ariel, vol. 1, pp. 209-234.
- Bond, G.; Broecker, W.; Johnsen, S.; Mcmanus, J.; Labeyrie, L.; & Bonani, G. (1993).** Correlations between climate records from North Atlantic sediments and Greenland ice. *Nature*, 365: 143-147.
- Bond, G. G.; Kromer, B.; Beer, J.; Muscheler, R.; Evans, M.; Showers, W.; Hoffmann, S.; Lotti-Bond, R.; Hajdas, I. & Bonani, G. (2001).** Persistent solar influence on North Atlantic climate during the Holocene, *Science*, 294: 2130-2136.
- Bradley, R. (2003).** Climate forcing during the Holocene. *PAGES*, 11, 2 i 3.
- Bradley, R.S. & Jones, P.D. (eds) (1992).** *Climate since 1500 A.D.* Routledge. London. 679 pp.
- Bradley, R.S.; Hughes, M. & Diaz, H. (2003).** Climate in Medieval Time. *Science*, 302: 404-405.
- Brewer, S., Cheddadi, R., De Beaulieu, J.L. & Reille, M. (2002).** The spread of deciduous Quercus throughout Europe since the last glacial period. *Forest Ecology and Management*, 156: 27-48.
- Broecker, W.S. (2000).** Was a change in thermohaline circulation responsible for the Little Ice Age? *Proc. National Academic Sciences*, 97 (4): 1339-1342.
- Brown, R. J. E. (1973).** Influence of climate and terrain factors on ground temperatures at three locations in the permafrost region of Canada. A: Permafrost; North American Contribution, Second International Permafrost Conference, Yakutsk, URSS. Washington. National Academy of Science, nº 2115, pp. 27-34.
- Burjachs, F.; Giral, S.; Riera, S.; Roca, R. & Julià, R. (1996).** Evolución paleoclimática durante el último ciclo glacial en la vertiente mediterránea de la Península Ibérica. *Notes Geografía Física* 25: 21-39.
- Burjachs, F.; Expósito, I. & Yll, I.E.. (2007).** Tendencias, crisis y desfases de la vegetación y el clima en el litoral mediterráneo de la Península Ibérica durante el Holoceno. I Congreso Nacional sobre Cambio Global. Universidad Carlos III de Madrid. Poster. Getafe, 25-27 abril 2007.
- Cabrera, LL. & Anadón, P. (2003).** Desarrollo de los proyectos IGCP 219 y 324 (GLOPALS-GGLAB) en España: Kerry Kelts como elemento dinamizador. A: Valero-Garcés, B.L. (ed.). *Limnogeología en España: un tributo a Kerry Kelts*. Consejo Superior de Investigaciones Científicas. pp. 37-57.
- Cacho, I.; Grimalt, J.; Canals, M.; Sierro, F.; Flores, J. & Shackleton, N. (1999).** Dansgaard-Oeschger and Heinrich event imprints in Alboran Sea paleotemperatures, *Paleoceanography*, 14(6): 698-705.
- Cacho, I.; Grimalt, J.; Sierro, F.J.; Flores, J.A.; Shackleton, N. & Canals, M. (2000).** Evidence for enhanced Mediterranean thermohaline circulation during rapid climatic coolings. *Earth and Planetary Science Letters*, 183: 417-429.

- Cacho, I.; Grimalt, J.O.; Canals, M.; Scaffi, L.; Shackleton, N.J.; Schönfeld, J. & Zhan, R. (2001).** Variability of the western Mediterranean Sea surface temperature during the last 25,000 years and its connection with the Northern Hemisphere climatic changes. *Paleoceanography* 16: 40-52.
- Carrión, J.S. (2003).** Sobresaltos en el bosque mediterráneo: incidencia de las perturbaciones observables en una escala paleoecológica. *Ecosistemas*, 3.
- Carrión, J.S.; Munuera, M.; Dupré, M. & Andrade, A. (2001).** Abrupt vegetation changes in the Segura Mountains of southern Spain throughout the Holocene. *Journal of Ecology*, 89: 783-797.
- Carrión, J.; Fuentes, N.; González-Sampériz, P.; Sánchez Quirante, L.; Finlaysson, C.; Fernández, S. & Andrade, A. (2007).** Holocene environmental change in a montane region of southern Europe with a long history of human settlement. *Quaternary Science Reviews*, 26:1455-1475.
- Castillo, A.; del Valle, M.; Rubio-Campos, J.C. & Fernández-Rubio, R. (1996).** Síntesis hidrológica del macizo de Sierra Nevada (Granada y Almería). *1ª Conferencia Internacional Sierra Nevada. 20-26 marzo de 1996*: 389-413.
- Chappellaz, J.; Blunier, T.; Kints, S.; Dallenbach, A.; Barnola, J.M.; Schwander, J.; Raynaud, D. & Stauffer, B. (1997).** Changes in the atmospheric CH<sub>4</sub> gradient between Greenland and Antarctica during the Holocene. *Journal of Geophysical Research*, 102: 15987-15997.
- Cheddadi, R.; Lamb, H.F.; Guiot, J. & van der Kaars, S. (1998).** Holocene climatic change in Morocco: a quantitative reconstruction from pollen data. *Clim. Dyn.*, 14: 883-890.
- Chueca, J. & Julián, A. (1995).** Cuantificación de movimientos de masa lentos en medios de montaña: Pirineo Central. *Lurralde*, 18: 173-196.
- Chueca, J.; Peña Monné, J.L.; Lampre F. & Julián, A. (1998).** La Pequeña Edad de Hielo en el Pirineo central y meridional. Inferencias paleoambientales a partir de datos geomorfológicos. In: Gómez Ortiz i Pérez Alberti (Eds.). *Las huellas glaciares de las montañas españolas*. Servicio de publicaciones Universidad de Santiago. pp. 307-328.
- Climap (1981).** Seasonal reconstructions of the Earth's surface at the last glacial maximum, *Geol. Soc. Am. Map Chart Ser.*, MC-36.
- Colom, G. (1967).** Les lacs du Burdigalien supérieur de l'île de Majorque (Baléares) et le rôle des méolositres (Diatomées) dans la formation de leurs varves. *Bull. Soc. Geol. France*, 7, 9: 835-843.
- Combourieu-Nebout, N.; Turon, J. L.; Zahn, R.; Capotondi, L.; Londeix, L. & Pahnke, K. (2002).** Enhanced aridity and atmospheric high-pressure stability over the western Mediterranean during the North Atlantic cold events of the past 50 k.y., *Geology*, 30: 863-866.
- Creus, J. & García Ruiz, J.M. (1977).** Observaciones sobre soliflucción en el alto valle del río Gállego. *Trabajos sobre Neógeno- Cuaternario*, 6: 57-64.
- Crowdace I.W.; Rindby A. & Rothwell R.G. (2006).** ITRAX: description and evaluation of a new sediment core scanner. In: Rothwell, R.G. (ed.) *New techniques in sediment core analysis*. London, UK, Geological Society of London, 51-63.
- Daansgard, W., Johnsen, S.J., Clausen, Dahl-Jensen, D., Gundestrup, N.S., Hammer, C.U., Hvidberg, C.S., Steffensen, J.P., Sveinbjornsdottir, A.E., Jouzel, J. & Bond, G. (1993).** Evidence for general instability of past climate from a 250-kyr ice-core record, *Nature*, 364: 218-220.

- Damnati, B. (2000).** Holocene lake records in the Northern Hemisphere of Africa. *Journal of African Earth Sciences* 31: 253–262.
- De Beaulieu, J.L.; Miras, Y.; Andrieu-Ponel, V. & Guiter, F. (2005).** Vegetation dynamics in north-western Mediterranean regions: instability of the Mediterranean bioclimate. *Plant Biosystems*, Vol. 139 (2): 114 – 126.
- deMenocal, P.B. (2001).** Cultural responses to climate change during the late Holocene, *Science*, 292: 667-673.
- Desprat, S.; Goñi, M.F.S. & Loutre, M.F. (2003).** Revealing climatic variability of the last three millennia in northwestern Ibera using pollen influx data. *Earth and Planetary Science Letters* 213: 63-78.
- Dyke, A. S. (1981).** Late Holocene solifluction rates and radiocarbon soil ages, central Canadian Arctic. *Geological Survey of Canada Paper*, 81: 17–22.
- ELDP (1995).** *European Lakes Drilling Program*. European Science Foundation.
- Elliott, G. & Worsley, P. (1999).** The sedimentology, stratigraphy and <sup>14</sup>C dating of a turf-banked solifluction lobe: evidence for Holocene slope instability at Okstindan, northern Norway. *Journal of Quaternary Science*, 14 (2): 175-188.
- Esteban Amat, A. (1995).** *Evolución del paisaje durante los últimos 10.000 años en las montañas del Mediterráneo Occidental: ejemplos del Pirineo Oriental y Sierra Nevada*. Tesis doctoral. Universitat de Barcelona.
- FAO (2006).** Guidelines for soil description. United Nations, 133 pp.
- Florschütz, F.; Menéndez Amor J. & Wijmstra, T.A. (1971).** Palinology of a thick Quaternary succession in southern Spain. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 10: 233–264.
- French, H.M. (1996).** *The Periglacial Environment*, 2nd ed. Longman, Harlow. 341 pp.
- Friend, P.J. (1988).** *High latitude gelifluction lobes as paleoclimatic indicators*. Unpublished master's thesis, University of Alberta, Edmonton, Alberta.
- Galop, D. (1997).** *La forêt, l'homme et le troupeau. Six millénaires d'anthropisation du massif pyrénéen de la Garonne à la Méditerranée*. Tesis doctoral, Université Toulouse le Mirail. 322 pp.
- Gamper, M.V. (1983).** Controls and rates of movement of solifluction lobes in the eastern Swiss Alps., *Proc. IV Int. Conf. Permafrost*, Fairbanks, pp. 328-333.
- García Ruiz, J.M.; Chueca, P. & Julián, A. (2004).** Los movimientos en masa del Alto Gállego. In: Peña, J.L.; Longares, L.A. & Sánchez, M. *Geografía Física de Aragón: aspectos generales y temáticos*. 141-152.
- Gasse, F. (2000).** Hydrological changes in the African tropics since the Last Glacial Maximum. *Quaternary Science Reviews*, 19: 189–211.
- Gasse, F., 2002.** Diatom-inferred salinity and carbonate oxygen isotopes in Holocene waterbodies of the western Sahara and Sahel (Africa). *Quaternary Science Reviews*, 21: 717-767.
- Gerrard, J. (2000).** *Fundamentals of soils*. Routledge, London, New York. 230 pp.
- Gil García, M.J.; Ruiz Zapata, M.B.; Santisteban, J.I.; Mediavilla, R.; López Pamo, E. & Dabrio, C.J. (2006).** Late Holocene environments in Las Tablas de Daimiel (South central Iberian Peninsula, Spain). *Vegetation History and Archaeobotany*. 10.1007/s00334-006-0047-9.



- Gómez Ortiz A (1980).** *Estudio geomorfológico del Pirineo catalán: morfogénesis glacial y periglacial de los altos niveles y vertientes meridionales de los macizos de Calmquerdós, Tossa Plana de Llès y Port Negre (Cerdanya-Alt Urgell)*. Tesis doctoral, Departament de Geografia, Universitat de Barcelona, Barcelona.
- Gómez Ortiz, A. (1987).** Morfología glaciaria en la vertiente meridional de Sierra Nevada (Área Veleta-Mulhacén). *Estudios Geográficos*, 193: 527-558.
- Gómez Ortiz, A. & Salvador Franch, F. (1992).** Aportaciones significativas al conocimiento del glaciario y periglaciario de Sierra Nevada. *Notes de Geografia Física*, 20-21: 89-101.
- Gómez Ortiz, A.; Simón Torres, M.; Salvador Franch, F.; Sánchez Gómez & Esteban Amat, A. (1993).** El Cuaternario de los niveles cimeros de Sierra Nevada (Guía de la Excursión). In *Periglaciario en la península Ibérica, Canarias y Baleares. Guía de la excursión (Granada-Sierra Nevada, 1993)*. SEG. Barcelona, 72 pp.
- Gómez Ortiz, A. (1997).** Hugo Obermaier Grat (1877-1946): su contribución al conocimiento del glaciario de las montañas españolas y particularmente al de Sierra Nevada. A: *Los paisajes cuaternarios de Sierra Nevada. Hugo Obermaier*. Colección Sierra Nevada y la Alpujarra, 19. Granada, 63 pp.
- Gómez Ortiz, A.; Palacios Estremera, D.; Ramos Saínz, M.; Schulte, L.; Salvador Franch, F. & Tanarro García, L. (1999).** Degradación de permafrost en Sierra Nevada. Repercusiones geomorfológicas: el caso del Corral del Veleta. Resultados preliminares. A Gómez Ortiz, A. & Martí Bono, C. M. (eds). *Boletín de la Asociación de Geógrafos Españoles (AGE)*, 27, pp. 7-21.
- Gómez Ortiz, A. & Pérez González, A. (eds) (2001):** *Evolución reciente de la Geomorfología española (1980-2000). Aportación española a la V Conferencia Internacional de Geomorfología (Tokyo 2001)*. Sociedad Española de Geomorfología. Servei de Gestió i Evolució del Paisatge (UB). Barcelona. 427 pp.
- Gómez Ortiz, A. (coord.) (2002).** *Mapa geomorfológico de Sierra Nevada*. Consejería de Medio Ambiente, Junta de Andalucía.
- Gómez Ortiz, A.; Palacios Estremera, D.; Luengo Nicolau, E.; Tanarro García, L. M.; Schulte, L.; Ramos Saínz, M. & Salvador Franch, F. (2002).** Inestabilidad de taludes y cubierta nival en áreas marginales de permafrost. El caso de la pared norte del picacho del Veleta (Sierra Nevada, España). *Revista de Geografía*, Vol. 1. pp. 35-56.
- Gómez Ortiz, A.; Palacios, D. & Ramos, M. (2004).** Permafrost, evolution of associated landforms and thermal control in the Veleta Cirque (Sierra Nevada, Spain). New results. *Boletín de la Real Sociedad Española de Historia Natural (Sec. Geol.)*, 99 (1-4): 47-63.
- Gómez Ortiz, A.; Schulte, L.; Salvador Franch, F.; Palacios, D.; Sanz De Galdeano, C.; Sanjosé Blasco, J. J.; Tanarro García, L.M. & Atkinson, A. (2005).** Field trip to Sierra Nevada massif glacial geomorphology and present cold processes. *Sixth International Conference on Geomorphology. Field Trip Guides*: 309-326.
- Gómez, Ortiz, A.; Milheiro, B. & Serrano, D. (2008).** Aguas, nieves e hielos en los paisajes de Sierra Nevada. El interés de la información gráfica de los libros de época. *Cuadernos de investigación Geográfica*, 34: 99-118.

- González Trueba, J.J.; Martín Moreno, R.; Martínez de Pisón, E. & Serrano, E. (2008).** Little Ice Age glaciation and current glaciers in the Iberian Peninsula. *The Holocene*, 18 (4): 551-568.
- Gorbutnov, A. & Seversky, E.V. (1999).** Solifluction in the mountains of Central Asia: distribution, morphology and processes. *Permafrost and periglacial processes*, 10: 81-89.
- Grab, S. (2000).** Stone-banked lobes and environmental implications, High Drakensberg, Southern Africa. *Permafrost and periglacial processes*, 11: 177-187.
- Grimalt, M. & Rodríguez, A. (1994).** El modelado periglacial en Baleares. Estado de la cuestión. *Periglaciario en la Península Ibérica, Canarias y Baleares*. Servicio de Publicaciones Universidad Granada: 189-201.
- Håkanson, L. & Jansson, M. (2002).** *Principles of Lake Sedimentology*, Springer-Verlag, Berlin. 312 pp.
- Harris, C. (1981).** Periglacial Mass-Wasting: A Review of Research. *BGRG Research Monograph*, vol.4.
- Harris, C.; Davies, M. & Coutard, J.P. (1997).** Rates and processes of periglacial solifluction: an experimental approach. *Earth Surface Processes and Landforms*, 22: 849-868.
- Harris, C.; Davies, M. & Rea, B. (2003).** Gelifluction: viscous flow or plastic creep? *Earth Surface Processes Landforms*, 28: 1289-1301.
- Harris, C.; Smith, J.S.; Davies, M.C.R. & Rea, B.R. (2007).** An investigation of periglacial slope stability in relation to soil properties based on physical modelling in the geotechnical centrifuge. *Geomorphology*, 93: 437-459.
- Heinrich, H. (1988)** Origin and consequence of cycling ice rafting in the Northeast Atlantic Ocean during the past 130, 000 years. *Quaternary Research*, 29: 142-152.
- Heiri, O.; Lotter, A.F. & Lemcke, G. (2000).** Loss of ignition as a method for estimating organic and carbonate content in sediments: Reproducibility and comparability of results. *Journal of Paleolimnology*, 25: 101-110.
- Hellmann, G. (1881).** Der südlichste Gletscher Europas. *Zeitschrift Gesellschaft für Erdkunde*, 16: 362-367.
- Hempel, L. (1960).** Límites altitudinales geomorfológicos en Sierra Nevada. *Estudios Geográficos*, 78: 81-93.
- Higashi, A. & Corté, A.E. (1971).** Solifluction: a model experiment. *Science*, 171: 480-482.
- Hoelzle, M.; Wegmann, M. & Krummenacher, B. (1999).** Miniature temperature dataloggers for mapping and monitoring of permafrost in high mountain areas: first experience from the Swiss Alps. *Permafrost and Periglacial Processes* 10: 113-124.
- Höfer, H. V. (1879).** Gletscher und Eiszeitstudien. *Sitzungsberichte der Academie der Wissenschaften in Wien, Mathematische-naturwissenschaftliche Klasse*, 79.
- Hoffert, M. & Covey, C. (1992).** Deriving global climate sensitivity from paleoclimate reconstructions, *Nature*, 360 (6404): 573-576.
- Holzhauser, H.; Magny, M. & Zumbühl H.J. (2005).** Glacier and lake-level variations in west-central Europe over the last 3500 years. *The Holocene* 15 (6): 789-801.
- Hormes, A.; Müller, B.U. & Schlüchter, C. (2001).** The Alps with little ice: evidence for eight Holocene phases of reduced glacier extent in the Central Swiss Alps. *The Holocene*, 11: 255-265.

- Hormes A.; Beer J. & Schlüchter C. (2006).** A geochronological approach to understanding the role of solar activity on Holocene glacier length variability in the Swiss Alps. *Geografiska Annaler: Series A, Physical Geography* 88: 281–294.
- Hugenholtz, C.H. & Lewkowicz, A.G. (2002).** Morphometry and environmental characteristics of turf-banked solifluction lobes, Kluane Range, Yukon Territory, Canada. *Permafrost Periglac. Process.* 13: 301–313.
- Hughes, M.K. & Gramlich L.J. (1996).** Multimillennial dendroclimatic records from Western North America. In: Jones, P.D.; Bradley, R.S. & Jouzel, J. (ed.). *The impacts of climate variability and forests.* Springer, Berlin, pp. 99-107.
- Hughes, P.D & Woodward, J. C. (2008).** Timing of glaciation in the Mediterranean mountains during the last cold stage. *Journal Of Quaternary Science*, 23(6-7): 575–588.
- Huntley, B. & Prentice, I.C. (1993).** *Holocene vegetation and climates of Europe. Global climates since the last glacial maximum.* University of Minnesota, Minneapolis, pp. 136-168.
- IPCC (2001).** *Intergovernmental Panel on Climate Change. World Meteorological Organization.*
- IPCC (2007).** *Intergovernmental Panel on Climate Change. World Meteorological Organization.*
- Jackson, J.A. (1997).** *Glossary of Geology.* American Geological Institute. Fourth Edition, Virginia. 769 pp.
- Jacobeit, J., Glaser, R., Luterbacher, J., Nonnenmacher, M. & Wanner, H. (2002).** Links Between Flood Events In Central Europe Since AD 1500 And The Large-Scale Atmospheric Circulation. In: Torndycraft, V.R.; Benito, G.; Barriendos, M. & Llasat, M.C. *Paleofloods, Historical Floods and Climate Variability: Applications in Flood Risk Assessment.* Proceedings of PHEFRA Workshop, Barcelona, 16 – 19th October, pp. 269-274.
- Jacoby G.C.; Workman, K.W. & D'Arrigo, R.D. (1999).** Laki eruption of 1783, tree rings and disaster for northwest Alaska Inuit. *Quaternary Science Reviews*, 18: 1365-1371.
- Jaesche, P.; Veit, H. & Huwe, B. (2003).** Snow cover and soil moisture controls on solifluction in an area of seasonal frost, Eastern Alps. *Permafrost and Periglacial Processes*, 14: 399-410.
- Jalut, G.; Esteban, A.; Bonnet, L. ; Gauquelin, T. & Fontugne, M. (2000).** Holocene climatic changes in the western Mediterranean, from south-east France to south-east Spain. *Palaeogeography, Palaeoclimatology, Palaeoecology* 160: 255-290.
- Jansà Guardiola, J. J. (1966).** *Meteorología del Mediterráneo Occidental.* Servicio Meteorológico Nacional. Serie A (Memorias), 43. Madrid.
- Jansen J.H.F.; Van der Gaast S.J.; Koster B. & Vaars A.J. (1998).** CORTEX, a shipboard XRF-scanner for element analyses in split sediment cores. *Marine Geology*, 151:143–153.
- Jiang, H.; Eriksson, J.; Schulz, M.; Knudsen, K.L. & Seidenkrantz, M.S. (2005).** Evidence for solar forcing of sea-surface temperature on the North Icelandic Shelf during the late Holocene. *Geology, Geological Society of America*, 1: 73-76.
- Jiménez, J.I. (1996).** Régimen de precipitación en Sierra Nevada y cambio climático. *1ª Conferencia Internacional Sierra Nevada*, 20-26 de marzo de 1996, pp. 23-40.
- Johnsen, S.; Dahl-Jensen, D.; Gunderstrup, N.; Steffensen, J.P.; Clausen, H.B.; Miller, H.; Masson-Delmotte, V.; Sveinbjörnsdóttir, A.E. & White, J. (2001).** Oxygen isotope and paleotemperature

- records from six Greenland ice-core stations: Camp Century, Dye-3, GRIP, GISP2, Renland and NorthGRIP. *Journal of Quaternary Science*, 16: 299-308.
- Jonasson, M. (1993).** Solifluction in northern Finland: past and present. In: Frenzel, B., Matthews, J.A., Gläser, B. (eds). *Solifluction and Climatic Variation in the Holocene*. Paläoklimaforschung, Vol. 11: 179-195.
- Julià, R. (1977).** *Estudio de la cuenca lacustre de Banyoles-Besalú*. Tesi doctoral. Universitat de Barcelona, 180 pp.
- Julià, R.; Burjachs, F. & Giral, S. (1996).** Redepositional or Climate Event in Iberia during the Boreal. *Journal of Quaternary Science*, 12 (1): 73-75.
- King, L. & Smith, E. (1993).** Solifluction and thermal erosion in high arctic ecosystems, Northern Spitsbergen: processes and effects. In: Frenzel, B., Matthews, J.A., Gläser, B. (eds). *Solifluction and Climatic Variation in the Holocene*. Paläoklimaforschung, Vol. 11: 267-278.
- Kinnard, C. & Lewkowicz, A.G. (2005).** Movement, moisture and thermal conditions at a turf-banked solifluction lobe, Kluane Range, Yukon Territory, Canada. *Permafrost and Periglacial Processes*, 16: 261-275.
- Kinnard, C. & Lewkowicz, A.G. (2006).** Frontal advance of turf-banked solifluction lobes, Kluane Range, Yukon Territory, Canada. *Geomorphology*, 73: 261-276.
- Koç, N.; Jansen, E. & Hafliðason, H. (1993).** Paleoceanographic reconstructions of surface ocean conditions in the Greenland, Iceland and Norwegian Seas through the last 14 ka based on diatoms. *Quaternary Science Reviews* 12: 115-140.
- Koshkarova, V.L. & Koshkarov, A.D. (2004).** Regional signatures of changing landscape and climate of northern central Siberia in the Holocene. *Russian Geology and Geophysics* 45 (6): 672-685.
- Kuhlemann, J.; Rohling, E.J.; Kumrei, I.; Kubik, P.; Ivy-Ochs, S. & Kucera, M. (2008).** Regional synthesis of Mediterranean atmospheric circulation during the Last Glacial Maximum. *Science*, 321 (5894): 1338-1340.
- Labeyrie, L.; Cole, J.; Alverson, K. & Stocker, T. (2003).** The history of climate dynamics in the Late Quaternary. A: Alverson, K.D.; Bradley, R.S. i Pedersen, T.F. *Paleoclimate, global change and the future*. Springer, Berlin, 33-61 pp.
- Lacomba, J.A. (2001).** Historia de Andalucía. Ágora, 2a ed. Málaga. 434 pp.
- Lamb, H. (1970).** Volcanic dust in the atmosphere; with a chronology and assessment of its meteorological significance. *Philosophical Transactions of the Royal Society of London*, A266: 425-533.
- Lamb, H.H.. (1977).** *Climate, present, past and future*. Vol. 2.; William Claus and Sons, 835 pp.
- Lampre, F. (1994).** La línea de equilibrio glacial y los suelos helados en el macizo de la Maladeta (Pirineo Aragonés): evolución desde la Pequeña Edad de Hielo y situación actual. A: Martí Bono i García Ruiz (Eds.). *El glaciarismo surpirenaico: Nuevas aportaciones*. Geoforma. Logroño. pp. 125-142.
- Last, W.M. (2001).** Textural analysis in lake sediments. In: Last, W.M. & Smol, J.P (eds). *Tracking Environmental Changes Using Lake Sediments*. Kluwer Academic Publishers. Holanda. Vol. 2: 41-73.
- Lean, J.; Beer, J. & Bradley, R. (1995).** Reconstruction of solar irradiance since 1610: Implications for climate change. *Geophysical Research Letters*, vol. 22 (23): 3195-3198.

- Leira, M. (2005).** Diatom responses to Holocene environmental changes in a small lake in northwest Spain. *Quaternary International*, 140–141: 90–102.
- Lewkowicz, AG. (1988).** Slope processes. In Clark, MJ (Eds). *Advances in Periglacial Geomorphology*, John Wiley and Sons. Chichister, 325-368.
- Lhenaff, R. (1977).** *Recherches géomorphologiques sur les Cordilleres Betiques centro-occidentales (Espagne)*. Thèse. Université de Lille.
- Long, A.J.; Roberts, D.H. & Daws, S. (2006).** Early Holocene history of the west Greenland Ice Sheet and the GH-8.2 event. *Quaternary Science Reviews*, 25: 904–922.
- López Moreno, J.I. (2000).** *Los glaciares de alto valle del Gállego (Pirineo aragonés) desde la Pequeña Edad de Hielo. Implicaciones en la evolución de las temperaturas*. Ed. Geoforma. Logroño. 77 pp.
- Löye-Pilot, M.D., Martin, J.M. & Morelli, J., (1986).** Influence of Saharan dust on the rainfall acidity and atmospheric input to the Mediterranean. *Nature* 321: 427–428.
- Luterbacher, J., Dietrich, D., Xoplaki, E., Grosjean, M., & Wanner, H. (2004).** Extreme seasonal and annual temperature variability, trends and extremes since 1500. *Science*, 303: 1499–1503.
- Mackay, A.,; Battarbee, R.; Birks, J. & Oldfield, F. (2003).** *Global change in the Holocene*. Holder Headline, London. 527 pp.
- Magny, M.; Bégeot, C.; Guiot, J. & Peyron, O. (2003).** Contrasting patterns of hydrological changes in Europe in response to Holocene climate cooling phases. *Quaternary Science Reviews* 22: 1589–1596.
- Mahaney, W. & Sanmugadas, K. (1990).** Extractable Al and Fe in early to Middle Quaternary paleosoils in Gorges valley, Mount Kenya, East Africa. *Catena*, 17: 563-572.
- Maher, B.A. & Thompson. R. (1995).** Paleorainfall reconstructions from pedogenic magnetic susceptibility variations in the Chinese loess and paleosoils. *Quaternary Research*, 44: 383-391.
- Mailänder, R. & Veit, H. (2001).** Periglacial cover-beds on the Swiss Plateau: indicators of soil, climate and landscape evolution during the Late Quaternary. *Catena*, 45 (4): 251-272.
- Mangerud, J.; Andersen, S.T.; Berglund, B.E. & Donner, J.J. (1974).** Quaternary stratigraphy of Norden, a proposal for terminology and classification. *Boreas* 3: 109–127.
- Mann, M.E.; Bradley, R.S. & Hughes, M.K. (1998).** Global-scale temperature patterns and climate forcing over the past six centuries. *Nature*, 392: 779-785.
- Margalef, R. (1947).** Observaciones micropaleontológicas sobre los sedimentos lacustres miocénicos de Libros (Teruel). *Estudios Geológicos*, 5: 171-177.
- Margalef, R. (1957).** Paleoecología del lago miocénico de la Cerdanya.. *Instituto B. Biol. Apl.*, 25: 131-137.
- Martín Civantos, J. M. (2007).** *Poblamiento y territorio medieval en el Zenete (Granada)*. Universidad de Granada.
- Martín García, J.M.; Aranda, V.; Gámiz, E.; Bech, J. & Delgado, R. (2004).** Are Mediterranean mountain Entisols weakly developed? The case of Orthents from Sierra Nevada (Southern Spain). *Geoderma*, 118: 115-130.
- Martínez Cortizas, A.; Pontevedra-Pombal, X.; García-Rodeja, E.; Novoa-Muñoz, J.C. & Shotyk, W. (1999).** Mercury in a Spanish peat bog: archive of climate change and atmospheric metal deposition. *Science* 284: 939–942.



- Martínez-Parras, J.M.; Peinado, M. & Alcaraz, F. (1987).** Datos sobre la vegetación de Sierra Nevada. *Lazaroa* 7: 515-533.
- Masson-Delmotte, V.; Jouzel, J.; Landais, A.; Stievernard, M.; Johnsen, S.J.; White, J.W.C.; Werner, M.; Sveinbjornsdottir, A. & Fuhrer, K. (2005).** GRIP deuterium excess reveals rapid and orbital-scale changes in Greenland moisture origin. *Science*, 309: 118-121.
- Matsuoka, N. (1996).** Soil moisture variability in relation to diurnal frost heaving on Japanese high mountain slopes. *Permafrost Periglacial Processes*, 7: 139-151.
- Matsuoka, N. (2001).** Solifluction rates, processes and landforms: a global review. *Earth-Science Reviews*, 55: 107-134.
- Matsuoka, N. (2006).** Monitoring periglacial processes: towards a construction of a global network. *Geomorphology*, 80: 20-31.
- Matsuoka, N. & Humlun, O. (2003).** Monitoring Periglacial Processes: New Methodology and Technology. *Permafrost and Periglacial Processes*, 14: 299–303.
- Matsuoka, N., Ikeda, A. & Date, T. (2005).** Morphometric analysis of solifluction lobes and rock glaciers in the Swiss Alps. *Permafrost Periglac. Process*, 16: 99-113.
- Matthews, J.A.; Ballantyne C.K.; Harris, C.; Berrisford, M.S. (1993).** Solifluction and climatic variation in the Holocene: discussion and synthesis. In: Frenzel, B., Matthews, J.A., Gläser, B. (eds). *Solifluction and Climatic Variation in the Holocene*. Paläoklimaforschung, Vol. 11. Gustav Fischer Verlag, Stuttgart: 339-361.
- Matthews, J.A.; Seppälä, M. & Dresser P.Q. (2005).** Holocene solifluction, climate variation and fire in a subarctic landscape at Pippokangas, Finnish Lapland, based on radiocarbon-dated buried charcoal. *Journal of Quaternary Science*, 20: 533-548.
- May, J.H.; Zech, R. & Veit, H. (2008).** Late Quaternary paleosol-sediment-sequences and landscape evolution along the Andean piedmont, Bolivian Chaco. *Geomorphology*. 98 (1-2): 34-54.-
- Mayewski, P.A.; Rohling, E.E.; Stager, J.C.; Karlen, W.; Maasch, K.A.; Meeker, L.D.; Meyerson, E.A.; Gasse, F.; Van Kreveld, S.; Holmgren, K.; Lee-Thorp, J.; Rosqvist, G.; Rack, F.; Staubwasser, M.; Schneider R.R. & Steig, E.J. (2004).** Holocene climate variability. *Quaternary Research*, 62: 243-255.
- Medina-Sánchez, J. M.; Villar-Argaiz, M.; Sánchez-Castillo, P.; Cruz-Pizarro, L. & Carrillo, P. (1999).** Structure changes in a planktonic food web: biotic and abiotic controls. *Journal of Limnology*, 58: 213-222.
- Meeker, L.D. & Mayewski, P.A. (2002).** A 1400-year high-resolution record of atmospheric circulation over the North Atlantic and Asia. *Holocene* 12: 257–266.
- Mehra, O.P. & Jackson, M.L. (1960).** Iron oxide removal from soils and clays by dithionit-citrate systems buffered with sodium bicarbonate. *Clays and Clay Minerals*, 7: 317-327.
- Meier, M.F. & Post, A.S. (1962).** Recent variations in mass net budgets of glaciers in western North America. *International Association of Scientific Hydrology*, 58: 63-77.
- Menéndez Amor J. & Florschütz, F. (1964).** Results of the preliminary palynological investigation of samples from a 50 m boring in southern Spain. *Boletín de la Real Sociedad Española de Historia Natural (Sec.Geología)* 62: 251–255.

- Messerli, B. (1965).** *Beiträge zur geomorphologie der Sierra Nevada (Andalusien)*. Juris Verlag. Zürich.
- Meyers, P.A. & Ishiwatari, R. (1995).** Organic matter accumulation records in lake sediments. In: Lerman, A.; Imbroden, D. i Gat, J. (eds.). *Physics and Chemistry of lakes*. Springer. Germany, 332 pp.
- Meyers, P.A. & Teranes, J.L. (2001).** Sediment organic matter. In: Last, W.M. & Smol, J.P (eds). *Tracking Environmental Changes Using Lake Sediments*. Kluwer Academic Publishers, Dordrecht. Vol. 2: 239–270.
- Miehe, G. (1995).** On the connexion of vegetation dynamics with climatic changes in High Asia. *Palaogeography, Palaeoclimatology, Palaeoecology*, 120: 5-24.
- Molero Mesa, J. & Pérez Raya, F. (1987).** *La flora de Sierra Nevada*. Universidad de Granada.
- Morales-Baquero, R.; Pulido-Villena, E. & Reche, I. (2006).** Atmospheric inputs of phosphorus and nitrogen to the Southwest Mediterranean region: Biogeochemical responses of high mountain lakes. *Limnology and Oceanography* 51: 830-837
- Moreno A.; Nave S.; Kuhlmann H.; Canals M.; Targarona J.; Freudenthal T., & Abrantes F. (2002)** Productivity response in the North Canary Basin to climate changes during the last 250,000 yr: a multi-proxy approach. *Earth Planetary Science Letters*, 196: 147–159.
- Moreno, A.; Cacho, I.; Canals, M.; Grimalt, J.O.; Sánchez-Goñi, M.F.; Shackleton, N. & Sierro, F.J. (2005).** Links between marine and atmospheric processes oscillating on a millennial time-scale. A multi-proxy study of the last 50,000 yr from the Alboran sea. *Quaternary Science Reviews* 24: 1623–1636.
- Morin H. & Payette S. (1988).** Holocene gelifluction in a snow-patch environment at the forest–yundra transition along the eastern Hudson Bay Coast, Canada. *Boreas* 17: 79–88.
- Nesje, A. & Dahl, S.O. (1993).** Lateglacial and Holocene glacier fluctuations and climate variations in western Norway: a review. *Quaternary Science Reviews*, 12: 255-261.
- Nesje, A. & Dahl, S.O. (2000).** *Glaciers and environmental change*. Arnold, London.
- Nowaczyk, N.R. (2001).** Logging of magnetic susceptibility. In: Last, W.M. & Smol, J.P (eds). *Tracking Environmental Changes Using Lake Sediments*. Kluwer Academic Publishers, Dordrecht. Vol.1: 155–170.
- Nuzhet Dalfes, H.; Kukla, G. & Weiss, H. (eds) (1996).** Third millenium BC climate change and Old World collapse. NATO ASI Series, Serie I. *Global Environmental Change*: 49. Springer, 709 pp.
- Oliva, M.; López-Bustins, J.A.; Barriendos, M.; Muedra, C. & Martín-Vide, J. (2006).** Reconstrucción histórica de la oscilación del Mediterráneo Occidental (WEMO) e inundaciones en el Levante Peninsular (1500-2000). in Cuadrat Prats, J.M.; Saz Sánchez, M.A.; Vicente Serrano, S.M.; Lanjeri, S.; De Luis Arrillaga, M. & González-Hidalgo, J.C. (Eds): *Clima, Sociedad y Medio Ambiente. Publicaciones de la Asociación Española de Climatología*, AEC, Serie A, nº 5, 241-250.
- Oliva, M. (2008).** Lóbulos de soliflución y fluctuaciones paleoclimáticas durante el Holoceno en Sierra Nevada. *Publicaciones de la Asociación Española de Climatología*, Serie A, nº 6, pp. 189-198.
- Oliva, M. & Moreno I. (2008).** Sierra Nevada, nexo entre dos patrones de teleconexión: la NAO y la WeMO. *Publicaciones de la Asociación Española de Climatología*, Serie A, nº 6, pp. 199-208.
- Oliva, M.; Schulte, L. & Gómez Ortiz, A. (2008).** Solifluction Lobes in Sierra Nevada (Southern Spain): Morphometry and Palaeoenvironmental Changes. *Proceedings of the IX International Conference on Permafrost*. Fairbanks (Alaska), 29-3 July 2008.

- Oliva, M.; Schulte, L. & Gómez Ortiz, A. (accepted).** Approach to solifluction lobes in Sierra Nevada (Southern Spain): morphometry, dynamism and palaeoenvironmental implications. *Permafrost and Periglacial Processes*.
- Ortiz, A.; Torres, T.; Delgado, A.; Julià, R.; Lucini, M. & Llamas, F.J. (2004).** The palaeoenvironmental and palaeohydrological evolution of Padul Peat Bog (Granada, Spain) over one million years, from elemental, isotopic and molecular organic geochemical proxies. *Organic Geochemistry*, 35: 1243–1260.
- O'Sullivan, P. (2004).** Paleolimnology. In: O'Sullivan, P.E. & Reynolds, C.S. (Eds.). *The lakes handbook. Limnology and limnetic ecology*. Vol. 1. Blackwell Publishing. UK. 699 pp.
- Palacios, D; Gómez Ortiz, A.; Fernández Fernández, A.; Zamorano Orozco, J.J.; Tanarro García, L.M.; Salvador Franch, F.; Schulte, L. & Ramos Sainz, M. (2000).** La formación de debris flows en el Corral del Veleta (Sierra Nevada, España). Geomorfología para el Tercer Milenio y Sociedad. *VI Reunión Nacional de Geomorfología*, Madrid. pp. 133.
- Palacios, D.; Andrés, N. de & Luengo, E. (2003).** Distribution and effectiveness of nivation in Mediterranean mountains: Peñalara (Spain). *Geomorphology* 54, 157-178.
- Pallàs, R.; Rodes, A.; Braucher, R.; Carcaillet, J.; Ortuno, M.; Bordonau, J.; Bourles, D.; Vilaplana, J.M.; Masana, E. & Santanach, P. (2006).** Late Pleistocene and Holocene glaciation in the Pyrenees: A critical review and new evidence from <sup>10</sup>Be exposure ages, south-central Pyrenees. *Quaternary Science Reviews*, 25: 2937-2963.
- Pauling, A. ; Luterbacher, J & Wanner, H. (2003).** Evaluation of proxies for European and Horth Atlantic temperature field reconstructions. *Geophysical Research Letters*, 30 (2-1): 2-4.
- Pauling, A.; Luterbacher, J.; Casty, C. & Wanner, H. (2006).** Five hundred years of gridded high-resolution precipitation reconstructions over Europe and the connection to large-scale circulation. *Climate Dynamics*, 26: 387-405.
- Pérez Alberti, A., Valcárcel Díaz, M. & Blanco Chao, R. (2004).** Pleistocene glaciation in Spain. In Ehlers, J. & Gibbard, P.L.(editors), *Quaternary glaciations – extent and chronology*. Vol. 2: Part I. Europe. Elsevier, 389–94.
- Pérez Raya, F.; López Nieto, J.; Molero, Mesa, J. & Valle, F. (1990).** *Vegetación de Sierra Nevada*. Granada, Ayuntamiento de Granada.
- Pons, A. & Reille, M. (1988).** The Holocene and upper Pleistocene pollen record from Padul (Granada, Spain): a new study. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 66: 243–263.
- Porter, S. G. & Denton, G. H. (1967).** Chronology of Neoglaciation in the North American Cordillera. *Am. J. Sci.* 265: 177-210.
- Porter, S.C. (1986).** Pattern and forcing of northern hemisphere glacier variations during the last millenium. *Quaternary Research*, 26: 27-48.
- Porter, S.C. (2001).** Snowline depression in the tropics during the Last Glaciation. *Quaternary Science Reviews* 20: 1067-1091.
- Price, L.W. (1969).** The collapse of solifluction lobes as a factor in vegetating blockfield. *Arctic* 22: 395-402.
- Price, L.W.. (1971).** Vegetation, microtopography, and depth of active layer on different exposures in subarctic alpine tundra. *Ecology*, 52 (4): 638-647.



- Prohom, M.J.; Esteban, P.; Martín Vide, J. & Jones, P.D. (2003).** Surface atmospheric circulation over Europe following major tropical volcanic eruptions, 1780-1995. *Volcanism and the Earth's Atmosphere, Geophysical Monograph* 139: 273-281. American Geophysical Union.
- Pulido-Villena, E. (2004).** *El papel de la deposición atmosférica en la biogeoquímica de lagunas de alta montaña (Sierra Nevada, España)*. Tesis doctoral. Universitat de Granada. 296 pp.
- Pulido-Villena, E.; Reche, I. & Morales-Baquero, R. (2005).** Food web reliance on allochthonous carbon in two high mountain lakes with contrasting catchments: a stable isotope approach. *Can. J. Fish. Aquat. Sci.*, 62: 2640-2648.
- Pulido-Villena, E.; Reche, I. & Morales-Baquero, R. (2006).** Significance of atmospheric inputs of calcium over the southwestern Mediterranean region: High mountain lakes as tools for detection. *Global Biogeochemical Cycles* 20: GB2012, doi:10.1029/2005GB002662
- Ramrath, A.; Sadori, L. & Negendank, J.F.W. (2000).** Sediments from Lago di Mezzano, central Italy: a record of Lateglacial/Holocene climatic variations and anthropogenic impact. *The Holocene* 10: 87-95.
- Rapp, A. (1960).** Recent development of mountain slopes in Kärkevagge and surroundings, Northern Scandinavia. *Geografiska Annaler*, 42A: 65-200.
- Raynaud, D.; Barnola, J.M.; Chappellaz, J.; Blunier, T.; Indermühle, A. & Stauffer, B. (2000).** The ice record of greenhouse gases: a view in the context of future changes. *Quaternary Science Reviews* 19: 9-17.
- Reimer, RW; Remmele, S; Southon, JR; Stuiver, M.; Talamo, S; Taylor, FW; Van der Plicht, J; Weyhenmeyer, CE (2004).** IntCal04 Atmospheric radiocarbon age calibration, 26-0 ka BP. *Radiocarbon* 46: 1029-1058
- Rivas Martínez, S. (1983).** Nuevo índice de termicidad para la región mediterránea. *VIII Reunión-Ponencia de Bioclimatología del CSIC, Zaragoza*.
- Rodrigo, F.S. (1994).** *Cambio climático natural. La Pequeña Edad de Hielo en Andalucía*. Tesis doctoral. Universitat de Granada.
- Rodrigo, F.S., Esteban-Parra, M.J., Pozo-Vázquez, D. & Castro-Díaz, Y. (1999).** A 500-year precipitation record in southern Spain. *International Journal of Climatology*, 19: 1233-1253.
- Rodrigo, F.S.; Esteban-Parra, M.J.; Pozo-Vázquez, D. & Castro-Díez, Y. (2000).** Rainfall variability in southern Spain on decadal to centennial time scales. *International Journal of Climatology*, 20: 721-732.
- Rodríguez, A.; Montávez, J.P.; Roldán, C. & Jiménez, J.I. (1996).** Análisis y contraste de las series climatológicas de Granada y Sierra Nevada. *1ª Conferencia Internacional Sierra Nevada*, 20-26 de marzo de 1996. pp. 71-85.
- Rogerson, M.; Rohling, E.J.; Weaver, P.P. & Murray, J.W. (2004).** The Azores Front since the Last Glacial Maximum. *Earth and Planetary Science Letters* 222: 779-789.
- Röhl, U. & L. Abrams (2000).** High resolution, downhole, and nondestructive core measurements from Sites 999 and 1001 in the Caribbean Sea: Application to the late Paleocene thermal maximum, *Proc. Ocean Drill. Program, Sci. Results*, 165: 191-204, doi:10.2973/odp.proc.sr.165.009.2000.
- Rohling, E. J.; Abu-Zied, R. H.; Casford, J. S. L.; Hayes, A. & Hoogakker B. A. (2008).** The Mediterranean Sea: present and past. In: Woodward, J. C. (ed.) *The Physical Geography of the Mediterranean*. Oxford, UK, Oxford University Press.

- Royo Gómez, J. (1922).** El Mioceno continental ibérico y su fauna malacológica. *Com. Inv. Paleontol. Prehistóricas* 30: 227 pp.
- Rute, L. (1889).** *La Sierra Nevada*. Nouvelle Revue Internationale. Imprimerie Charaire et fils. Paris.
- Sánchez Gómez, S. (1990).** *Aplicación del estudio de suelos a la dinámica de la cuenca del río Lanjarón. Relación suelos-geomorfología*. Tesis doctoral. Universitat de Granada.
- Sánchez, S.; Simón, M., García, I. & Gómez Ortiz, A. (1998).** Morfogénesis de un sistema nival de Sierra Nevada: Laguna Seca (Provincia de Granada). *Cuaternario y Geomorfología*, 2: 99-105.
- Sanz De Galdeano, C. & López-Garrido, A.C. (1999).** Nature and impact of the Neotectonic deformation in the western Sierra Nevada (Spain). *Geomorphology*, 30: 259-272.
- Scheffer, F. & Schachtschabel, P. (2002).** *Lehrbuch der Bodenkunde*. Spektrum Akademischer Verlag.
- Schulte, L. (2002a).** Climatic and human influence on river systems and glacier fluctuations in southeast Spain. *Quaternary International* 93-94: 85-100.
- Schulte, L. (2002b).** *Evolución cuaternaria de la depresión de Vera y de Sorbas oriental (SE Península Ibérica). Reconstrucción de las fluctuaciones paleoclimáticas a partir de estudios morfológicos y edafológicos*. Servei de Publicacions de la Universitat de Barcelona. Barcelona.
- Schulte, L. (2002c).** Aplicación del método de datación del Pb<sup>210</sup> a la determinación cronológica de procesos geomórficos. In: Pérez-González, A.; Vegas, J. & Machado, J. (eds). *Aportaciones de la geomorfología de España en el inicio del tercer milenio*. pp. 483-490.
- Schulte, L.; Gómez Ortiz & Passmore, D.G. (2000).** Sobre la posible existencia de restos periglaciares pleistocenos en los llanos del sureste de la Península Ibérica. In: Peña Monné, Sánchez Fabre & Lozano (eds). *Procesos y formas periglaciares en la montaña mediterránea*. Instituto de Estudios Turolenses. pp. 281-297.
- Schulte, L.; Marcos García-Blanco, J.de; Gómez Ortiz, A.; Palacios-Estremera, D.; Tanarro García, L.M.; Fernández Fernández, A. & Ramos Sainz, M. (2002).** Evolución glacial y periglacial del Circo del Mulhacén (Sierra Nevada, Península Ibérica). In: Pérez-González, Vegas & Machado (eds): *Aportaciones a la Geomorfología de España en el inicio del tercer milenio*. pp. 491-499.
- Schulte, L.; Julià, R.; Oliva, M.; Burjachs, F.; Veit H. & Carvalho, F. (2008).** Sensitivity of alpine fluvial environments in the Swiss Alps to climate forcing during the Late Holocene. *Sediment Dynamics in Changing Environments*. Proceedings of a symposium held in Christchurch, New Zealand, December 2008). IAHS Publ. 325.
- Schulte, L.; Veit, H.; Burjachs, F.; Julià, R. (2009).** Lütschine fan delta response to climate variability and land use in the Bernese Alps during the last 2400 years. *Geomorphology*.
- Schwertmann, U. (1964).** Differenzierung der Eisenoxide des Bodens durch Extraktion mit saurer Ammoniumoxalat-Lösung. *Z. Pflanzenernähr., Düng., Bodenkunde*, 105: 194-202.
- Schwertmann, U. & Fitzpatrick, R.W. (1992).** Iron minerals in surface environments. In: Skinner, H.C.W. and Fitzpatrick, R.W. (editors). *Biomineralization Processes of Iron and Manganese* Catena Verlag, Cremlingen-Destedt, Germany, pp. 7-30.
- Seppälä, M. (1993).** Solifluction in northern Finland: past and present. In: Frenzel, B., Matthews, J.A., Gläser, B. (eds). *Solifluction and Climatic Variation in the Holocene*. Paläoklimaforschung, Vol. 11: 59-70.

- Shindell, T.; Schmidt, G.A.; Mann, M.; Rind, D. & Waple, A. (2001).** Solar forcing of regional climate change during the Maunder Minimum. *Science*, 294: 2149-2152.
- Simon, M.; Sánchez, S. & Garca, I. (2000).** Soil-landscape evolution on a Mediterranean high mountain. *Catena*, 39: 211-231.
- Smith, D. J. (1987).** Late Holocene solifluction in the Mount Rae area, southern Canadian Rocky Mountains. *Canadian Journal of Earth Sciences*, 24: 1634–1642.
- Smith, M.W. (1975).** Microclimatic influences on ground temperatures and permafrost distribution, Mackenzie Delta, Northwest Territories. *Canadian Journal of Earth Sciences*, 12: 1421-1438.
- Smol, J.P.; Birks, H.J.B. & Last, W.M. (eds) (2001).** Tracking Environmental Changes Using Lake Sediments. Kluwer Academic Publishers, Dordrecht. Vol. 3 i 4.
- Solanski, S.K; Usoskin I.G.; Kromer, B.; Schüssler, M. & Beer, J. (2004).** Unusual activity of the Sun during recent decades compared to the previous 11.000 anys. *Nature*, 431: 1084-1087.
- Soon, W. & Baliunas, S. (2003).** Proxy climatic and environmental changes of the past 1000 years. *Climate Research*, 23: 89-110.
- Sowers, T.; Alley, R.B. & Jubenville, J. (2003).** Ice Core Records of Atmospheric N<sub>2</sub>O Covering the Last 106,000 Years. *Science*, 301: 945-948.
- Steinmann, S. (1978).** Postglaziale Reliefgeschichte und gegenwärtige Vegetationsdifferenzierung in der alpinen Stufe der Südtiroler Dolomiten (Puez- und Sellagruppe). *Landschaftsgenese und Landschaftsökologie* 2:1–93.
- Stuiver, M., Braziunas, T.F., Grootes, P.M. & Zielinski, G.A. (1997).** Is There Evidence for Solar Forcing of Climate in the GISP2 Oxygen Isotope Record? *Quaternary Research* 48 (3): 259-266.
- Thorn, C.E. (1988).** *An Introduction to Theoretical Geomorphology*. Unwin Hyman, Boston. 247 pp.
- Titos Martínez, M. (1995).** *El tranvía de Sierra Nevada*. Málaga. Ed. Arguval.
- Titos Martínez, M. (1997).** *Sierra Nevada: una gran historia*. Universidad de Granada- CETURSA Sierra Nevada, S.A. vol. 2. Granada.
- Ulfstedt A.C. (1993).** Solifluction in the Swedish mountains: distribution in relation to vegetation and snow cover. In: Frenzel, B., Matthews, J.A., Gläser, B. (eds). *Solifluction and Climatic Variation in the Holocene*. Paläoklimaforschung, Vol. 11: 217-224.
- Usoskin, I.G.; Mursula, K.; Solanski, S.; Schüssler, M. i Alanko, K. (2004).** Reconstruction of solar activity for the last millenium using 10Be data. *Astronomy&Astrophysics*, 413: 745-751.
- Veit, H. (1988).** Fluviale und solifluidale Morphodynamik des Spät- und Postglazials in einem zentralalpinen Flusseinzugsgebiet (südliche Hohe Tauern, Osttirol). *Bayreuther Geowissenschaftliche Arbeiten*, Band 13: 1-167, Bayreuth.
- Veit, H. (1993).** Holocene solifluction in the Austrian and southern Tyrolean Alps: dating and climatic implications. In: Frenzel, B., Matthews, J.A., Gläser, B. (eds). *Solifluction and Climatic Variation in the Holocene*. Paläoklimaforschung, Vol. 11: 23-32.
- Veit, H. (2002).** Relict soils as paleoclimatic indicators: examples from the Austrian Alps and the Central Andes. *17th WCSS*, 14-21 Agost 2002 (Tailândia).
- Veit, H. & Höfner, T. (1993).** Permafrost, gelifluction and fluvial sediment transfer in the Alpine/Subnival ecotone, Central Alps, Austria: present, past and future. *Z. Geomorph. N.F.*, 92: 71-84.

- Vieira, G.; Mora, C. & Ramos, M. (2003).** Ground temperature regimes and geomorphological implications in a Mediterranean mountain (Serra da Estrela, Portugal), *Geomorphology* 52: 57-72.
- Vliet-Lanoe, B. (1993).** Solifluction in the North Atlantic Arctic and Subarctic: the microstratigraphic record of changing climate during the second half of the Holocene. In: Frenzel, B., Matthews, J.A., Gläser, B. (eds). *Solifluction and Climatic Variation in the Holocene*. Paläoklimaforschung, Vol. 11: 279-294.
- Wahrhaftig, C. & Cox, A. (1959).** Rock glaciers in the Alaska Range, *Geological Society of America Bulletin*, 70: 383–436.
- Wanner, H.; Holzhauser, H.; Pfister, C. & Zumbühl, H. (2000).** Interannual to century scale climate variability in the European Alps. *Erdkunde*, 54: 62-69.
- Wanner, H.; Beck, C.; Brádzil, R.; Casty, C.; Deutsch, M.; Glaser, R.; Jacobeit, J.; Luterbacher, J.; Pfister, C.; Pohl, S.; Sturm, K.; Werner, P.C. & Xoplaki, E. (2004).** Dynamic and socioeconomic aspects of historical floods in central Europe. *Erdkunde*, 58: 1-16.
- Wanner, H.; Beer, J.; Crowley, T.; Cubasch, U.; Flückiger, J.; Goosse, H.; Grosjean, M.; Kaplan, J.; Küttel, M.; Solomina, O.; Stocker, T.; Tarasov, P.; Wagner, M. & M. Widmann, M. (in press).** Mid to late Holocene climate change - a comprehensive review, *Quaternary Science Reviews*.
- Washburn, A. (1979).** *Geocryology: a Survey of Periglacial Processes and Environments*. Arnold, London. 406 pp.
- Wiersma, A.P. & Renssen, H. (2006).** Model–data comparison for the 8.2 ka BP event: confirmation of a forcing mechanism by catastrophic drainage of Laurentide Lakes. *Quaternary Science Reviews* 25: 63–88.
- Winiger, L. & Marti, T. (2006).** *Holocene solifluction activity in northern Iceland*. Diplomarbeit. Universität de Berna, 160 pp.
- Wu, H.; Guiot, J.; Brewer, S. & Guo, Z. (2007).** Climatic changes in Eurasia and Africa at the Last Glacial Maximum and mid-Holocene: reconstruction from pollen data using inverse vegetation modelling. *Climate Dynamics* 29: 211–229.
- Yu, Z.C. & Wright, H.E. (2001).** Response of interior North America to abrupt climate oscillations in the North Atlantic region during the last deglaciation. *Earth-Science Reviews*, 52: 333-369.
- Zorita, E., Von Storch, H., Gonzalez-Rouco, F.J., Cubasch, U., Luterbacher, J., Legutke, S., Fisher-Bruns, I. & Schlese, U. (2004).** Climate evolution in the last five centuries simulated by an atmospheric-ocean model: global temperatures, the North Atlantic Oscillation and the late Maunder Minimum. *Meteorologische Zeitschrift*, 13 (4): 271-289.

