

## **6.- BIBLIOGRAFÍA**



**Ali Azam, T. y Ishihama, A.** (1999). Twelve species of the nucleoid-associated protein from *Escherichia coli*. Sequence recognition specificity and DNA binding affinity. *The Journal of Biological Chemistry*. **274**: 33105-33113.

**Ali Azam, T., Iwata, A., Nishimura, A., Ueda, S. y Ishihama, A.** (1999). Growth phase-dependent variation in protein composition of the *Escherichia coli* nucleoid. *Journal of Bacteriology*. **181**: 6361-6370.

**Ali, B.M., Amit, R., Braslavsky, I., Oppenheim, A.B., Gileadi, O. y Stavans, J.** (2001). Compaction of single DNA molecules induced by binding of integration host factor (IHF). *Proc. Natl. Acad. Sci. USA*. **98**: 10658-10663.

**Amster-Choder, O., Houtman, F. y Wright, A.** (1989). Protein phosphorylation regulates transcription of the  $\beta$ -glucoside utilization operon in *E. coli*. *Cell*. **58**: 847-855.

**Amster-Choder, O. y Wright, A.** (1990). Regulation of activity of a transcriptional anti-terminator in *E. coli* by phosphorylation. *Science*. **249**: 540-542.

**Atlung, T. y Ingmer, H.** (1997). H-NS: a modulator of environmentally regulated gene expression. *Molecular Microbiology*. **24**: 7-17.

**Babu, M.M. y Teichmann, S.A.** (2003). Evolution of transcription factors and the gene regulatory network in *Escherichia coli*. *Nucleic Acids Research*. **31**: 1234-1244.

**Bailey, M.J.A., Koronakis, V., Schmoll, T. y Hughes, C.** (1992). *Escherichia coli* HlyT protein, a transcriptional activator of haemolysin synthesis and secretion, is encoded by the *rfaH* (*sfrB*) locus required for expression of sex factor and lipopolysaccharide genes. *Molecular Microbiology*. **6**: 1003-1012.

**Balsalobre, C., Johansson, J., Uhlin, B.E., Juárez, A. y Muñoz, F.J.** (1999). Alterations in protein expression caused by the *hha* mutation in *Escherichia coli*: influence of growth medium osmolarity. *Journal of Bacteriology*. **181**: 3018-3024.

**Balsalobre, C., Juárez, A., Madrid, C., Mouriño, M., Prenafreta, A. y Muñoa, F.J.** (1996). Complementation of the *hha* mutation in *Escherichia coli* by the *ymoA* gene from *Yersinia enterocolitica*: dependence on the gene dosage. *Microbiology*. **142**: 1841-1846.

**Berlyn, M.K.B., Low, K.B. y Rudd, K.E.** (1996). Integrated linkage map of *Escherichia coli*. A: F. C. Neidhardt *et al.*, ed., *Escherichia coli* and *Salmonella*: Cellular and molecular biology, 2nd ed., vol. 2. Washington DC: ASM Press.

**Bertin, P., Benhabiles, N., Krin, E., Laurent-Winter, C., Tendeng, C., Turlin, E., Thomas, A., Danchin, A. y Brasseur, R.** (1999). The structural and functional organization of H-NS-like proteins is evolutionarily conserved in Gram-negative bacteria. *Molecular Microbiology*. **31**: 319-329.

**Bertin, P., Hommais, F., Krin, E., Soutourina, O., Tendeng, C., Derzelle, S. y Danchin, A.** (2001). H-NS and H-NS-like proteins in Gram-negative bacteria and their multiple role in the regulation of bacterial metabolism. *Biochimie*. **83**: 235-241.

**Birnboim, H.C.** (1983). A rapid alkaline extraction method for the isolation of plasmid DNA. *Methods in Enzymology*. **100**: 243-254.

**Bloch, V., Yang, Y., Margeat, E., Chavanieu, A., Augé, M.T., Robert, B., Arold, S., Rimsky, S. y Kochoyan, M.** (2003). The H-NS dimerization domain defines a new fold contributing to DNA recognition. *Nature Structural Biology*. **10**: 212-218.

**Blomfield, I.C., Vaughn, V., Rest, R.F. y Eisenstein, B.I.** (1991). Allelic exchange in *Escherichia coli* using the *Bacillus subtilis* *sacB* gene and a temperature-sensitive pSC101 replicon. *Molecular Microbiology*. **5**: 1447-1457.

**Bolívar, F., Rodríguez, R.L., Greene, P.J., Betlach, M.C., Heyneker, H.L. y Boyer, H.W.** (1977). Construction and characterization of new cloning vehicles. II. A multipurpose cloning system. *Gene*. **2**: 95-113.

**Bracco, L., Kotlarz, D., Kolb, A., Diekmann, S. y Buc, H.** (1989). Synthetic curved DNA sequences can act as a transcriptional activators in *Escherichia coli*. *EMBO Journal*. **8**: 4289-4296.

**Bradford, M.M.** (1976). A rapid and sensitive method for the quantitation of micrograms quantities of proteins utilizing the principle of protein-dye binding. *Analytical Biochemistry*. **72**: 248-254.

**Brandi, A., Spurio, R., Gualerzi, C.O. y Pon, C.L.** (1999). Massive presence of the *Escherichia coli* "major cold-shock protein" CspA under non-stress conditions. *EMBO Journal*. **18**: 1653-1659.

**Braun, V., Schonherr, R. y Hobbie, S.** (1993). Enterobacterial haemolysins: activation, secretion and pore formation. *Trends in Microbiology*. **1**: 211-216.

**Brown, E.D., Vivas, E.I., Walsh, C.T. y Kolter, R.** (1995). *MurA* (*MurZ*), the enzyme that catalyzes the first committed step in peptidoglycan biosynthesis, is essential in *Escherichia coli*. *Journal of Bacteriology*. **177**: 4194-4197.

**Browning, D.F. y Busby, S.J.W.** (2004). The regulation of bacterial transcription initiation. *Nature Reviews*. **2**: 1-9.

**Burland, V., Shao, Y., Perna, N.T., Plunkett, G., Sofia, H.J. y Blattner, F.R.** (1998). The complete DNA sequence and analysis of the large virulence plasmid of *Escherichia coli* O157:H7. *Nucleic Acids Research*. **26**: 4196-4204.

**Burnette, W.N.** (1981). "Western blotting": Electrophoretic transfer of proteins from sodium dodecyl sulfate-polyacrilamide gels to unmodified nitrocellulose and radiographic detection with antibody and radioiodinated protein A. *Analytical Biochemistry*. **112**: 195-203.

**Caramel, A. y Schnetz K.** (1998). Lac and lambda repressors relieve silencing of the *Escherichia coli* *bgl* promoter. Activation by alteration of a repressing nucleoprotein complex. *Journal of Molecular Biology*. **284**: 875-883

**Carmona, M., Balsalobre, C., Muñoa, F.J., Mouriño, M., Jubete, Y., De la Cruz, F. y Juárez, A.** (1993). *Escherichia coli hha* mutants, DNA supercoiling and expression of haemolysin genes from recombinant plasmid pANN202-312. *Molecular Microbiology*. **9**: 1011-1018.

**Carniel, E., Mazigh, D. y Mollaret, H.H.** (1987). Expression of iron-regulated proteins in *Yersinia* species and their relation to virulence. *Infection immunology*. **55**: 277-280.

**Chang, A.C.Y. y Cohen, S.N.** (1978). Construction and characterization of amplifiable multicopy DNA cloning vehicles derived from the P15A cryptic miniplasmid. *Journal of Bacteriology*. **134**: 1141-1156.

**Cohen, S.N., Chang, A.C.Y. y Hsu, L.** (1972). Nonchromosomal antibiotic resistance in bacteria: genetic transformation of *Escherichia coli* by R-factor DNA. *Proceedings of the National Academy of Sciences of the United States of America*. **69**: 2110-2114.

**Coker, C., Bakare, O.O. y Mobley, H.L.** (2000). H-NS is a repressor of the *Proteus mirabilis* urease transcriptional activator gene *ureR*. *Journal of Bacteriology*. **182**: 2649-2653.

**Conchas, R.F. y Carniel, E.** (1990). A highly efficient electroporation system for transformation of *Yersinia*. *Gene*. **87**: 133-137.

**Cornelis, G.R.** (2002). *Yersinia* type III secretion: send in the effectors. *The Journal of Cell Biology*. **158**: 401-408.

**Cornelis, G.R., Boland, A., Boyd, A.P., Geuijen, C., Iriarte, M., Neyt, C., Sory, M.P. y Stainier, I.** (1998). The virulence plasmid of *Yersinia*, an antihost genome. *Microbiology and Molecular Biology Reviews*. **62**: 1315-1352.

**Cornelis, G.R. y Colson, C.** (1975). Restriction of DNA in *Yersinia enterocolitica* detected by recipient ability for a derepressed R factor from *Escherichia coli*. *Journal of General Microbiology*. **87**: 285-291.

**Cornelis, G., Sluiter, C., Delor, I., Gelb, D., Kanninga, K., Lambert de Rouvroit, C., Sory, M.P., Vanooteghem, J.C. y Michaelis T.** (1991). *YmoA*, a *Yersinia enterocolitica* chromosomal gene modulating the expression of virulence functions. *Molecular Microbiology*. **5**: 1023-1034.

**Cornelis, G., Sluiter, C., Lambert de Rouvroit, C. y Michiels, T.** (1989). Homology between VirF, the transcriptional activator of the *Yersinia* virulence regulon, and AraC, the *Escherichia coli* arabinose operon regulator. *Journal of Bacteriology*. **171**: 254-282.

**Cornelis, G., Vanooteghem, J.C. y Sluiter, C.** (1987). Transcription of the *yop* regulon from *Y. enterocolitica* requires trans acting pYV and chromosomal genes. *Microbial Pathogenesis*. **2**: 367-379.

**Csonka, L.N. y Hanson A.D.** (1991). Prokaryotic osmoregulation: genetics and physiology. *Annual Reviews of Microbiology*. **45**: 569-606.

**Cunningham, T.P., Montelaro, R.C. y Rushlow, K.E.** (1993). Lentivirus envelope sequences and proviral genomes are stabilized in *Escherichia coli* when cloned in low-copy-number plasmid vector. *Gene*. **9**: 93-98.

**Cusick, M.E. y Belfort, M.** (1998). Domain structure and RNA annealing activity of the *Escherichia coli* regulatory protein StpA. *Molecular Microbiology*. **28**: 847-857.

**Dame, R.T. y Goosen, N.** (2002). HU: promoting or counteracting DNA compaction? *FEBS Letters*. **529**: 151-156.

**Dame, R.T., Wyman, C., Wurm, R., Wagner, R. y Gossen, N.** (2002). Structural basis for H-NS mediated trapping of RNA polymerase in the open initiation complex at the *rrnB* P1. *Journal Biological Chemistry*. **277**: 2146-2150.

**Datsenko K.A. y Wanner, B.L.** (2000). One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proceedings of the National Academy of Sciences*. **97**: 6640-6645.

**Datta, N. y Hedges R.W.** (1972). Trimethoprim resistance conferred by W plasmids in *Enterobacteriaceae*. *Journal of General Microbiology*. **72**: 349-355.

**DeBord, K.L., Galanopoulos, N.S. y Schneewind, O.** (2003). The *ttsA* gene is required for low-calcium-induced type III secretion Yop proteins and virulence of *Yersinia enterocolitica* W22703. *Journal of Bacteriology*. **185**: 3499-3507.

**Deighan, P., Beloin, C. y Dorman, C.J.** (2003). Three-way interactions among the Sfh, StpA and H-NS nucleoid-associated proteins of *Shigella flexneri* 2a strain 2457T. *Molecular Microbiology*. **48**: 1401-1416.

**De la Cruz, F., Carmona, M. y Juárez, A.** (1992). The Hha protein from *Escherichia coli* is highly homologous to the YmoA protein from *Yersinia enterocolitica*. *Molecular Microbiology*. **6**: 3451-3454.

**De Lorenzo, V., Herrero, M., Jakubzik, U. y Timmis, K.N.** (1990). Mini-Tn5 transposon derivatives for insertion mutagenesis, promoter probing, and chromosomal insertion of cloned DNA in Gram-negative Eubacteria. *Journal of Bacteriology*. **172**: 6568-6572.

**Dersch, P., Schmidt, K. y Bremer, E.** (1993). Synthesis of the *Escherichia coli* K-12 nucleoid-associated DNA-binding protein H-NS is subjected to growth-phase control and autoregulation. *Molecular Microbiolgy*. **8**: 875-889.

**Dole, S., Klingen, Y., Nagarajavel, V. y Schnetz, K.** (2004a). The protease Lon and the RNA-Binding protein Hfq reduce silencing of the *Escherichia coli bgl* operon by H-NS. *Journal of Bacteriology*. **186**: 2708-2716.

**Dole, S., Nagarajavel, V. y Schnetz, K.** (2004). The histone-like nucleoid structuring protein H-NS represses the *Escherichia coli bgl* operon downstream of the promoter. *Molecular Microbiology*. **52**: 589-600.

**Dorman, C.J. y Deighan, P.** (2003). Regulation of gene expression by histone-like proteins in bacteria. *Current Opinion*. **13**: 179-184.



**Dorman, C.J., Hinton, J.C.D. y Free, A.** (1999). Domain organization and oligomerization among H-NS-like nucleoid-associated proteins in bacteria. *Trends in Microbiology*. **7**: 124-128.

**Dorman, C.J.** (2004). H-NS: A universal regulator for a dynamic genome. *Nature reviews (Microbiology)*. **2**: 391-400.

**Dove, S.L., Darst, S.A. y Hochschild, A.** (2003). Region 4 of  $\sigma$  as a target for transcription regulation. *Molecular Microbiology*. **48**: 863-874.

**Dower, W.J., Miller, F.J. y Ragsdale, C.W.** (1988). High efficiency transformation of *E. coli* by high voltage electroporation. *Nucleic Acids Research*. **16**: 6127-6145.

**Drlica, K.** (1992). Control of bacterial DNA supercoiling. *Molecular Microbiology*. **6**: 425-33.

**Ebright, R.H.** (2000). RNA polymerase: structural similarities between bacterial RNA polymerase and eukaryotic RNA polymerase II. *Journal of Molecular Biology*. **304**: 687-698.

**Ebright, R.H. y Busby, S.** (1995). The *Escherichia coli* RNA polymerase  $\alpha$  subunit: structure and function. *Current Opinion in Genetics and Development*. **5**: 197-203.

**Edwards, R.A., Linda, H.K. y Schifferli, D.M.** (1998). Improved allelic exchange vectors and their use to analyze 987P fimbria gene expression. *Gene*. **207**: 149-157.

**Eisenstadt, E., Carlton, B.C. y Brown, B.J.** (1994). Gene Mutation. A: *Methods for General and Molecular Bacteriology*, págs. 297-315. Gerhardt, P., Murray, R.G.E., Wood, W.A. y Krieg, N.R. American Society for Microbiology, Washington, D.C.

**Elliot, S.J., Sperandio, V., Giron, J.A., Shin, S., Mellies, J.L., Wainwright, L., Hutcheson, S.W., McDaniel, T.K. y Kaper, J.B.** (2000). The locus of enterocyte effacement (LEE)- encoded regulator controls expression of both LEE- and no LEE-

encoded virulence factors in enteropathogenic and enterohemorrhagic *Escherichia coli*. *Infection and Immunity*. **68**: 6115-6126.

**Ellison, D.W., Young, B., Nelson, K. y Miller, V.L.** (2003). YmoA negatively regulates expression of invasins from *Yersinia enterocolitica*. *Journal of Bacteriology*. **185**: 7153-7159.

**Esposito, D., Petrovic, A., Harris, R., Ono, S., Eccleston, J.F., Mbabaali, A., Haq, I., Higgins, C.F., Hinton, J.C., Driscoll, P.C., y Ladbury, J.E.** (2002). H-NS oligomerization domain structure reveals the mechanism for high order self-association of the intact protein. *Journal of Molecular Biology*. **324**: 841-850.

**Fahlen, T.F., Mathur, N. y Jones, B.D.** (2000). Identification and characterization of mutants with increased expression of *hilA*, the invasion gene transcriptional activator of *Salmonella typhimurium*. *FEMS Immunology Medicine Microbiology*. **28**: 25-35.

**Fahlen, T.F., Wilson, R.B., Boddicker, J.D. y Jones, B.D.** (2001). Hha is a negative modulator of transcription of *hilA*, the *Salmonella enterica* serovar *Thyphimurium* invasion gene transcriptional activator. *Journal of Bacteriology*. **183**: 6620-6629.

**Falconi, M., Brandi, A., La Teana, A., Gualerzi, C.O. y Pon C.L.** (1996). Antagonistic involvement of FIS and H-NS proteins in transcriptional control of *hns* expression. *Molecular Microbiology*. **19**: 965-975.

**Falconi, M., Colonna, B., Prosseda, G., Micheli, G. y Gualerzi, C.O.** (1998). Thermoregulation of *Shigella* and *Escherichia coli* EIEC pathogenicity. A temperature-dependent structural transition of DNA modulates accessibility of *virF* promoter to transcriptional repressor H-NS. *EMBO Journal*. **17**: 7033-7043.

**Falconi, M., Higgins, N.P., Spurio, R., Pon, C.L. y Gualerzi, C.O.** (1993). Expression of the gene encoding the major bacterial nucleoid protein H-NS is subject to transcriptional auto-repression. *Molecular Microbiology*. **10**: 273-282.

**Falconi, M., McGovern, V., Gualerzi, G., Hillyard, D. y Higgins, N.P.** (1991). Mutations altering chromosomal protein H-NS induce mini-Mu transposition. *New Biologist*. **3**: 615-625.

**Falconi, M., Prosseda, G., Giangrossi, M., Beghetto, E. y Colonna, B.** (2001). Involvement of FIS in the H-NS-mediated regulation of *virF* gene of *Shigella* and enteroinvasive *Escherichia coli*. *Molecular Microbiology*. **42**: 439-452.

**Finkel, S.A. y Johnson, R.C.** (1992). The Fis protein: it's not just for DNA inversion anymore. *Molecular Microbiology*. **6**: 3257-3265.

**Forns, Núria.** (2003). Termoregulación de la conjugación del plásmido R27 de *Salmonella typhi*: papel de las proteínas asociadas al nucleóide Hha y H-NS. *Memoria para el diploma de estudios avanzados*.

**Forns, N., Baños, R.C., Balsalobre, C., Juárez, A. y Madrid, C.** (2005). Temperature-dependent conjugative transfer of R27: role of chromosomal- and plasmid-encoded Hha and H-NS proteins. *Journal of Bacteriology*, *In press*.

**Free, A. y Dorman, C.J.** (1997). The *Escherichia coli stpA* gene is transiently expressed during growth in rich medium and is induced in minimal medium and by stress conditions. *Journal of Bacteriology*. **179**: 909-918.

**Free, A., Williams, R.M. y Dorman, C.J.** (1998). The StpA protein functions as a molecular adapter to mediate repression of the *bgl* operon by truncated H-NS in *Escherichia coli*. *Journal of Bacteriology*. **180**: 994-997.

**Freundlich, M., Ramani, N., Mathew, E., Sirko, A. y Tsui, P.** (1992). The role of integration host factor in gene expression in *Escherichia coli*. *Molecular Microbiology*. **6**: 2557-2563.

**Friedrich, K., Gualerzi, C.O., Lammi, M., Losso, M.A. y Pon, C.L.** (1988). Proteins from the prokaryotic nucleoid. Interaction of nucleic acids with the 15 kDa *Escherichia coli* histone-like protein H-NS. *FEBS Letters*. **229**: 197-202.

**Gabant P., Szpirer C.Y., Couturier M. y Faelen M.** (1998). Direct selection cloning vectors adapted to the genetic analysis of gram-negative bacteria and their plasmids. *Gene*. **207**: 87-92.

**García, J., Cordeiro, T.N., Nieto, J.M., Pons, I., Juárez, A. y Pons, M.** (2005). Interaction between the bacterial nucleoid associated proteins Hha and H-NS involves a conformational change of Hha. *Biochemical Journal* (Publicación on-line).

**Glaser, P., Ladant, D., Selzer, D., Pichot, F., Ullmann, A. y Daunchin, A.** (1988). The calmodulin-sensitive adenylate-cyclase of *Bordetella pertussis*: cloning and expression in *Escherichia coli*. *Molecular Microbiology*. **2**: 19-30.

**Godessart, N., Muñoa, F.J., Regué, M. y Juárez, A.** (1988). Chromosomal mutations that increase the production of a plasmid-encoded haemolysin in *Escherichia coli*. *Journal of General Microbiology*. **134**: 2779-2787.

**Goebel, W. y Schrempf, H.** (1971). Isolation and characterization of supercoiled circular deoxyribonucleic acid from beta-haemolytic strains of *Escherichia coli*. *Journal of Bacteriology*. **106**: 311-317.

**Goldberg, M.D., Johnson, M., Hilton, J.C.D. y Williams, P.H.** (2001). Role of the nucleoid-associated protein Fis in the regulation of virulence properties of enteropathogenic *Escherichia coli*. *Molecular Microbiology*. **41**: 549-559.

**González-Gil, G., Bringmann, P. y Kahmann, R.** (1996). FIS is a regulator of metabolism in *Escherichia coli*. *Molecular Microbiology*. **22**: 21-29.

**Goosen, N y van de Putte, P.** (1995). The regulation of transcription initiation by integration host factor. *Molecular Microbiology*. **16**: 1-7.

**Göransson, M., Sonden, B., Nilsson, P., Dagberg, B., Forsman, K., Emanuelsson, K. y Uhlin, B.E.** (1990). Transcriptional silencing and thermoregulation of gene expression in *Escherichia coli*. *Nature*. **344**: 682-685.

**Gottesman, S.** (1984). Bacterial regulation: global regulatory networks. *Annual Reviews Genetics*. **18**: 415-441.

**Goyard, S. y Bertin, P.** (1997). Characterization of BpH3 and H-NS-like protein in *Bordetella pertussis*. *Molecular Microbiology*. **24**: 815-823.

**Grindley, N.D.F., Grindley, J.N. y Anderson, E.S.** (1972). R factor compatibility groups. *Molecular General Genetics*. **119**: 287-297.

**Gruber, T.M. y Gross, C.A.** (2003). Multiple sigma subunits and the partitioning of bacterial transcription space. *Annual Reviews in Microbiology*. **57**: 441-466.

**Gulati, A. y Mahadevan, S.** (2000). Mechanism of catabolite repression in the *bgl* operon of *Escherichia coli*: involvement of the anti-terminator BglG, CRP-cAMP and EIIA<sup>Glc</sup> in mediating glucose effect downstream of transcription initiation. *Gene Cells*. **5**: 239-250.

**Hamilton, C.M., Aldea, M., Washburn, B.K., Babitzke, P. y Kushner, S.R.** (1989). New method for generating deletions and gene replacements in *Escherichia coli*. *Journal of Bacteriology*. **171**: 4617-4622.

**Hanahan, D., Jessee, J. y Bloom, F.R.** (1991). Plasmid transformation of *Escherichia coli* and other bacteria. *Methods in Enzymology*. **204**: 63-113.

**Hardie, K.R., Issartel, J.P., Koronakis, E., Hughes, C. y Koronakis, V.** (1991). In vitro activation of *Escherichia coli* prohaemolysin to the mature membrane-targeted toxin requires HlyC and a low molecular-weight cytosolic polypeptid. *Molecular Microbiology*. **5**: 1669-1679.

**Hatfield, G.W. y Benham, C.J.** (2002). DNA topology-mediated control of global gene expression in *Escherichia coli*. *Annual Review in Genetics*. **36**: 175-203.

**Heroven, A.K., Nagel, G., Tran, H.J., Parr, S. y Dersch, P.** (2004). RovA is autoregulated and antagonizes H-NS-mediated silencing of invasion and *rovA* expression in *Yersinia pseudotuberculosis*. *Molecular Microbiology*. **53**: 871-888.

**Herrero, M., de Lorenzo, V. y Timmis, K.N.** (1990). Transposon vectors containing non-antibiotic resistance selection markers for cloning and stable chromosomal insertion of foreign genes in gram-negative bacteria. *Journal Bacteriology*. **172**: 6557-6567.

**Hess, J., Wels, W., Vogel, M. y Goebel, W.** (1986). Nucleotide sequence of a plasmid-encoded haemolysin determinant and its comparison with a corresponding chromosomal haemolysin sequence. *FEMS Microbiology Letters*. **34**: 1-11.

**Higgins, C.F., Dorman, C.J., Stirling, D.A., Waddel, L., Booth, I.R., May, G. y Bremer, E.** (1988). A physiological role for DNA supercoiling in the osmotic regulation of gene expression in *S. typhimurium* and *E. coli*. *Cell*. **52**: 569-584.

**Hochschild, A. y Dove, S.L.** (1998). Protein-protein contacts that activate and repress prokaryotic transcription. *Cell*. **92**: 597-600.

**Hochuli, E., Bannwarth, W., Dobeli, H., Gentz, R. y Stüber, D.** (1988). Genetic approach to facilitate purification of recombinant proteins with a novel metal chelate absorbent. *Bio/Technology*. **6**: 1321-1325.

**Holland, I.B., Mackman, N. y Nicaud, J.M.** (1986). Secretion of proteins from bacteria. *Biotechnology*. **4**: 427-431.

**Hommais, F., Krin, E., Laurent-Winter, C., Soutourina, O., Malpertuy, A., Le Caer, J.P., Danchin, A. y Bertin, P.** (2001). Large-scale monitoring of pleiotropic regulation of gene expression by the prokaryotic nucleoid-associated protein, H-NS. *Molecular Microbiology*. **40**: 20-36.

**Houman, F., Díaz-Torres, M.R. y Wright, A.** (1990). Transcriptional antitermination in the *bgl* operon of *E. coli* is modulated by a specific RNA binding protein. *Cell*. **62**: 1153-63.

**Hsieh, L.S., Burger, R.M. y Drlica, K.** (1991a). Bacterial DNA supercoiling and [ATP]/[ADP] ratio: changes associated with a transition to anaerobic growth. *Journal of Molecular Biology*. **219**: 443-450.

**Hsieh, L.S., Rouvière-Yaniv, J. y Drlica, K.** (1991b). Bacterial DNA supercoiling and [ATP]/[ADP] ratio: changes associated with salt shock. *Journal of Bacteriology*. **173**: 3914-3917.

**Hughes, K. y Mathee, K.** (1998). The anti-sigma factors. *Annual Reviews in Microbiology*. **52**: 231-286.

**Hulton, C.S.J., Seirafi, A., Hinton, J.C.D., Sidebotham, J.M., Waddell, L., Pavitt, G.D., Owen-Hughes, T., Spassky, A., Buc, H. y Higgins, C.F.** (1990). Histone-like protein H1 (H-NS), DNA supercoiling, and gene expression in bacteria. *Cell*. **63**: 631-642.

**Ishihama, A.** (2000). Functional modulation of *Escherichia coli* RNA polymerase. *Annual Reviews in Microbiology*. **54**: 499-518.

**Issartel, J.P., Koronakis, V. y Hughes, C.** (1991). Activation of *E.coli* prohaemolysin to the mature toxin by acyl carrier protein-dependent fatty acylation. *Nature*. **351**: 759-761.

**Jarvis, K.G., Giron, J.A., Jerse, A.E., McDaniel, T.K., Domnenberg, M.S. y Kaper, J.B.** (1995). Enteropathogenic *Escherichia coli* contains a putative type III secretion system necessary for the export of proteins involved in attaching and effacing lesion formation. *Proc. Natl. Acad. Sci. USA*. **92**: 7996-8000.

**Jayaraman, K., Müller-Hill, B. y Rickenberg, H.** (1966). Inhibition of the synthesis of  $\beta$ -Galactosidase in *Escherichia coli* by 2-Nitrophenyl- $\beta$ -D-Fucoside. *Journal of Molecular Biology*. **18**: 339-343.

**Johansson, J., Balsalobre, C., Wang, S.Y., Urbonaviciene, J., Jin, D.J., Sondén, B. y Uhlin, B.E.** (2000). Nucleoid proteins stimulate stringently controlled bacterial promoters: A link between the cAMP-CRP and the (p)ppGpp regulons in *Escherichia coli*. *Cell*. **102**: 475-485.

**Johansson, J., Dagberg, B., Richet, E. y Uhlin, B.E.** (1998). H-NS and StpA proteins simulate expression of the maltose regulon in *Escherichia coli*. *Journal of Bacteriology*. **180**: 6117-6125.

**Johansson, J., Eriksson, S., Sondén, B., Wai, S.N. y Uhlin, B.E.** (2001). Heteromeric interactions among nucleoid-associated bacterial proteins: Localization of StpA-stabilizing regions in H-NS of *Escherichia coli*. *Journal of Bacteriology*. **183**: 2343-2347.

**Johansson, J. y Uhlin, B.E.** (1999). Differential protease-mediated turnover of H-NS and StpA revealed by a mutation altering protein stability and stationary-phase survival of *Escherichia coli*. *Proceedings of the National Academy of Sciences, USA*. **96**: 10776-10781.

**Juárez, A., Hughes, C., Vogel, M. y Goebel, W.** (1984). Expression and regulation of the plasmid-encoded haemolysin determinant of *Escherichia coli*. *Molecular and General Genetics*. **197**: 196-203.

**Jubete, Y., Zabala, J.C., Juárez, A. y De la Cruz, F.** (1995). *hlyM*, a transcriptional silencer downstream of the promoter in the *hly* operon of *Escherichia coli*. *Journal of Bacteriology*. **177**: 242-246.

**Kar, S. y Adhya, S.** (2001). Recruitment of HU by piggyback: a special role of GalR in repressosome assembly. *Genes and Development*. **15**: 2273-2281.

**Karunakaran, P., Endresen, D.T., Ertesvag, H., Blatny, J.M. y Valla, S.** (1999). A small derivative of the broad-host-range plasmid RK2 which can be switched from a replicating to a non-replicating state as a response to an externally added inducer. *FEMS Microbiology Letters*. **180**: 221-227.

**Koronakis, V., Cross, M. y Hughes, C.** (1988). Expression of the *Escherichia coli* haemolysin secretion gene *hlyB* involves transcript anti-terminator within the *hly* operon. *Nucleic Acids Research*. **16**: 4789-4800.

**Laemmli, U.K.** (1970). Cleavage of structural proteins during the assembly of head of the bacteriophage T4. *Nature*. **227 (259)**: 680-685.



**Lambert de Rouvroit, C., Sluifers, C. y Cornelis, G.R.** (1992). Role of the transcriptional activator, VirF, and temperature in the expression of the pYV plasmid genes of *Yersinia enterocolitica*. *Molecular Microbiology*. **6**: 395-409.

**Lejeune, P., Bertin, P., Walon, C., Willemot, K., Colson, C. y Danchin, A.** (1989). A locus involved in kanamycin, chloramphenicol and L-serine resistance is located in the *bglY-galU* region of the *Escherichia coli* K12 chromosome. *Molecular General Genetic*. **218**: 361-363.

**Lejeune, P. y Danchin, A.** (1990). Mutations in the *bglY* gene increase the frequency of spontaneous deletions in *Escherichia coli* K12. *Proceedings of the Natural Academy of Sciences, USA*. **87**: 360-363.

**Link, A.J., Philips, D. y Church, G.M.** (1997). Methods for generating precise deletions and insertions in the genome of wild type *Escherichia coli*: applications to open reading frame characterization. *Journal of Bacteriology*. **179**: 6228-6237.

**Liu, Q. y Richardson, C.** (1993). Gene 5.5 protein of bacteriophage T7 inhibits the nucleoid protein H-NS of *Escherichia coli*. *Proceedings of the Natural Academy of Sciences, USA*. **90**: 1761-1765.

**Lupas, A.** (1996). Coiled coils: new structures and new functions. *Trends in Biochemistry Science*. **21**: 375-382.

**Madrid, C., Nieto, J.M. y Juárez, A.** (2002b). Role of the Hha/YmoA family of proteins in the thermoregulation of the expression of virulence factors. *International Journal of Medical Microbiology*. **291**: 425-432.

**Madrid, C., Nieto, J.M., Paytubi, S., Falconi, F., Gualerzi, C.O. y Juárez, A.** (2002a). Temperature- and H-NS-Dependent regulation of a plasmid-encoded virulence operon expressing *Escherichia coli* haemolysin. *Journal of Bacteriology*. **184**: 5058-5066.

**Mahadevan, S., Reynolds, A.E. y Wright, A.** (1987). Positive and negative regulation of the *bgl* operon of *Escherichia coli*. *Journal of Bacteriology*. **169**: 2570-2578.

**Mahan, M.J., Slauch, J.M. y Mekalanos, J.** (1996). Environmental regulation of virulence gene expression in *Escherichia*, *Salmonella*, and *Shigella* spp. A.: Neidhardt F.C. *et al.*, ed., *Escherichia coli* and *Salmonella: Cellular and molecular biology*. **Pág:** 2803-2815. American Society for Microbiology, Washington.

**Maher, D. y Taylor, D.E.** (1993). Host range and transfer efficiency of incompatibility group HI plasmid. *Canadian Journal of Microbiology*. **39:** 581-587.

**Marsh, P.** (1986). p<sub>tac85</sub>, an *Escherichia coli* vector for expression of non-fusion proteins. *Nucleic Acids Research*. **14:** 3603.

**Marshal, D. G., Sheenan, B. J. y Dorman, C. J.** (1999). A role for the leucine-responsive regulator protein and integration host factor in the regulation of the *Salmonella* plasmid virulence (*spv*) locus in *Salmonella typhimurium*. *Molecular Microbiology*. **34:** 134-145.

**Martínez-Antonio, A. y Collado-Vides, J.** (2003). Identifying global regulators in transcriptional regulatory networks in bacteria. *Current Opinion in Microbiology*. **6:** 482-489.

**Martínez, E. y de la Cruz, F.** (1988). Transposon Tn2 *I* encodes a *recA*-independent site-specific integration system. *Molecular General Genetics*. **211:** 320-325.

**Maurelli, A.T. y Sansonetti, P.J.** (1988). Identification of a chromosomal gene controlling temperature-regulated expression of *Shigella* virulence. *Proceedings of the natural academy of sciences of United States of America*. **85:** 2820-2824.

**McLeod, S.M. y Johnson, R.C.** (2001). Control of transcription by nucleoid proteins. *Current Opinion in Microbiology*. **4:** 152-159.

**Menestrina, G., Moser, C., Pellet, S. y Welch, R.** (1994). Pore-formation by *Escherichia coli* haemolysin (HlyA) and other members of the RTX toxins family. *Toxicology*. **87:** 249-267.

**Michiels, T., Vanooteghem, J.C., Lambert de Rouvroit, C., China, B., Gustin, A., Boudry, P. y Cornelis, G.R.** (1991), Analysis of *virC*, an operon involved in the secretion of Yop proteins by *Yersinia enterocolitica*. *Journal of Bacteriology*. **173**: 4994-5009.

**Mikulskis, A.V. y Cornelis, G.** (1994). A new class of proteins regulating gene expression in enterobacteria. *Molecular Microbiology*. **11**: 77-86.

**Miller, J.H.** (1992). *A Short Course in Bacterial Genetics. A Laboratory Manual and Handbook for Escherichia coli and Related Bacteria*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N. Y.

**Miller, V. y Mekalanos, J.** (1988). A novel suicide vector and its use in construction of insertion mutations: osmoregulation of outer membrane proteins and virulence determinants in *Vibrio cholerae* requires *toxR*. *Journal Bacteriology*. **170 (6)**: 2575-2583.

**Mouriño, M., Balsalobre, C., Madrid, C., Nieto, J.M., Prenafreta, A., Muñoa, F.J. y Juárez, A.** (1998). Osmolarity modulates the expression of the Hha protein from *Escherichia coli*. *FEMS Microbiology Letters*. **160**: 225-229.

**Mouriño, M., Madrid, C., Balsalobre, C., Prenafreta, A., Muñoa, F.J., Blanco, J., Blanco, M., Blanco, J.E. y Juárez, A.** (1996). The Hha protein as a modulator of expression of virulence factors in *Escherichia coli*. *Infection and Immunity*. **64**: 2881-2884.

**Mouriño, M., Muñoa, F.J., Balsalobre, C., Díaz, P., Madrid, C. y Juárez, A.** (1994). Environmental regulation of the  $\alpha$ -haemolysin expression in *Escherichia coli*. *Microbial Pathogenesis*. **16**: 249-259.

**Muffler, A., Fischer, D., Altuvia, S., Storz, G. y Hengge-Aronis, R.** (1996). The RNA-binding protein HF-I, known as a host factor for phage Q $\beta$  RNA replication, is essential for *rpoS* translation in *Escherichia coli*. *Genes and Development*. **10**: 1143-1151.

**Mukerji, M. y Mahadevan, S.** (1997). Characterization of the negative elements involved in silencing the *bgl* operon of *Escherichia coli*: possible roles for DNA gyrase, H-NS and CRP-cAMP in regulation. *Molecular Microbiology*. **24**: 617-627.

**Mullis K., Faloona F., Scharf S., Saiki R., Horn G. y Erlich H.** (1986). Specific enzymatic amplification of DNA in vitro: the polymerase chain reaction. *Cold Spring Harb Symp Quant Biol.* **51** Pt **1**:263-73

**Neidhardt, F. C. y Savageau, M.A.** (1996). Regulation beyond the operon. Neidhardt, F.C., Curtis III, R., Ingraham, J.L., Lin, E.C.C., Low, K.B., Magasanik, B., Reznikoff, W.S., Riley, M., Schaechter, M. y Umbarger, H.E. (Eds.). *A Escherichia coli and Salmonella: Cellular and molecular biology*. **Pág:** 1310-1324. American Society for Microbiology, Washington.

**Nieto, J.M. y Juárez, A.** (1996). A new member of the Hha/YmoA class of bacterial regulators in plasmid R100 of *Escherichia coli*? *Molecular Microbiology.* **19**: 407.

**Nieto, J.M. y Juárez, A.** (1999). The putative Orf4 of Broad-Host-Range conjugative plasmid R446 could be related to the H-NS family of bacterial nucleoid-associated proteins. *Plasmid.* **41**: 125-127.

**Nieto, J.M., Bailey, M.J.A., Hughes, C. y Koronakis, V.** (1996). Suppression of transcription polarity in the *Escherichia coli* haemolysin operon by a short upstream element shared by polysaccharide and DNA transfer determinants. *Molecular Microbiology.* **19**: 705-713.

**Nieto, J.M., Carmona, M., Bolland, S., Jubete, Y., De la Cruz, F. y Juárez, A.** (1991). The *hha* gene modulates haemolysin expression in *Escherichia coli*. *Molecular Microbiology.* **5**: 1285-1293.

**Nieto, J.M., Madrid, C., Miquelay, E., Parra, J.L., Rodríguez, S. y Juárez, A.** (2002). Evidence for direct protein-protein interaction between members of enterobacterial Hha/YmoA and H-NS families of proteins. *Journal of Bacteriology.* **184**: 629-635.

**Nieto, J.M., Madrid, C., Prenafeta, A., Miquelay, E., Balsalobre, C., Carrascal, M. y Juárez, A.** (2000). Expression of the haemolysin operon in *Escherichia coli* is modulated by a nucleoid-protein complex that includes proteins Hha and H-NS. *Molecular General Genetics.* **263**: 349-358.

**Nieto, J.M., Prenafeta, A., Miquelay, E., Torrades, S. y Juárez, A.** (1998). Sequence, identification and effect on conjugation of the *rmoA* gene of plasmid R100-1. *FEMS Microbiology Letters*. **169**: 59-66.

**Noegel, A., Rdest, U. y Goebel, W.** (1981). Determinations of the functions of hemolytic plasmid pHly152 of *Escherichia coli*. *Journal of Bacteriology*. **145**: 233-247.

**Nye, M.B., Pfau, J.D., Skorupski, K. y Taylor, R.K.** (2000). *Vibrio cholerae* H-NS silences virulence gene expression at multiple steps in the *ToxR* regulatory cascade. *Journal of Bacteriology*. **182**: 4295-4303.

**O'Hanley, P., Marcus, R., Baek, K.Y., Denich, K. y Ji, G.E.** (1993). Genetic conservation of *hly* determinants and serological conservation of HlyA: Basis for developing a broadly cross-reactive subunit *Escherichia coli* alpha-haemolysin vaccine. *Infect and Immunity*. **61**: 1091-1097.

**Oberto, J. y Rouvière-Yaniv, J.** (1996). *Serratia marcescens* contains a heterodimeric HU protein like *Escherichia coli* and *Salmonella typhimurium*. *Journal of Bacteriology*. **178**: 293-297.

**Oshima, T., Wada, C., Kawagoe, Y., Ara, T., Maeda, M., Masuda, Y., Hiraga, S. y Mori, H.** (2002). Genome-wide analysis of deoxyadenosine methyltransferase-mediated control of gene expression in *Escherichia coli*. *Molecular microbiology*. **45**: 673-695.

**Owen-Hughes, T.A., Pavitt, G.D., Santos, D.S., Sidebotham, J.M., Hulton, C.S.J., Hinton, J.C.D. y Higgins, C.F.** (1992). The chromatin-associated protein H-NS interacts with curved DNA to influence DNA topology and gene expression. *Cell*. **71**: 255-265.

**Packman, L.C.** (1993). Protein chemical methods for molecular biologists. *Methods in Molecular and Cellular Biology*. **4**: 189-198.

**Paytubi, S., Madrid, C., Forns, N., Nieto, J.M., Balsalobre, C., Uhlin, B.E. y Juárez, A.** (2004). YdgT, the Hha paralogue in *Escherichia coli*, forms heteromeric complexes with H-NS and StpA. *Molecular Microbiology*. **54**: 251-263.

**Pérez-Rueda, E. y Collado-Vides, J.** (2000). The repertoire of DNA-binding transcriptional regulators in *Escherichia coli* K-12. *Nucleic Acids Research*. **28**: 1838-1847.

**Porter, M.E. y Dorman, C.J.** (1997). Differential regulation of the plasmid-encoded genes in the *Shigella flexneri* virulence regulon. *Molecular and General Genetics*. **256**: 93-103.

**Reusch, R.N., Shabalin, O., Crumbaugh, A., Wagner, R., Schroder, O. y Wurm, R.** (2002). Posttranslational modification of *E. coli* histone-like protein H-NS and bovine histones by short-chain poly-(R)-3-hydroxybutyrate (cPHB). *FEBS Letters*. **527**: 319-322.

**Rimsky, S.** (2004). Structure of the histone-like protein H-NS and its role in regulation and genome superstructure. *Current Opinion in Microbiology*. **7**: 109-114.

**Rimsky, S., Zuber, F., Buckle, M. y Buch. H.** (2001). A molecular mechanism for the repression of transcription by the H-NS protein. *Molecular Microbiology*. **42**: 1311-1323.

**Rodríguez, R.L. y Tait, R.C.** (1983). Recombinant DNA Techniques: An introduction. *The Benjamin/Cummings Publishing Co., Inc., Menlo Park, California*. **Págs**: 187-191.

**Rojo, F.** (1999). Repression of transcription initiation in bacteria. *Journal of Bacteriology*. **181**: 2987-2991.

**Rubirés, X., Saigi, F., Piqué, N., Climent, N., Merino, S., Albertí, S., Tomás, J.M. y Regué, M.** (1997). A gene (*wbbL*) from *Serratia marcescens* N28b (O4) complements the *rfb-50* mutation of *Escherichia coli* K-12 derivatives. *Journal of Bacteriology*. **179**: 7581-7586.

**Sanger, F. y Coulson A.R.** (1975). A rapid method for determining sequences in DNA by primed synthesis with DNA polymerase. *Journal of Molecular Biology*. **94(3)**: 441-448.

**Sambrook, J., Fritsch, E.F. y Maniatis, T.** (1989). *Molecular cloning: A Laboratory Manual*. 2ª Ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.

**Sarkar, G. y Sommer, S.** (1990). The 'Megaprimer' method of site-directed mutagenesis. *BioTechniques*. **8**: 404-407.

**Schaechter, M.** (2001). *Escherichia coli* and *Salmonella* 2000: the view from here. *Microbiology and Molecular Biology Reviews*. **65**: 119-130.

**Schaeffler, S.** (1967). Inducible system for utilization of  $\beta$ -glucosides in *Escherichia coli*. *Journal of Bacteriology*. **93**: 254-263.

**Schägger, H. y von Jagow, G.** (1987). Tricine-sodium dodecyl sulfate-polyacrilamide gel electrophoresis for the separation of proteins in the range from 1 to 100 kDa. *Analytical Biochemistry*. **166**: 368-379.

**Schechter, L.M., Jain, S., Akbar, S. y Lee, C.A.** (2003). The small nucleoid-binding proteins H-NS, HU and Fis affects *hila* expression in *Salmonella enterica* serovar *Typhimurium*. *Infection and Immunity*. **71**: 5432-5435.

**Schneider, R., Travers, A., Kutateladze, T. y Muskhelishvili, G.** (1999). A DNA architectural protein couples cellular physiology and DNA topology in *Escherichia coli*. *Molecular Microbiology*. **34**: 953-964.

**Schnetz, K.** (1995). Silencing of *Escherichia coli bgl* promoter by flanking sequence elements. *EMBO Journal*. **14**: 2545-2550.

**Schnetz, K.** (2002). Silencing of the *Escherichia coli bgl* operon by RpoS requires Crl. *Microbiology*. **148**: 2573-2578.

**Schnetz, K. y Rak, B.** (1988). Regulation of the *bgl* operon of *Escherichia coli* by transcriptional antitermination. *EMBO Journal*. **7**: 3271-3277.

**Schnetz, K., Toloczyki, C. y Rak, B.** (1987). Beta-glucoside (*bgl*) operon of *Escherichia coli* K-12: nucleotide sequence genetic organization and possible evolutionary relationship to regulatory components of two *Bacillus subtilis* genes. *Journal of Bacteriology*. **169**: 2579-2590.

**Schröder, O. y Wagner, R.** (2002). The bacterial regulatory protein H-NS- A versatile modulator of nucleic acid structures. *Biological Chemistry*. **383**: 945-960.

**Sharma, V.K. y Zuerner, R.L.** (2004). Role of *hha* and *ler* in transcriptional regulation of the *esp* operon of enterohemorrhagic *Escherichia coli* O157:H7. *Journal of Bacteriology*. **186**: 7290-7301.

**Shen-Orr, S.S., Milo, R., Mangan, S. y Alon, U.** (2002). Network motifs in the transcriptional regulation network of *Escherichia coli*. *Nature Genetics*. **31**: 64–68.

**Sherburne, C.K., Lawley, T.D., Gilmour, M.W., Blattner, F.R., Burland, V., Grotbeck, E., Rose, D.J. y Taylor, D.E.** (2000). The complete DNA sequence and analysis of R27, a large IncHI plasmid from *Salmonella typhi* that is temperature sensitive for transfer. *Nucleic Acids Research*. **28**: 2177-2186.

**Sherrat, D.J.** (2003). Bacterial chromosome dynamics. *Science*. **301**: 780-785.

**Shi, X. y Bennett, G.N.** (1994). Plasmids bearing *hfq* and the *hns*-like gene *stpA* complement *hns* mutants in modulating *arginine decarboxilase* gene expression in *Escherichia coli*. *Journal of Bacteriology*. **176**: 6769-6775.

**Shindo, H., Iwaki, T., Ieda, R., Kurumizaka, H., Ueguchi, C., Mizuno, T., Morikawa, S., Nakamura, H. y Kuboniwa, H.** (1995). Solution structure of the DNA binding domain of a nucleoid-associated protein, H-NS, from *Escherichia coli*. *FEBS Letters*. **360**: 125-131.

**Sieber, V., Martinez, C.A. y Arnold, F.H.** (2001). Libraries of hybrid proteins from distantly related sequences. *Nature biotechnology*. **19**: 456-460.

**Silhavy, T.J., Berman, M.L. y Enquist, L.W.** (1984). Experiments with gene fusions. Cold Spring Harbor Laboratory, Cold Spring Harbor, N. Y.



**Simon, R., Hotte, B., Klauke, B. y Kosier, B.** (1991). Isolation and characterization of insertion sequence elements from gram-negative bacteria by using new broad-host-range, positive selection vectors. *Journal of Bacteriology*. **173**: 1502-1508.

**Simon, R., Priefer, U. y Puhler, A.** (1983). A broad host range mobilization system for in vivo genetic engineering: transposon mutagenesis in gram negative bacteria. *Bio/Technology*. **1**: 784-791.

**Skerra, A.** (1994). Use of the tetracycline promoter for the tightly regulated production of a murine antibody fragment in *Escherichia coli*. *Gene*. **151**: 131-135.

**Smith, H.** (1968). Biochemical challenge of microbial pathogenicity. *Bacteriological Reviews*. **32**: 164-184.

**Smyth, C.P., Lundbäck, T., Renzoni, D., Siligardi, G., Bevil, R., Layton, M., Sidebotham, J.M., Hinton, J.C.D., Driscoll, P.C., Higgins, C.F. y Ladbury, J.E.** (2000). Oligomerization of the chromatin-structuring protein H-NS. *Molecular Microbiology*. **36**: 962-972.

**Sondén, B. y Uhlin, B.E.** (1996). Coordinated and differential expression of histone-like proteins in *Escherichia coli*: regulation and function of the H-NS analog StpA. *EMBO Journal*. **15**: 4970-4980.

**Sonnenfield, J.M., Burns, C.M., Higgins, C.F. y Hinton, J.C.** (2001). The nucleoid-associated protein StpA binds curved DNA, has a greater DNA-binding affinity than H-NS and is present in significant levels in *hns* mutants. *Biochimie*. **83**: 243-249.

**Spassky, A., Rimsky, S., Garreau, H. y Buc. H.** (1984). H1a, an *E. coli* DNA-binding protein which accumulates in stationary phase, strongly compacts DNA *in vitro*. *Nucleic Acids Research*. **12**: 5321-5340.

**Spurio, R., Falconi, M., Brandi, A., Pon, C.L. y Gualerzi, C.O.** (1997). The oligomeric structure of nucleoid protein H-NS is necessary for recognition of intrinsically curved DNA and for DNA bending. *EMBO Journal*. **16**: 1795-1805.

**Strathdee, C.A. y Lo, R.Y.C.** (1987). Extensive homology between the leucotoxin of *Pasteurella hemolytica* A1 and the  $\alpha$ -haemolysin of *Escherichia coli*. *Infection and Immunity*. **155**: 3233-3236.

**Studier, F.W. y Moffatt, B.A.** (1986). Use of bacteriophage T7 RNA polymerase to direct selective high-level expression of cloned genes. *Journal of Molecular Biology*. **189**: 113-130.

**Studier, F.W., Rosengerg, A.H., Dunn, J.J. y Dubendorff, J.W.** (1990). Use of T7 RNA polymerase to direct expression of cloned genes. *Methods in Enzymology*. **185**: 60-89.

**Taylor, L.W. y Levine, J.G.** (1980). Studies of temperature-sensitive transfer and maintenance of H incompatibility group plasmids. *Journal of General Microbiology*. **116**: 475-484.

**Tendeng, C., Badaut, C., Krin, E., Gounon, P., Ngo, S., Danchin, A., Rimsky, S. y Bertin, P.** (2000). Isolation and characterization of *vicH*, encoding a new regulator in *Vibrio cholerae*. *Journal of Bacteriology*. **182**: 2026-2032.

**Tendeng, C. y Bertin, P.** (2003). H-NS in Gram-negative bacteria: a family of multifaceted proteins. *Trends in Microbiology*. **11**: 511-518.

**Thieffry, D., Huerta, A.M., Pérez-Rueda, E. y Collados-Vides, J.** (1998). From specific gene regulation to genomic networks: a global analysis of transcriptional regulation in *Escherichia coli*. *Bioassays*. **20**: 433-440.

**Tietze, E. y Tschäpe, H.** (1994). Temperature-dependent expression of conjugation pili by IncM plasmid-harboring bacteria: identification of plasmid-encoded regulatory functions. *Journal of Basic Microbiology*. **34**: 105-116.

**Tippner, D. y Wagner, R.** (1995). Fluorescence analysis of the *Escherichia coli* transcription regulator H-NS reveals two distinguishable complexes dependent on binding to specific or nonspecific DNA sites. *The Journal of Biological Chemistry*. **270**: 22243-22247.

**Trachman, J.D. y Maas, W.K.** (1998). Temperature regulation of heat-labile enterotoxin (LT) synthesis in *Escherichia coli* is mediated by an interaction of H-NS protein with the LT A-subunit DNA. *Journal of Bacteriology*. **180**: 3715-3718.

**Towbin, H., Staehelin, T. y Gordon, J.** (1979). Electrophoretic transfer of proteins from polyacrilamide gels to nitrocellulose sheets: Procedure and some applications. *Proceedings of the Natural Academy of Sciences of the United States of America*. **76**: 4350-4354.

**Tupper, A.E., Owen-Hughes, T.A., Ussery, D.W., Santos, D.S., Ferguson, D.J.P., Sidebotham, J.M., Hinton, J.C.D. y Higgins, C.F.** (1994). The chromatin-associated protein H-NS alters DNA topology in vitro. *EMBO Journal*. **13**: 258-268.

**Ueguchi, C., Kakeda, M. y Mizuno, T.** (1993). Autoregulatory expression of the *Escherichia coli hns* gene encoding a nucleoid protein: H-NS functions as a repressor of its own transcription. *Molecular and General Genetics*. **236**: 171-178.

**Ueguchi, C., Suzuki, T., Yoshida, T., Tanaka, K.I. y Mizuno, T.** (1996). Systematic mutational analysis revealing the functional domain organization of *Escherichia coli* nucleoid protein H-NS. *Journal of Molecular Biology*. **263**: 149-162.

**Ussery, D.W., Hinton, J.C.D., Jordi, B.J.A. M., Granum, P.E., Seirafi, A., Stephen, R.J., Tupper, A.E., Berridge, G., Sidebotham, J.M. y Higgins, C.D.** (1994). The chromatin associated protein H-NS. *Biochimie*. **76**: 968-980.

**Varshavsky, A.J., Nedospasov, S.A., Bakayev, V.V., Bakayeva, T.G. y Georgiev, G.P.** (1977). Histone-like proteins in the purified *Escherichia coli* deoxyribonucleo-protein. *Nucleic Acids Research*. **4**: 2725-2745.

**Venkatesan, M.M., Goldberg, M.B., Rose, D.J., Grotbeck, E.J., Burland, V. y Blattner, F.R.** (2001). Complete DNA sequence and analysis of the large virulence plasmid of *Shigella flexnerii*. *Infection and Immunity*. **69**: 3271-3285.

**Vogel, M. Hess, J., Then, I, Juárez, A. y Goebel, W.** (1988). Characterization of a sequence (*hlyR*) which enhances synthesis and secretion of haemolysin in *Escherichia coli*. *Molecular and General Genetics*. **212**: 76-84.

**Wagner, R.** (2000). Lessons from a manifold regulated system. *Transcription Regulation in Prokaryotes*. Oxford: Oxford University Press. **Pág.:** 309-330.

**Wandersman, C. y Delepelaire, P.** (1990). TolC, an *Escherichia coli* outer membrane protein required for haemolysin secretion. *Proceedings of the Natural Academy of Sciences, USA*. **87**: 4776-4780.

**Warren, J.W., Walker, J.R., Roth, J.R. y Altman, E.** (2000). Construction and characterization of a highly regulable expression vector, pLAC11, and its multipurpose derivatives, pLAC22 and pLAC33. *Plasmid*. **44**: 138-151.

**Way, J.C., Davis, M.A., Morisato, D., Roberts, D.E. y Kleckner, N.** (1984). New Tn10 derivatives for transposon mutagenesis and for the construction of lacZ operon fusions by transposition. *Gene*. **32**: 369-379.

**Welch, R.A., Dellinger, E.P., Minshew, B. y Falkow, S.** (1981). Haemolysin contributes to virulence of extra-intestinal *E. coli* infections. *Nature*. **294**: 665-667.

**Welch, R.A. y Pellett, S.** (1988). Transcriptional organization of the *Escherichia coli* haemolysin genes. *Journal of Bacteriology*. **170**: 1622-1630.

**Williams, R.M. y Rimsky, S.** (1997). Molecular aspects of the *E.coli* nucleoid protein H-NS: a central controller of gene regulatory networks. *FEMS Microbiology Letters*. **156**: 175-185.

**Williams, R.M., Rimsky, S. y Buc, H.** (1996). Probing the structure, function, and interactions of the *Escherichia coli* H-NS and StpA proteins by using dominant negative derivatives. *Journal of Bacteriology*. **178**: 4335-4343.

**Williamson, H.S. y Free, A.** (2005). A truncated H-NS-like protein from enteropathogenic *Escherichia coli* acts as an H-NS antagonist. *Molecular Microbiology*. **55**: 808-827.

**Wilson, R.L., Libby, S.J., Freet, A.M., Boddicker, J.D., Fahlen, T.F. y Jones, B.D.** (2001). Fis, a DNA nucleoid-associated protein, is involved in *Salmonella typhimurium* SPI-1 invasion gene expression. *Molecular Microbiology*. **39**: 79-88.

**Wilson, J.W., Schurr, M.J., LeBlanc, C.L., Ramamurthy, R., Buchanan, K.L. y Nickerson, C.A.** (2002). Mechanisms of bacterial pathogenicity. *Postgraduate Medical Journal*. **78**: 216-224.

**Wösten, M.M.S.M.** (1998). Eubacterial sigma-factors. *FEMS Microbiology Reviews*. **22**: 127-150.

**Xu, J. y Johnson, R.C.** (1995). Identification of genes negatively regulated by Fis: Fis and RpoS comodulate growth-phase dependent gene expression in *Escherichia coli*. *Journal of Bacteriology*. **177**: 938-947.

**Yamada, H., Muramatsu, S. y Sasakawa, C.** (1990). An *Escherichia coli* protein that preferentially binds to sharply curved DNA. *Journal of Biochemistry*. **108**: 420-425.

**Yanisch-Perron, C., Vieira, J. y Messing, M.** (1985). Improved M13 phage cloning vectors and host strains: nucleotide sequencing of the M13mp18 and pUC19 vectors. *Gene*. **33**: 103-119.

**Yee, A., Chang, X., Pineda-Lucena, A., Wu, B., Semesi, A., Le, B., Ramelot, T., Lee, G.M., Bhattacharyya, S., Gutierrez, P., Denisov, A., Lee, C.H., Cort, J.R., Kozlov, G., Liao, J., Finak, G., Chen, L., Wishart, D., Lee, W., McIntosh, L.P., Gehring, K., Kennedy, M.A., Edwards, A.M. y Arroxwmith, C.H.** (2002). An NMR approach to structural proteomics. *Proceedings of the Natural Academy of Sciences, USA*. **99**: 1825-30.

**Yoshioka, Y., Ohtsubo, H. y Ohtsuvo, E.** (1987). Repressor gene *finO* in plasmids R100 and F: constitutive transfer of plasmid F is caused by insertion of IS3 into *F finO*. *Journal of Bacteriology*. **169**: 619-623.

**Yother, J., Chamness, T.W. y Goguen, J.** (1986). Temperature-controlled plasmid regulon associated with low calcium response in *Yersinia pestis*. *Journal of Bacteriology*. **165**: 704-711.

**Zhang, A. y Belfort, M.** (1992). Nucleotide sequence of a newly-identified *Escherichia coli* gene, *stpA*, encoding an H-NS-like protein. *Nucleic Acids Research*. **20**: 67.

**Zhang, A., Derbyshare, V., Galloway Salvo, J.L. y Belfort, M.** (1995). *Escherichia coli* protein StpA stimulates self-splicing by promoting RNA assembly *in vitro*. *RNA*. **1**: 783-793.

**Zhang, A., Rimsky, S., Reaban, M.E., Buc, H. y Belfort, M.** (1996). *Escherichia coli* protein analogs StpA and H-NS: regulatory loops, similar and disparate effects on nucleic acid dynamics. *EMBO Journal*. **15**: 1340-1349.