



Paper del canal de clorur CIC-2 en les patologies de la mielina

Tanit Arnedo Llena

ADVERTIMENT. La consulta d'aquesta tesi queda condicionada a l'acceptació de les següents condicions d'ús: La difusió d'aquesta tesi per mitjà del servei TDX (www.tdx.cat) i a través del Dipòsit Digital de la UB (diposit.ub.edu) ha estat autoritzada pels titulars dels drets de propietat intel·lectual únicament per a usos privats emmarcats en activitats d'investigació i docència. No s'autoritza la seva reproducció amb finalitats de lucre ni la seva difusió i posada a disposició des d'un lloc aliè al servei TDX ni al Dipòsit Digital de la UB. No s'autoritza la presentació del seu contingut en una finestra o marc aliè a TDX o al Dipòsit Digital de la UB (framing). Aquesta reserva de drets afecta tant al resum de presentació de la tesi com als seus continguts. En la utilització o cita de parts de la tesi és obligat indicar el nom de la persona autora.

ADVERTENCIA. La consulta de esta tesis queda condicionada a la aceptación de las siguientes condiciones de uso: La difusión de esta tesis por medio del servicio TDR (www.tdx.cat) y a través del Repositorio Digital de la UB (diposit.ub.edu) ha sido autorizada por los titulares de los derechos de propiedad intelectual únicamente para usos privados enmarcados en actividades de investigación y docencia. No se autoriza su reproducción con finalidades de lucro ni su difusión y puesta a disposición desde un sitio ajeno al servicio TDR o al Repositorio Digital de la UB. No se autoriza la presentación de su contenido en una ventana o marco ajeno a TDR o al Repositorio Digital de la UB (framing). Esta reserva de derechos afecta tanto al resumen de presentación de la tesis como a sus contenidos. En la utilización o cita de partes de la tesis es obligado indicar el nombre de la persona autora.

WARNING. On having consulted this thesis you're accepting the following use conditions: Spreading this thesis by the TDX (www.tdx.cat) service and by the UB Digital Repository (diposit.ub.edu) has been authorized by the titular of the intellectual property rights only for private uses placed in investigation and teaching activities. Reproduction with lucrative aims is not authorized nor its spreading and availability from a site foreign to the TDX service or to the UB Digital Repository. Introducing its content in a window or frame foreign to the TDX service or to the UB Digital Repository is not authorized (framing). Those rights affect to the presentation summary of the thesis as well as to its contents. In the using or citation of parts of the thesis it's obliged to indicate the name of the author.

Paper del canal de clorur CIC-2 en les patologies de la mielina.

Tesi doctoral: Tanit Arnedo Llana
Departament de Ciències Fisiològiques II,
Unitat de Fisiologia
Universitat de Barcelona 2015

Director: Raúl Estévez Povedano
Departament de Ciències Fisiològiques II, UB

BIBLIOGRAFIA

Abascal, F., and Zardoya, R. (2012). LRRC8 proteins share a common ancestor with pannexins, and may form hexameric channels involved in cell-cell communication. *Bioessays* 34, 551–560.

Abrams, C.K., and Scherer, S.S. (2012). Gap junctions in inherited human disorders of the central nervous system. *Biochim. Biophys. Acta* 1818, 2030–2047.

Accardi, A., and Pusch, M. (2000). Fast and slow gating relaxations in the muscle chloride channel CLC-1. *J. Gen. Physiol.* 116, 433–444.

Ahmed, N., Ramjeesingh, M., Wong, S., Varga, A., Garami, E., and Bear, C.E. (2000). Chloride channel activity of CLC-2 is modified by the actin cytoskeleton. *Biochem. J.* 352 Pt 3, 789–794.

Altevogt, B.M., and Paul, D.L. (2004). Four classes of intercellular channels between glial cells in the CNS. *J. Neurosci.* 24, 4313–4323.

Alvarez-Maubecin, V., Garcia-Hernandez, F., Williams, J.T., and Van Bockstaele, E.J. (2000). Functional coupling between neurons and glia. *J. Neurosci.* 20, 4091–4098.

Ambrosini, E., Serafini, B., Lanciotti, A., Tosini, F., Scialpi, F., Psaila, R., Raggi, C., Di Girolamo, F., Petrucci, T.C., and Aloisi, F. (2008). Biochemical characterization of MLC1 protein in astrocytes and its association with the dystrophin-glycoprotein complex. *Mol. Cell. Neurosci.* 37, 480–493.

Amiry-Moghaddam, M., and Ottersen, O.P. (2003). The molecular basis of water transport in the brain. *Nat. Rev. Neurosci.* 4, 991–1001.

Araque, A., Parpura, V., Sanzgiri, R.P., and Haydon, P.G. (1998). Glutamate-dependent astrocyte modulation of synaptic transmission between cultured hippocampal neurons. *Eur. J. Neurosci.* 10, 2129–2142.

Arnedo, T., Aiello, C., Jeworutzki, E., Dentici, M.L., Uziel, G., Simonati, A., Pusch, M., Bertini, E., and Estévez, R. (2014a). Expanding the spectrum of megalencephalic leukoencephalopathy with subcortical cysts in two patients with GLIALCAM mutations. *Neurogenetics* 15, 41–48.

Arnedo, T., Tania, L., Jeworutzki, E., Capdevila-nortes, X., and Pusch, M. (2014b). Functional Analyses of Mutations in HEPACAM Causing Megalencephalic Leukoencephalopathy. *Hum. Mut.* 35, 1175–1178.

Arreola, J., Begenisich, T., and Melvin, J.E. (2002). Conformation-dependent regulation of inward rectifier chloride channel gating by extracellular protons. *J. Physiol.* 541, 103–112.

Ashcroft, F.M. (2006). From molecule to malady. *Nature* 440, 440–447.

Bachoo, R.M., Kim, R.S., Ligon, K.L., Maher, E.A., Brennan, C., Billings, N., Chan, S., Li, C., Rowitch, D.H., Wong, W.H., et al. (2004). Molecular diversity of astrocytes with implications for neurological disorders. *Proc. Natl. Acad. Sci. U. S. A.* 101, 8384–8389.

Bali, M., Lipecka, J., Edelman, A., and Fritsch, J. (2001). Regulation of CLC-2 chloride channels in T84 cells by TGF- α . *Am. J. Physiol. Cell Physiol.* 280, C1588–C1598.

Bateman, A. (1997). The structure of a domain common to archaebacteria and the homocystinuria disease protein. *Trends Biochem. Sci.* 22, 12–13.

Becker, D., Blase, C., Bereiter-Hahn, J., and Jendrach, M. (2005). TRPV4 exhibits a functional role in cell-volume regulation. *J. Cell Sci.* *118*, 2435–2440.

Becker, D., Bereiter-Hahn, J., and Jendrach, M. (2009). Functional interaction of the cation channel transient receptor potential vanilloid 4 (TRPV4) and actin in volume regulation. *Eur. J. Cell Biol.* *88*, 141–152.

Di Bella, D., Pareyson, D., Savoiaro, M., Farina, L., Ciano, C., Caldarazzo, S., Sagnelli, A., Bonato, S., Nava, S., Bresolin, N., et al. (2014). Subclinical leukodystrophy and infertility in a man with a novel homozygous CLCN2 mutation. *Neurology* *83*, 1217–1218.

Benarroch, E.E. (2005). Neuron-astrocyte interactions: partnership for normal function and disease in the central nervous system. *Mayo Clin. Proc.* *80*, 1326–1338.

Benesova, J., Rusnakova, V., Honsa, P., Pivonkova, H., Dzamba, D., Kubista, M., and Anderova, M. (2012). Distinct expression/function of potassium and chloride channels contributes to the diverse volume regulation in cortical astrocytes of GFAP/EGFP mice. *PLoS One* *7*, e29725.

Benfenati, V., and Ferroni, S. (2010). Water transport between CNS compartments: functional and molecular interactions between aquaporins and ion channels. *Neuroscience* *168*, 926–940.

Benfenati, V., Amiry-Moghaddam, M., Caprini, M., Mylonakou, M.N., Rapisarda, C., Ottersen, O.P., and Ferroni, S. (2007a). Expression and functional characterization of transient receptor potential vanilloid-related channel 4 (TRPV4) in rat cortical astrocytes. *Neuroscience* *148*, 876–892.

Benfenati, V., Nicchia, G.P., Svelto, M., Rapisarda, C., Frigeri, A., and Ferroni, S. (2007b). Functional down-regulation of volume-regulated anion channels in AQP4 knockdown cultured rat cortical astrocytes. *J. Neurochem.* *100*, 87–104.

Benfenati, V., Caprini, M., Dovizio, M., Mylonakou, M.N., Ferroni, S., Ottersen, O.P., and Amiry-Moghaddam, M. (2011). An aquaporin-4/transient receptor potential vanilloid 4 (AQP4/TRPV4) complex is essential for cell-volume control in astrocytes. *Proc. Natl. Acad. Sci. U. S. A.* *108*, 2563–2568.

Bernard, P., and Couturier, M. (1992). Cell killing by the F plasmid CcdB protein involves poisoning of DNA-topoisomerase II complexes. *J. Mol. Biol.* *226*, 735–745.

Blackburn, D., Sargsyan, S., Monk, P.N., and Shaw, P.J. (2009). Astrocyte function and role in motor neuron disease: a future therapeutic target? *Glia* *57*, 1251–1264.

Blanz, J., Schweizer, M., Auberson, M., Maier, H., Muenscher, A., Hübner, C.A., and Jentsch, T.J. (2007). Leukoencephalopathy upon disruption of the chloride channel ClC-2. *J. Neurosci.* *27*, 6581–6589.

Blattner, R., Von Moers, A., Leegwater, P.A.J., Hanefeld, F.A., Van der Knaap, M.S., and Köhler, W. (2003). Clinical and genetic heterogeneity in megalencephalic leukoencephalopathy with subcortical cysts (MLC). *Neuropediatrics* *34*, 215–218.

Boespflug-Tanguy, O., Labauge, P., Fogli, A., and Vaurs-Barriere, C. (2008). Genes involved in leukodystrophies: a glance at glial functions. *Curr. Neurol. Neurosci. Rep.* 8, 217–229.

Boor, I., Nagtegaal, M., Kamphorst, W., van der Valk, P., Pronk, J.C., van Horsen, J., Dinopoulos, A., Bove, K.E., Pascual-Castroviejo, I., Muntoni, F., et al. (2007). MLC1 is associated with the dystrophin-glycoprotein complex at astrocytic endfeet. *Acta Neuropathol.* 114, 403–410.

Boor, P.K.I., de Groot, K., Waisfisz, Q., Kamphorst, W., Oudejans, C.B.M., Powers, J.M., Pronk, J.C., Scheper, G.C., and van der Knaap, M.S. (2005). MLC1: a novel protein in distal astroglial processes. *J. Neuropathol. Exp. Neurol.* 64, 412–419.

Bösl, M.R., Stein, V., Hübner, C., Zdebik, A.A., Jordt, S.E., Mukhopadhyay, A.K., Davidoff, M.S., Holstein, A.F., and Jentsch, T.J. (2001). Male germ cells and photoreceptors, both dependent on close cell-cell interactions, degenerate upon CIC-2 Cl(-) channel disruption. *EMBO J.* 20, 1289–1299.

Bossi, E., Fabbrini, M.S., and Ceriotti, A. (2007). Exogenous protein expression in *Xenopus* oocytes: basic procedures. *Methods Mol. Biol.* 375, 107–131.

Bovolenta, P., Liem, R.K., and Mason, C.A. (1984). Development of cerebellar astroglia: transitions in form and cytoskeletal content. *Dev. Biol.* 102, 248–259.

Branicky, R., Miyazaki, H., Strange, K., and Schafer, W.R. (2014). The Voltage-Gated Anion Channels Encoded by *clh-3* Regulate Egg Laying in *C. elegans* by Modulating Motor Neuron Excitability. 34, 764–775.

Brignone, M.S., Lanciotti, A., Macioce, P., Macchia, G., Gaetani, M., Aloisi, F., Petrucci, T.C., and Ambrosini, E. (2011). The beta1 subunit of the Na,K-ATPase pump interacts with megalencephalic leukoencephalopathy with subcortical cysts protein 1 (MLC1) in brain astrocytes: new insights into MLC pathogenesis. *Hum. Mol. Genet.* 20, 90–103.

Brignone, M.S., Lanciotti, A., Visentin, S., De Nuccio, C., Molinari, P., Camerini, S., Diociaiuti, M., Petrini, S., Minnone, G., Crescenzi, M., et al. (2014). Megalencephalic leukoencephalopathy with subcortical cysts protein-1 modulates endosomal pH and protein trafficking in astrocytes: relevance to MLC disease pathogenesis. *Neurobiol. Dis.* 66, 1–18.

Brockmann, K., Finsterbusch, J., Terwey, B., Frahm, J., and Hanefeld, F. (2003). Megalencephalic leukoencephalopathy with subcortical cysts in an adult: quantitative proton MR spectroscopy and diffusion tensor MRI. *Neuroradiology* 45, 137–142.

Bruzzone, R., Hormuzdi, S.G., Barbe, M.T., Herb, A., and Monyer, H. (2003). Pannexins, a family of gap junction proteins expressed in brain. *Proc. Natl. Acad. Sci. U. S. A.* 100, 13644–13649.

Butt, A.M., and Kalsi, A. (2006). Inwardly rectifying potassium channels (Kir) in central nervous system glia: a special role for Kir4.1 in glial functions. *J. Cell. Mol. Med.* 10, 33–44.

Capdevila-Nortes, X., López-Hernández, T., Ciruela, F., and Estévez, R. (2012). A modification of the split-tobacco etch virus method for monitoring interactions between membrane proteins in mammalian cells. *Anal. Biochem.* 423, 109–118.

Capdevila-Nortes, X., López-Hernández, T., Apaja, P.M., López de Heredia, M., Sirisi, S., Callejo, G., Arnedo, T., Nunes, V., Lukacs, G.L., Gasull, X., et al. (2013). Insights into MLC pathogenesis: GlialCAM is an MLC1 chaperone required for proper activation of volume-regulated anion currents. *Hum. Mol. Genet.* 22, 4405–4416.

Chever, O., Djukic, B., McCarthy, K.D., and Amzica, F. (2010). Implication of Kir4.1 channel in excess potassium clearance: an in vivo study on anesthetized glial-conditional Kir4.1 knock-out mice. *J. Neurosci.* 30, 15769–15777.

Clark, S., Jordt, S.E., Jentsch, T.J., and Mathie, A. (1998). Characterization of the hyperpolarization-activated chloride current in dissociated rat sympathetic neurons. *J. Physiol.* 506 (Pt 3, 665–678.

Clayton, G.H., Owens, G.C., Wolff, J.S., and Smith, R.L. (1998). Ontogeny of cation-Cl⁻ cotransporter expression in rat neocortex. *Brain Res. Dev. Brain Res.* 109, 281–292.

Cornejo, I., Niemeyer, M.I., Zúñiga, L., Yusef, Y.R., Sepúlveda, F. V, and Cid, L.P. (2009). Rapid recycling of ClC-2 chloride channels between plasma membrane and endosomes: role of a tyrosine endocytosis motif in surface retrieval. *J. Cell. Physiol.* 221, 650–657.

Cortez, M.A., Li, C., Whitehead, S.N., Dhani, S.U., D'Antonio, C., Huan, L.J., Bennett, S.A.L., Snead, O.C., and Bear, C.E. (2010). Disruption of ClC-2 expression is associated with progressive neurodegeneration in aging mice. *Neuroscience* 167, 154–162.

Cuppoletti, J., Tewari, K.P., Sherry, A.M., Kupert, E.Y., and Malinowska, D.H. (2001). ClC-2 Cl⁻ channels in human lung epithelia: activation by arachidonic acid, amidation, and acid-activated omeprazole. *Am. J. Physiol. Cell Physiol.* 281, C46–C54.

Cuppoletti, J., Tewari, K.P., Sherry, A.M., Ferrante, C.J., and Malinowska, D.H. (2004). Sites of protein kinase A activation of the human ClC-2 Cl⁻ channel. *J. Biol. Chem.* 279, 21849–21856.

D'Agostino, D., Bertelli, M., Gallo, S., Cecchin, S., Albiero, E., Garofalo, P.G., Gambardella, A., St Hilaire, J.-M., Kwiecinski, H., Andermann, E., et al. (2004). Mutations and polymorphisms of the CLCN2 gene in idiopathic epilepsy. *Neurology* 63, 1500–1502.

D'Ambrosio, R., Gordon, D.S., and Winn, H.R. (2002). Differential role of KIR channel and Na⁽⁺⁾/K⁽⁺⁾-pump in the regulation of extracellular K⁽⁺⁾ in rat hippocampus. *J. Neurophysiol.* 87, 87–102.

Dani, J.W., Chernjavsky, A., and Smith, S.J. (1992). Neuronal activity triggers calcium waves in hippocampal astrocyte networks. *Neuron* 8, 429–440.

Depienne, C., Bugiani, M., Dupuits, C., Galanaud, D., Touitou, V., Postma, N., van Berkel, C., Polder, E., Tollard, E., Darios, F., et al. (2013). Brain white matter oedema due to ClC-2 chloride channel deficiency: an observational analytical study. *Lancet. Neurol.* 12, 659–668.

Dhani, S.U., Kim Chiaw, P., Huan, L.-J., and Bear, C.E. (2008). ATP depletion inhibits the endocytosis of ClC-2. *J. Cell. Physiol.* 214, 273–280.

Dibaj, P., Kaiser, M., Hirrlinger, J., Kirchhoff, F., and Neusch, C. (2007). Kir4.1 channels regulate swelling of astroglial processes in experimental spinal cord edema. *J. Neurochem.* *103*, 2620–2628.

Duan, D., Ye, L., Britton, F., Horowitz, B., and Hume, J.R. (2000). A novel anionic inward rectifier in native cardiac myocytes. *Circ. Res.* *86*, E63–E71.

Duarri, A., Teijido, O., López-Hernández, T., Scheper, G.C., Barriere, H., Boor, I., Aguado, F., Zorzano, A., Palacín, M., Martínez, A., et al. (2008). Molecular pathogenesis of megalencephalic leukoencephalopathy with subcortical cysts: mutations in MLC1 cause folding defects. *Hum. Mol. Genet.* *17*, 3728–3739.

Duarri, A., Lopez de Heredia, M., Capdevila-Nortes, X., Ridder, M.C., Montolio, M., López-Hernández, T., Boor, I., Lien, C.-F., Hagemann, T., Messing, A., et al. (2011). Knockdown of MLC1 in primary astrocytes causes cell vacuolation: a MLC disease cell model. *Neurobiol. Dis.* *43*, 228–238.

Dubey, M., Bugiani, M., Ridder, M.C., Postma, N.L., Brouwers, E., Polder, E., Jacobs, J.G., Baayen, J.C., Klooster, J., Kamermans, M., et al. (2014). Mice with Megalencephalic Leukoencephalopathy with Cysts: A developmental angle. *Ann. Neurol.*

Duffield, M., Rychkov, G., Bretag, A., and Roberts, M. (2003). Involvement of helices at the dimer interface in ClC-1 common gating. *J. Gen. Physiol.* *121*, 149–161.

Duffy, S., and MacVicar, B.A. (1995). Adrenergic calcium signaling in astrocyte networks within the hippocampal slice. *J. Neurosci.* *15*, 5535–5550.

Dutzler, R. (2004). Structural basis for ion conduction and gating in ClC chloride channels. *FEBS Lett.* *564*, 229–233.

Dutzler, R. (2007). A structural perspective on ClC channel and transporter function. *FEBS Lett.* *581*, 2839–2844.

Dutzler, R., Campbell, E.B., and MacKinnon, R. (2003). Gating the selectivity filter in ClC chloride channels. *Science* *300*, 108–112.

Ernest, N.J., Weaver, A.K., Van Duyn, L.B., and Sontheimer, H.W. (2005). Relative contribution of chloride channels and transporters to regulatory volume decrease in human glioma cells. *Am. J. Physiol. Cell Physiol.* *288*, C1451–C1460.

Estévez, R., and Jentsch, T.J. (2002). ClC chloride channels: correlating structure with function. *Curr. Opin. Struct. Biol.* *12*, 531–539.

Estévez, R., Pusch, M., Ferrer-Costa, C., Orozco, M., and Jentsch, T.J. (2004). Functional and structural conservation of CBS domains from ClC chloride channels. *J. Physiol.* *557*, 363–378.

Fahlke, C., Yu, H.T., Beck, C.L., Rhodes, T.H., and George, A.L. (1997). Pore-forming segments in voltage-gated chloride channels. *Nature* *390*, 529–532.

Favre-Kontula, L., Rolland, A., Bernasconi, L., Karmirantzou, M., Power, C., Antonsson, B., and Boschert, U. (2008). GlialCAM, an immunoglobulin-like cell adhesion molecule is expressed in glial cells of the central nervous system. *Glia* 56, 633–645.

Ferroni, S., Marchini, C., Nobile, M., and Rapisarda, C. (1997). Characterization of an inwardly rectifying chloride conductance expressed by cultured rat cortical astrocytes. *Glia* 21, 217–227.

Fischer, G., and Kettenmann, H. (1985). Cultured astrocytes form a syncytium after maturation. *Exp. Cell Res.* 159, 273–279.

Fritsch, J., and Edelman, A. (1996). Modulation of the hyperpolarization-activated Cl⁻ current in human intestinal T84 epithelial cells by phosphorylation. *J. Physiol.* 490 (Pt 1, 115–128.

Furukawa, T., Ogura, T., Katayama, Y., and Hiraoka, M. (1998). Characteristics of rabbit ClC-2 current expressed in *Xenopus* oocytes and its contribution to volume regulation. *Am. J. Physiol.* 274, C500–C512.

Gaudry, J.-P., Arod, C., Sauvage, C., Busso, S., Dupraz, P., Pankiewicz, R., and Antonsson, B. (2008). Purification of the extracellular domain of the membrane protein GlialCAM expressed in HEK and CHO cells and comparison of the glycosylation. *Protein Expr. Purif.* 58, 94–102.

Goodenough, D.A., and Paul, D.L. (2009). Gap junctions. *Cold Spring Harb. Perspect. Biol.* 1, a002576.

Gorospe, J.R., Singhal, B.S., Kainu, T., Wu, F., Stephan, D., Trent, J., Hoffman, E.P., and Naidu, S. (2004). Indian Agarwal megalencephalic leukodystrophy with cysts is caused by a common MLC1 mutation. *Neurology* 62, 878–882.

Graham, F.L., Smiley, J., Russell, W.C., and Nairn, R. (1977). Characteristics of a human cell line transformed by DNA from human adenovirus type 5. *J. Gen. Virol.* 36, 59–74.

Gründer, S., Thiemann, A., Pusch, M., and Jentsch, T.J. (1992). Regions involved in the opening of ClC-2 chloride channel by voltage and cell volume. *Nature* 360, 759–762.

Haj-Yasein, N.N., Jensen, V., Østby, I., Omholt, S.W., Voipio, J., Kaila, K., Ottersen, O.P., Hvalby, Ø., and Nagelhus, E.A. (2012). Aquaporin-4 regulates extracellular space volume dynamics during high-frequency synaptic stimulation: a gene deletion study in mouse hippocampus. *Glia* 60, 867–874.

Halassa, M.M., Fellin, T., Takano, H., Dong, J.-H., and Haydon, P.G. (2007). Synaptic islands defined by the territory of a single astrocyte. *J. Neurosci.* 27, 6473–6477.

Hewett, J.A. (2009). Determinants of regional and local diversity within the astroglial lineage of the normal central nervous system. *J. Neurochem.* 110, 1717–1736.

Hinzpeter, A., Lipecka, J., Brouillard, F., Baudoin-Legros, M., Dadlez, M., Edelman, A., and Fritsch, J. (2006). Association between Hsp90 and the ClC-2 chloride channel upregulates channel function. *Am. J. Physiol. Cell Physiol.* 290, C45–C56.

Hoegg-Beiler, M.B., Sirisi, S., Orozco, I.J., Ferrer, I., Hohensee, S., Auberson, M., Gödde, K., Vilches, C., de Heredia, M.L., Nunes, V., et al. (2014). Disrupting MLC1 and GlialCAM and ClC-2

interactions in leukodystrophy entails glial chloride channel dysfunction. *Nat. Commun.* *5*, 3475.

Hofmann, F., Lacinová, L., and Klugbauer, N. (1999). Voltage-dependent calcium channels: from structure to function. *Rev. Physiol. Biochem. Pharmacol.* *139*, 33–87.

Hosseinzadeh, Z., Bhavsar, S.K., and Lang, F. (2012). Downregulation of CLC-2 by JAK2. *Cell. Physiol. Biochem.* *29*, 737–742.

Hyzinski-García, M.C., Rudkouskaya, A., and Mongin, A.A. (2014). LRRC8A protein is indispensable for swelling-activated and ATP-induced release of excitatory amino acids in rat astrocytes. *J. Physiol.*

Iglesias, R., Dahl, G., Qiu, F., Spray, D.C., and Scemes, E. (2009). Pannexin 1: the molecular substrate of astrocyte “hemichannels”. *J. Neurosci.* *29*, 7092–7097.

Ilja Boor, P.K., de Groot, K., Mejaski-Bosnjak, V., Brenner, C., van der Knaap, M.S., Scheper, G.C., and Pronk, J.C. (2006). Megalencephalic leukoencephalopathy with subcortical cysts: an update and extended mutation analysis of MLC1. *Hum. Mutat.* *27*, 505–512.

Irrthum, A., Devriendt, K., Chitayat, D., Matthijs, G., Glade, C., Steijlen, P.M., Fryns, J.-P., Van Steensel, M.A.M., and Vikkula, M. (2003). Mutations in the transcription factor gene SOX18 underlie recessive and dominant forms of hypotrichosis-lymphedema-telangiectasia. *Am. J. Hum. Genet.* *72*, 1470–1478.

Jayakumar, A.R., and Norenberg, M.D. (2010). The Na-K-Cl Co-transporter in astrocyte swelling. *Metab. Brain Dis.* *25*, 31–38.

Jayakumar, A.R., Panickar, K.S., Curtis, K.M., Tong, X.Y., Moriyama, M., and Norenberg, M.D. (2011). Na-K-Cl cotransporter-1 in the mechanism of cell swelling in cultured astrocytes after fluid percussion injury. *J. Neurochem.* *117*, 437–448.

Jentsch, T.J., Friedrich, T., Schriever, A., and Yamada, H. (1999). The CLC chloride channel family. *Pflugers Arch.* *437*, 783–795.

Jeworutzki, E., López-Hernández, T., Capdevila-Nortes, X., Sirisi, S., Bengtsson, L., Montolio, M., Zifarelli, G., Arnedo, T., Müller, C.S., Schulte, U., et al. (2012). GlialCAM, a protein defective in a leukodystrophy, serves as a CLC-2 Cl(-) channel auxiliary subunit. *Neuron* *73*, 951–961.

Jeworutzki, E., Lagostena, L., Elorza-Vidal, X., López-Hernández, T., Estévez, R., and Pusch, M. (2014). GlialCAM, a CLC-2 Cl(-) channel subunit, activates the slow gate of CLC chloride channels. *Biophys. J.* *107*, 1105–1116.

Jordt, S.E., and Jentsch, T.J. (1997). Molecular dissection of gating in the CLC-2 chloride channel. *EMBO J.* *16*, 1582–1592.

Kaganovich, M., Peretz, A., Ritsner, M., Bening Abu-Shach, U., Attali, B., and Navon, R. (2004). Is the WKL1 gene associated with schizophrenia? *Am. J. Med. Genet. B. Neuropsychiatr. Genet.* *125B*, 31–37.

Kajita, H., Omori, K., and Matsuda, H. (2000). The chloride channel ClC-2 contributes to the inwardly rectifying Cl⁻ conductance in cultured porcine choroid plexus epithelial cells. *J. Physiol.* 523 Pt 2, 313–324.

Kalsi, A.S., Greenwood, K., Wilkin, G., and Butt, A.M. (2004). Kir4.1 expression by astrocytes and oligodendrocytes in CNS white matter: a developmental study in the rat optic nerve. *J. Anat.* 204, 475–485.

Kang, J., Jiang, L., Goldman, S.A., and Nedergaard, M. (1998). Astrocyte-mediated potentiation of inhibitory synaptic transmission. *Nat. Neurosci.* 1, 683–692.

Kariminejad, A., Rajaei, A., Ashrafi, M.R., Alizadeh, H., Tonekaboni, S.H., Malamiri, R.A., Ghofrani, M., Karimzadeh, P., Mohammadi, M.M., Baghalshooshtari, A., et al. (2014). Eight Novel mutations in MLC1 from 18 Iranian patients with Megalencephalic Leukoencephalopathy with Subcortical Cysts. *Eur. J. Med. Genet.*

Keeling, D.J., Fallowfield, C., Milliner, K.J., Tingley, S.K., Ife, R.J., and Underwood, A.H. (1985). Studies on the mechanism of action of omeprazole. *Biochem. Pharmacol.* 34, 2967–2973.

Kimelberg, H.K., Goderie, S.K., Higman, S., Pang, S., and Waniewski, R.A. (1990). Swelling-induced release of glutamate, aspartate, and taurine from astrocyte cultures. *J. Neurosci.* 10, 1583–1591.

Kimelberg, H.K., Macvicar, B.A., and Sontheimer, H. (2006). Anion channels in astrocytes: biophysics, pharmacology, and function. *Glia* 54, 747–757.

Van der Knaap, M.S., Valk, J., Barth, P.G., Smit, L.M., van Engelen, B.G., and Tortori Donati, P. (1995a). Leukoencephalopathy with swelling in children and adolescents: MRI patterns and differential diagnosis. *Neuroradiology* 37, 679–686.

Van der Knaap, M.S., Barth, P.G., Stroink, H., van Nieuwenhuizen, O., Arts, W.F., Hoogenraad, F., and Valk, J. (1995b). Leukoencephalopathy with swelling and a discrepantly mild clinical course in eight children. *Ann. Neurol.* 37, 324–334.

Van der Knaap, M.S., Barth, P.G., Vrensen, G.F., and Valk, J. (1996). Histopathology of an infantile-onset spongiform leukoencephalopathy with a discrepantly mild clinical course. *Acta Neuropathol.* 92, 206–212.

Van der Knaap, M.S., Lai, V., Köhler, W., Salih, M.A., Fonseca, M.-J., Benke, T.A., Wilson, C., Jayakar, P., Aine, M., Dom, L., et al. (2010). Megalencephalic leukoencephalopathy with cysts without MLC1 defect. *Ann. Neurol.* 67, 834–837.

Van der Knaap, M.S., Boor, I., and Estévez, R. (2012). Megalencephalic leukoencephalopathy with subcortical cysts: Chronic white matter oedema due to a defect in brain ion and water homeostasis. *Lancet Neurol.* 11, 973–984.

Kobe, B., and Kajava, A. V (2001). The leucine-rich repeat as a protein recognition motif. *Curr. Opin. Struct. Biol.* 11, 725–732.

Kofuji, P., and Newman, E.A. (2004). Potassium buffering in the central nervous system. *Neuroscience* 129, 1045–1056.

Koussa, S., Roukoz, H., Rizk, T., and Mégarbané, A. (2005). [Megalencephalic leukoencephalopathy with subcortical cysts: a study of a Lebanese family and a review of the literature]. *Rev. Neurol. (Paris)*. *161*, 183–191.

Lampe, P.D., and Lau, A.F. (2000). Regulation of gap junctions by phosphorylation of connexins. *Arch. Biochem. Biophys.* *384*, 205–215.

Lanciotti, A., Brignone, M.S., Camerini, S., Serafini, B., Macchia, G., Raggi, C., Molinari, P., Crescenzi, M., Musumeci, M., Sargiacomo, M., et al. (2010). MLC1 trafficking and membrane expression in astrocytes: role of caveolin-1 and phosphorylation. *Neurobiol. Dis.* *37*, 581–595.

Lanciotti, A., Brignone, M.S., Molinari, P., Visentin, S., De Nuccio, C., Macchia, G., Aiello, C., Bertini, E., Aloisi, F., Petrucci, T.C., et al. (2012). Megalencephalic leukoencephalopathy with subcortical cysts protein 1 functionally cooperates with the TRPV4 cation channel to activate the response of astrocytes to osmotic stress: dysregulation by pathological mutations. *Hum. Mol. Genet.* *21*, 2166–2180.

Landy, A. (1989). Dynamic, structural, and regulatory aspects of lambda site-specific recombination. *Annu. Rev. Biochem.* *58*, 913–949.

Larsen, B.R., Assentoft, M., Cotrina, M.L., Hua, S.Z., Nedergaard, M., Kaila, K., Voipio, J., and MacAulay, N. (2014). Contributions of the Na⁺/K⁺-ATPase, NKCC1, and Kir4.1 to hippocampal K⁺ clearance and volume responses. *Glia* *62*, 608–622.

Lee, L.H., Moh, M.C., Zhang, T., and Shen, S. (2009). The immunoglobulin-like cell adhesion molecule hepaCAM induces differentiation of human glioblastoma U373-MG cells. *J. Cell. Biochem.* *107*, 1129–1138.

Leegwater, P.A., Yuan, B.Q., van der Steen, J., Mulders, J., Könst, A.A., Boor, P.K., Mejaski-Bosnjak, V., van der Maarel, S.M., Frants, R.R., Oudejans, C.B., et al. (2001). Mutations of MLC1 (KIAA0027), encoding a putative membrane protein, cause megalencephalic leukoencephalopathy with subcortical cysts. *Am. J. Hum. Genet.* *68*, 831–838.

Leegwater, P.A.J., Boor, P.K.I., Yuan, B.Q., van der Steen, J., Visser, A., Könst, A.A.M., Oudejans, C.B.M., Schutgens, R.B.H., Pronk, J.C., and van der Knaap, M.S. (2002). Identification of novel mutations in MLC1 responsible for megalencephalic leukoencephalopathy with subcortical cysts. *Hum. Genet.* *110*, 279–283.

Liedtke, W., Edelmann, W., Bieri, P.L., Chiu, F.C., Cowan, N.J., Kucherlapati, R., and Raine, C.S. (1996). GFAP is necessary for the integrity of CNS white matter architecture and long-term maintenance of myelination. *Neuron* *17*, 607–615.

Liu, X., Bandyopadhyay, B.C., Bandyopadhyay, B., Nakamoto, T., Singh, B., Liedtke, W., Melvin, J.E., and Ambudkar, I. (2006). A role for AQP5 in activation of TRPV4 by hypotonicity: concerted involvement of AQP5 and TRPV4 in regulation of cell volume recovery. *J. Biol. Chem.* *281*, 15485–15495.

López-Hernández, T., Ridder, M.C., Montolio, M., Capdevila-Nortes, X., Polder, E., Sirisi, S., Duarri, A., Schulte, U., Fakler, B., Nunes, V., et al. (2011a). Mutant GlialCAM causes megalencephalic leukoencephalopathy with subcortical cysts, benign familial macrocephaly, and macrocephaly with retardation and autism. *Am. J. Hum. Genet.* *88*, 422–432.

López-Hernández, T., Sirisi, S., Capdevila-Nortes, X., Montolio, M., Fernández-Dueñas, V., Scheper, G.C., van der Knaap, M.S., Casquero, P., Ciruela, F., Ferrer, I., et al. (2011b). Molecular mechanisms of MLC1 and GLIALCAM mutations in megalencephalic leukoencephalopathy with subcortical cysts. *Hum. Mol. Genet.* *20*, 3266–3277.

Lu, L., Sirish, P., Zhang, Z., Woltz, R.L., Li, N., Timofeyev, V., Knowlton, A.A., Zhang, X.-D., Yamoah, E.N., and Chiamvimonvat, N. (2014). Regulation of Gene Transcription by Voltage-Gated L-Type Calcium Channel, Cav1.3. *J. Biol. Chem.*

MacVicar, B.A., Feighan, D., Brown, A., and Ransom, B. (2002). Intrinsic optical signals in the rat optic nerve: role for K(+) uptake via NKCC1 and swelling of astrocytes. *Glia* *37*, 114–123.

Maglione, M., Tress, O., Haas, B., Karram, K., Trotter, J., Willecke, K., and Kettenmann, H. (2010). Oligodendrocytes in mouse corpus callosum are coupled via gap junction channels formed by connexin47 and connexin32. *Glia* *58*, 1104–1117.

Mahmoud, I.G., Mahmoud, M., Refaat, M., Girgis, M., Waked, N., El Badawy, A., Selim, L., Hassan, S., and Abdel Aleem, A.K. (2014). Clinical, neuroimaging, and genetic characteristics of megalencephalic leukoencephalopathy with subcortical cysts in Egyptian patients. *Pediatr. Neurol.* *50*, 140–148.

Makara, J.K., Rappert, A., Matthias, K., Steinhäuser, C., Spät, A., and Kettenmann, H. (2003). Astrocytes from mouse brain slices express CIC-2-mediated Cl⁻ currents regulated during development and after injury. *Mol. Cell. Neurosci.* *23*, 521–530.

Malarkey, E.B., Ni, Y., and Parpura, V. (2008). Ca²⁺ entry through TRPC1 channels contributes to intracellular Ca²⁺ dynamics and consequent glutamate release from rat astrocytes. *Glia* *56*, 821–835.

Markovic, S., and Dutzler, R. (2007). The structure of the cytoplasmic domain of the chloride channel CIC-Ka reveals a conserved interaction interface. *Structure* *15*, 715–725.

McDonald, T.F., Pelzer, S., Trautwein, W., and Pelzer, D.J. (1994). Regulation and modulation of calcium channels in cardiac, skeletal, and smooth muscle cells. *Physiol. Rev.* *74*, 365–507.

Menichella, D.M., Goodenough, D.A., Sirkowski, E., Scherer, S.S., and Paul, D.L. (2003). Connexins are critical for normal myelination in the CNS. *J. Neurosci.* *23*, 5963–5973.

Menichella, D.M., Majdan, M., Awatramani, R., Goodenough, D.A., Sirkowski, E., Scherer, S.S., and Paul, D.L. (2006). Genetic and physiological evidence that oligodendrocyte gap junctions contribute to spatial buffering of potassium released during neuronal activity. *J. Neurosci.* *26*, 10984–10991.

Miller, C. (1982). Open-state substructure of single chloride channels from Torpedo electroplax. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* *299*, 401–411.

Miller, C. (2006). CIC chloride channels viewed through a transporter lens. *Nature* *440*, 484–489.

Moeser, A.J., Haskell, M.M., Shifflett, D.E., Little, D., Schultz, B.D., and Blikslager, A.T. (2004). ClC-2 chloride secretion mediates prostaglandin-induced recovery of barrier function in ischemia-injured porcine ileum. *Gastroenterology* 127, 802–815.

Moeser, A.J., Nighot, P.K., Engelke, K.J., Ueno, R., and Blikslager, A.T. (2007). Recovery of mucosal barrier function in ischemic porcine ileum and colon is stimulated by a novel agonist of the ClC-2 chloride channel, lubiprostone. *Am. J. Physiol. Gastrointest. Liver Physiol.* 292, G647–G656.

Moghaddam, B., and Adams, R.N. (1987). Regional differences in resting extracellular potassium levels of rat brain. *Brain Res.* 406, 337–340.

Moh, M.C., Zhang, C., Luo, C., Lee, L.H., and Shen, S. (2005). Structural and functional analyses of a novel ig-like cell adhesion molecule, hepaCAM, in the human breast carcinoma MCF7 cells. *J. Biol. Chem.* 280, 27366–27374.

Moh, M.C., Tian, Q., Zhang, T., Lee, L.H., and Shen, S. (2009). The immunoglobulin-like cell adhesion molecule hepaCAM modulates cell adhesion and motility through direct interaction with the actin cytoskeleton. *J. Cell. Physiol.* 219, 382–391.

Morales, M.M., Nascimento, D.S., Capella, M.A., Lopes, A.G., and Guggino, W.B. (2001). Arginine vasopressin regulates CFTR and ClC-2 mRNA expression in rat kidney cortex and medulla. *Pflugers Arch.* 443, 202–211.

Moser, T., Chow, R.H., and Neher, E. (1995). Swelling-induced catecholamine secretion recorded from single chromaffin cells. *Pflugers Arch.* 431, 196–203.

Nagelhus, E.A., Mathiisen, T.M., and Ottersen, O.P. (2004). Aquaporin-4 in the central nervous system: cellular and subcellular distribution and coexpression with KIR4.1. *Neuroscience* 129, 905–913.

Nagy, J.I., Dudek, F.E., and Rash, J.E. (2004). Update on connexins and gap junctions in neurons and glia in the mammalian nervous system. *Brain Res. Brain Res. Rev.* 47, 191–215.

Nascimento, D.S., Reis, C.U., Goldenberg, R.C., Ortiga-Carvalho, T.M., Pazos-Moura, C.C., Guggino, S.E., Guggino, W.B., and Morales, M.M. (2003). Estrogen modulates ClC-2 chloride channel gene expression in rat kidney. *Pflugers Arch.* 446, 593–599.

Nedergaard, M. (1994). Direct signaling from astrocytes to neurons in cultures of mammalian brain cells. *Science* 263, 1768–1771.

Nedergaard, M., Ransom, B., and Goldman, S.A. (2003). New roles for astrocytes: redefining the functional architecture of the brain. *Trends Neurosci.* 26, 523–530.

Nehrke, K., Arreola, J., Nguyen, H.-V., Pilato, J., Richardson, L., Okunade, G., Baggs, R., Shull, G.E., and Melvin, J.E. (2002). Loss of hyperpolarization-activated Cl⁻ current in salivary acinar cells from *Clcn2* knockout mice. *J. Biol. Chem.* 277, 23604–23611.

Neusch, C., Rozengurt, N., Jacobs, R.E., Lester, H.A., and Kofuji, P. (2001). Kir4.1 potassium channel subunit is crucial for oligodendrocyte development and in vivo myelination. *J. Neurosci.* 21, 5429–5438.

Newman, E.A. Regional specialization of retinal glial cell membrane. *Nature* 309, 155–157.

Nicchia, G.P., Srinivas, M., Li, W., Brosnan, C.F., Frigeri, A., and Spray, D.C. (2005). New possible roles for aquaporin-4 in astrocytes: cell cytoskeleton and functional relationship with connexin43. *FASEB J.* 19, 1674–1676.

Niemeyer, M.I., Cid, L.P., Zúñiga, L., Catalán, M., and Sepúlveda, F. V (2003). A conserved pore-lining glutamate as a voltage- and chloride-dependent gate in the ClC-2 chloride channel. *J. Physiol.* 553, 873–879.

Nighot, P.K., and Blikslager, A.T. (2010). ClC-2 regulates mucosal barrier function associated with structural changes to the villus and epithelial tight junction. *Am. J. Physiol. Gastrointest. Liver Physiol.* 299, G449–G456.

Nighot, P.K., and Blikslager, A.T. (2012). Chloride channel ClC-2 modulates tight junction barrier function via intracellular trafficking of occludin. *Am. J. Physiol. Cell Physiol.* 302, C178–C187.

Nighot, P.K., Moeser, A.J., Ryan, K.A., Ghashghaei, T., and Blikslager, A.T. (2009). ClC-2 is required for rapid restoration of epithelial tight junctions in ischemic-injured murine jejunum. *Exp. Cell Res.* 315, 110–118.

Nilius, B., and Droogmans, G. (2003). Amazing chloride channels: an overview. *Acta Physiol. Scand.* 177, 119–147.

Nilius, B., Eggermont, J., Voets, T., Buyse, G., Manolopoulos, V., and Droogmans, G. (1997). Properties of volume-regulated anion channels in mammalian cells. *Prog. Biophys. Mol. Biol.* 68, 69–119.

Nualart-Marti, A., Solsona, C., and Fields, R.D. (2013). Gap junction communication in myelinating glia. *Biochim. Biophys. Acta* 1828, 69–78.

Oberheim, N.A., Wang, X., Goldman, S., and Nedergaard, M. (2006). Astrocytic complexity distinguishes the human brain. *Trends Neurosci.* 29, 547–553.

Odermatt, B., Wellershaus, K., Wallraff, A., Seifert, G., Degen, J., Euwens, C., Fuss, B., Büssow, H., Schilling, K., Steinhäuser, C., et al. (2003). Connexin 47 (Cx47)-deficient mice with enhanced green fluorescent protein reporter gene reveal predominant oligodendrocytic expression of Cx47 and display vacuolized myelin in the CNS. *J. Neurosci.* 23, 4549–4559.

Ogata, K., and Kosaka, T. (2002). Structural and quantitative analysis of astrocytes in the mouse hippocampus. *Neuroscience* 113, 221–233.

Okada, Y. (1997). Volume expansion-sensing outward-rectifier Cl⁻ channel: fresh start to the molecular identity and volume sensor. *Am. J. Physiol.* 273, C755–C789.

Okada, Y., Shimizu, T., Maeno, E., Tanabe, S., Wang, X., and Takahashi, N. (2006). Volume-sensitive chloride channels involved in apoptotic volume decrease and cell death. *J. Membr. Biol.* 209, 21–29.

Okada, Y., Sato, K., and Numata, T. (2009). Pathophysiology and puzzles of the volume-sensitive outwardly rectifying anion channel. *J. Physiol.* *587*, 2141–2149.

Olsen, M.L., Higashimori, H., Campbell, S.L., Hablitz, J.J., and Sontheimer, H. (2006). Functional expression of Kir4.1 channels in spinal cord astrocytes. *Glia* *53*, 516–528.

Orkand, R.K., Nicholls, J.G., and Kuffler, S.W. (1966). Effect of nerve impulses on the membrane potential of glial cells in the central nervous system of amphibia. *J. Neurophysiol.* *29*, 788–806.

Ornellas, D.S., Nascimento, D.S., Christoph, D.H., Guggino, W.B., and Morales, M.M. (2002). Aldosterone and high-NaCl diet modulate CIC-2 chloride channel gene expression in rat kidney. *Pflugers Arch.* *444*, 193–201.

Padmanabhan, M., Cournoyer, P., and Dinesh-Kumar, S.P. (2009). The leucine-rich repeat domain in plant innate immunity: a wealth of possibilities. *Cell. Microbiol.* *11*, 191–198.

Pan, Z., Yang, H., Mergler, S., Liu, H., Tachado, S.D., Zhang, F., Kao, W.W.Y., Koziel, H., Pleyer, U., and Reinach, P.S. (2008). Dependence of regulatory volume decrease on transient receptor potential vanilloid 4 (TRPV4) expression in human corneal epithelial cells. *Cell Calcium* *44*, 374–385.

Pasantes-Morales, H., Lezama, R.A., Ramos-Mandujano, G., and Tuz, K.L. (2006). Mechanisms of cell volume regulation in hypo-osmolality. *Am. J. Med.* *119*, S4–S11.

Pascual-Castroviejo, I., van der Knaap, M.S., Pronk, J.C., García-Segura, J.M., Gutiérrez-Molina, M., and Pascual-Pascual, S.I. (2005). Vacuolating megalencephalic leukoencephalopathy: 24 year follow-up of two siblings. *Neurologia* *20*, 33–40.

Patrono, C., Di Giacinto, G., Eymard-Pierre, E., Santorelli, F.M., Rodriguez, D., De Stefano, N., Federico, A., Gatti, R., Benigno, V., Megarbané, A., et al. (2003). Genetic heterogeneity of megalencephalic leukoencephalopathy and subcortical cysts. *Neurology* *61*, 534–537.

Plotkin, M.D., Snyder, E.Y., Hebert, S.C., and Delpire, E. (1997). Expression of the Na-K-2Cl cotransporter is developmentally regulated in postnatal rat brains: a possible mechanism underlying GABA's excitatory role in immature brain. *J. Neurobiol.* *33*, 781–795.

Ponting, C.P. (1997). CBS domains in CIC chloride channels implicated in myotonia and nephrolithiasis (kidney stones). *J. Mol. Med. (Berl)*. *75*, 160–163.

Qiu, Z., Dubin, A.E., Mathur, J., Tu, B., Reddy, K., Miraglia, L.J., Reinhardt, J., Orth, A.P., and Patapoutian, A. (2014). SWELL1, a plasma membrane protein, is an essential component of volume-regulated anion channel. *Cell* *157*, 447–458.

Raffaele Di Barletta, M., Ricci, E., Galluzzi, G., Tonali, P., Mora, M., Morandi, L., Romorini, A., Voit, T., Orstavik, K.H., Merlini, L., et al. (2000). Different mutations in the LMNA gene cause autosomal dominant and autosomal recessive Emery-Dreifuss muscular dystrophy. *Am. J. Hum. Genet.* *66*, 1407–1412.

Ransom, C.B., Ransom, B.R., and Sontheimer, H. (2000). Activity-dependent extracellular K⁺ accumulation in rat optic nerve: the role of glial and axonal Na⁺ pumps. *J. Physiol.* *522 Pt 3*, 427–442.

Rash, J.E. (2010). Molecular disruptions of the panglial syncytium block potassium siphoning and axonal saltatory conduction: pertinence to neuromyelitis optica and other demyelinating diseases of the central nervous system. *Neuroscience* 168, 982–1008.

Rash, J.E., Duffy, H.S., Dudek, F.E., Bilhartz, B.L., Whalen, L.R., and Yasumura, T. (1997). Grid-mapped freeze-fracture analysis of gap junctions in gray and white matter of adult rat central nervous system, with evidence for a “panglial syncytium” that is not coupled to neurons. *J. Comp. Neurol.* 388, 265–292.

Rash, J.E., Yasumura, T., Davidson, K.G., Furman, C.S., Dudek, F.E., and Nagy, J.I. (2001). Identification of cells expressing Cx43, Cx30, Cx26, Cx32 and Cx36 in gap junctions of rat brain and spinal cord. *Cell Commun. Adhes.* 8, 315–320.

Ratté, S., and Prescott, S.A. (2011). ClC-2 channels regulate neuronal excitability, not intracellular chloride levels. *J. Neurosci.* 31, 15838–15843.

Ribeiro, M. de C., Hirt, L., Bogousslavsky, J., Regli, L., and Badaut, J. (2006). Time course of aquaporin expression after transient focal cerebral ischemia in mice. *J. Neurosci. Res.* 83, 1231–1240.

Ridder, M.C., Boor, I., Lodder, J.C., Postma, N.L., Capdevila-Nortes, X., Duarri, A., Brussaard, A.B., Estévez, R., Scheper, G.C., Mansvelder, H.D., et al. (2011). Megalencephalic leucoencephalopathy with cysts: defect in chloride currents and cell volume regulation. *Brain* 134, 3342–3354.

Rinke, I., Artmann, J., and Stein, V. (2010). ClC-2 voltage-gated channels constitute part of the background conductance and assist chloride extrusion. *J. Neurosci.* 30, 4776–4786.

Saiki, R.K., Gelfand, D.H., Stoffel, S., Scharf, S.J., Higuchi, R., Horn, G.T., Mullis, K.B., and Erlich, H.A. (1988). Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science* 239, 487–491.

Saint-Martin, C., Gouvain, G., Teodorescu, G., Gourfinkel-An, I., Fedirko, E., Weber, Y.G., Maljevic, S., Ernst, J.-P., Garcia-Olivares, J., Fahlke, C., et al. (2009). Two novel CLCN2 mutations accelerating chloride channel deactivation are associated with idiopathic generalized epilepsy. *Hum. Mutat.* 30, 397–405.

De Santiago, J.A., Nehrke, K., and Arreola, J. (2005). Quantitative analysis of the voltage-dependent gating of mouse parotid ClC-2 chloride channel. *J. Gen. Physiol.* 126, 591–603.

Sargiannidou, I., Markoullis, K., and Kleopa, K.A. (2010). Molecular mechanisms of gap junction mutations in myelinating cells. *Histol. Histopathol.* 25, 1191–1206.

Sawada, A., Takihara, Y., Kim, J.Y., Matsuda-Hashii, Y., Tokimasa, S., Fujisaki, H., Kubota, K., Endo, H., Onodera, T., Ohta, H., et al. (2003). A congenital mutation of the novel gene LRRC8 causes agammaglobulinemia in humans. *J. Clin. Invest.* 112, 1707–1713.

Scemes, E., Spray, D.C., and Meda, P. (2009). Connexins, pannexins, innexins: novel roles of “hemi-channels”. *Pflugers Arch.* 457, 1207–1226.

Scheper, G.C., van Berkel, C.G.M., Leisle, L., de Groot, K.E., Errami, A., Jentsch, T.J., and Van der Knaap, M.S. (2010). Analysis of CLCN2 as candidate gene for megalencephalic leukoencephalopathy with subcortical cysts. *Genet. Test. Mol. Biomarkers* 14, 255–257.

Schwiebert, E.M., Cid-Soto, L.P., Stafford, D., Carter, M., Blaisdell, C.J., Zeitlin, P.L., Guggino, W.B., and Cutting, G.R. (1998). Analysis of ClC-2 channels as an alternative pathway for chloride conduction in cystic fibrosis airway cells. *Proc. Natl. Acad. Sci. U. S. A.* 95, 3879–3884.

Sen, E., and Levison, S.W. (2006). Astrocytes and developmental white matter disorders. *Ment. Retard. Dev. Disabil. Res. Rev.* 12, 97–104.

Sener, R.N. (2003a). Leukoencephalopathy with patchy lesions, temporal cystic degeneration, and normal head size. *Comput. Med. Imaging Graph.* 27, 87–91.

Sener, R.N. (2003b). Proton MR spectroscopy demonstration of taurine peaks in megalencephalic leukoencephalopathy with cysts. *Comput. Med. Imaging Graph.* 27, 23–26.

Shin, K., Fogg, V.C., and Margolis, B. (2006). Tight junctions and cell polarity. *Annu. Rev. Cell Dev. Biol.* 22, 207–235.

Sík, A., Smith, R.L., and Freund, T.F. (2000). Distribution of chloride channel-2-immunoreactive neuronal and astrocytic processes in the hippocampus. *Neuroscience* 101, 51–65.

Simard, M., Arcuino, G., Takano, T., Liu, Q.S., and Nedergaard, M. (2003). Signaling at the gliovascular interface. *J. Neurosci.* 23, 9254–9262.

Singhal (1991). Megalencephalic leukoencephalopathy in India. In *Proceedings of the 8th Asian and Oceanian Congress of Neurolog.*,

Singhal, B.S., Gursahani, R.D., Udani, V.P., and Biniwale, A.A. (1996). Megalencephalic leukodystrophy in an Asian Indian ethnic group. *Pediatr. Neurol.* 14, 291–296.

Sirisi, S., Folgueira, M., López-Hernández, T., Minieri, L., Pérez-Rius, C., Gaitán-Peñas, H., Zang, J., Martínez, A., Capdevila-Nortes, X., De La Villa, P., et al. (2014). Megalencephalic leukoencephalopathy with subcortical cysts protein 1 regulates glial surface localization of GLIALCAM from fish to humans. *Hum. Mol. Genet.* 23, 5069–5086.

Smith, R.L., Clayton, G.H., Wilcox, C.L., Escudero, K.W., and Staley, K.J. (1995). Differential expression of an inwardly rectifying chloride conductance in rat brain neurons: a potential mechanism for cell-specific modulation of postsynaptic inhibition. *J. Neurosci.* 15, 4057–4067.

Song, X., Zhao, Y., Narcisse, L., Duffy, H., Kress, Y., Lee, S., and Brosnan, C.F. (2005). Canonical transient receptor potential channel 4 (TRPC4) co-localizes with the scaffolding protein ZO-1 in human fetal astrocytes in culture. *Glia* 49, 418–429.

Staley, K., Smith, R., Schaack, J., Wilcox, C., and Jentsch, T.J. (1996). Alteration of GABAA receptor function following gene transfer of the CLC-2 chloride channel. *Neuron* 17, 543–551.

Stölting, G., Teodorescu, G., Begemann, B., Schubert, J., Nabbout, R., Toliat, M.R., Sander, T., Nürnberg, P., Lerche, H., and Fahlke, C. (2013). Regulation of ClC-2 gating by intracellular ATP. *Pflugers Arch.* 465, 1423–1437.

Su, G., Kintner, D.B., and Sun, D. (2002a). Contribution of Na(+)-K(+)-Cl(-) cotransporter to high-[K(+)](o)- induced swelling and EAA release in astrocytes. *Am. J. Physiol. Cell Physiol.* **282**, C1136–C1146.

Su, G., Kintner, D.B., Flagella, M., Shull, G.E., and Sun, D. (2002b). Astrocytes from Na(+)-K(+)-Cl(-) cotransporter-null mice exhibit absence of swelling and decrease in EAA release. *Am. J. Physiol. Cell Physiol.* **282**, C1147–C1160.

Teijido, O., Martínez, A., Pusch, M., Zorzano, A., Soriano, E., Del Río, J.A., Palacín, M., and Estévez, R. (2004). Localization and functional analyses of the MLC1 protein involved in megalencephalic leukoencephalopathy with subcortical cysts. *Hum. Mol. Genet.* **13**, 2581–2594.

Teijido, O., Casaroli-Marano, R., Kharkovets, T., Aguado, F., Zorzano, A., Palacín, M., Soriano, E., Martínez, A., and Estévez, R. (2007). Expression patterns of MLC1 protein in the central and peripheral nervous systems. *Neurobiol. Dis.* **26**, 532–545.

Tewari, K.P., Malinowska, D.H., Sherry, A.M., and Cuppoletti, J. (2000). PKA and arachidonic acid activation of human recombinant ClC-2 chloride channels. *Am. J. Physiol. Cell Physiol.* **279**, C40–C50.

Theodosios, D.T., Poulain, D.A., and Oliet, S.H.R. (2008). Activity-dependent structural and functional plasticity of astrocyte-neuron interactions. *Physiol. Rev.* **88**, 983–1008.

Thiemann, A., Gründer, S., Pusch, M., and Jentsch, T.J. (1992). A chloride channel widely expressed in epithelial and non-epithelial cells. *Nature* **356**, 57–60.

Thompson, C.H., Olivetti, P.R., Fuller, M.D., Freeman, C.S., McMaster, D., French, R.J., Pohl, J., Kubanek, J., and McCarty, N.A. (2009). Isolation and characterization of a high affinity peptide inhibitor of ClC-2 chloride channels. *J. Biol. Chem.* **284**, 26051–26062.

Tinsa, F., Farid, O., Douira, W., Rodriguez, D., Burglen, L., Boussetta, K., and Bousnina, S. (2009). Megalencephalic leukoencephalopathy with subcortical cysts in a Tunisian boy. *J. Child Neurol.* **24**, 87–89.

Topcu, M., Saatci, I., Topcuoglu, M.A., Kose, G., and Kunak, B. (1998). Megalencephaly and leukodystrophy with mild clinical course: a report on 12 new cases. *Brain Dev.* **20**, 142–153.

Turner, J.R. (2006). Molecular basis of epithelial barrier regulation: from basic mechanisms to clinical application. *Am. J. Pathol.* **169**, 1901–1909.

Valverde, M.A., Cantero-Recasens, G., Garcia-Elias, A., Jung, C., Carreras-Sureda, A., and Vicente, R. (2011). Ion channels in asthma. *J. Biol. Chem.* **286**, 32877–32882.

Vázquez-Juárez, E., Ramos-Mandujano, G., Hernández-Benítez, R., and Pasantes-Morales, H. (2008). On the role of G-protein coupled receptors in cell volume regulation. *Cell. Physiol. Biochem.* **21**, 1–14.

Voss, F.K., Ullrich, F., Münch, J., Lazarow, K., Lutter, D., Mah, N., Andrade-Navarro, M.A., von Kries, J.P., Stauber, T., and Jentsch, T.J. (2014). Identification of LRRC8 heteromers as an essential component of the volume-regulated anion channel VRAC. *Science* 344, 634–638.

Wallraff, A., Köhling, R., Heinemann, U., Theis, M., Willecke, K., and Steinhäuser, C. (2006). The impact of astrocytic gap junctional coupling on potassium buffering in the hippocampus. *J. Neurosci.* 26, 5438–5447.

Walz, W. (2002). Chloride/anion channels in glial cell membranes. *Glia* 40, 1–10.

Wang, D.D., and Bordey, A. (2008). The astrocyte odyssey. *Prog. Neurobiol.* 86, 342–367.

Wang, F., Smith, N.A., Xu, Q., Fujita, T., Baba, A., Matsuda, T., Takano, T., Bekar, L., and Nedergaard, M. (2012). Astrocytes modulate neural network activity by Ca²⁺-dependent uptake of extracellular K⁺. *Sci. Signal.* 5, ra26.

Warth, A., Mittelbronn, M., and Wolburg, H. (2005). Redistribution of the water channel protein aquaporin-4 and the K⁺ channel protein Kir4.1 differs in low- and high-grade human brain tumors. *Acta Neuropathol.* 109, 418–426.

Wasseff, S.K., and Scherer, S.S. (2011). Cx32 and Cx47 mediate oligodendrocyte:astrocyte and oligodendrocyte:oligodendrocyte gap junction coupling. *Neurobiol. Dis.* 42, 506–513.

Xun, C., Luo, C., Wu, X., Zhang, Q., Yan, L., and Shen, S. (2010). Expression of hepaCAM and its effect on proliferation of tumor cells in renal cell carcinoma. *Urology* 75, 828–834.

Yamane, Y., Shiga, H., Asou, H., and Ito, E. (2002). GAP junctional channel inhibition alters actin organization and calcium propagation in rat cultured astrocytes. *Neuroscience* 112, 593–603.

Yan, Q., Malashkevich, V.N., Fedorov, A., Fedorov, E., Cao, E., Lary, J.W., Cole, J.L., Nathenson, S.G., and Almo, S.C. (2007). Structure of CD84 provides insight into SLAM family function. *Proc. Natl. Acad. Sci. U. S. A.* 104, 10583–10588.

Yankaskas, J.R., Marshall, B.C., Sufian, B., Simon, R.H., and Rodman, D. (2004). Cystic fibrosis adult care: consensus conference report. *Chest* 125, 1S – 39S.

Yiş, U., Scheper, G.C., Uran, N., Unalp, A., Cakmakçi, H., Hiz-Kurul, S., Dirik, E., and van der Knaap, M.S. Two cases with megalencephalic leukoencephalopathy with subcortical cysts and MLC1 mutations in the Turkish population. *Turk. J. Pediatr.* 52, 179–183.

Yum, S.W., Kleopa, K.A., Shumas, S., and Scherer, S.S. (2002). Diverse Trafficking Abnormalities of Connexin32 Mutants Causing CMTX. *Neurobiol. Dis.* 11, 43–52.

Zdebik, A.A., Cuffe, J.E., Bertog, M., Korbmayer, C., and Jentsch, T.J. (2004). Additional disruption of the ClC-2 Cl(-) channel does not exacerbate the cystic fibrosis phenotype of cystic fibrosis transmembrane conductance regulator mouse models. *J. Biol. Chem.* 279, 22276–22283.

Zhang, H., and Verkman, A.S. (2008). Evidence against involvement of aquaporin-4 in cell-cell adhesion. *J. Mol. Biol.* 382, 1136–1143.

Zhang, Y., Zhang, H., Feustel, P.J., and Kimelberg, H.K. (2008). DCPIB, a specific inhibitor of volume regulated anion channels (VRACs), reduces infarct size in MCAo and the release of glutamate in the ischemic cortical penumbra. *Exp. Neurol.* *210*, 514–520.

Zúñiga, L., Niemeyer, M.I., Varela, D., Catalán, M., Cid, L.P., and Sepúlveda, F. V (2004). The voltage-dependent ClC-2 chloride channel has a dual gating mechanism. *J. Physiol.* *555*, 671–682.