

- Capapé, C., Zaouali, J. 1977. Contribution á la biologie des Scyliorhinidae des côtes tunisiennes. VI.- Galeus melastomus Rafinesque, 1810. Répartition géographique et bathymétrique, sexualité, reproduction, fécondité. Cah. Biol. Mar. 18: 449-463
- Carpine, C. 1970a. Ecologie de l'étage bathyal dans la Méditerranée occidentale. Mém. Inst. océanogr. (Monaco) 2: 1-146.
- Carpine, C. 1970b. Une expérience du chalutage profond (recherche de la "Caravelle" engloutie au large de Nice). Bull. Inst. océanogr. 69(1408): 16 pp.
- Carrassón, M., Matallanas, J. 1990. Preliminary data about the feeding habits of some deep-sea Mediterranean fishes. J. Fish Biol. 36: 461- 463
- Carrassón, M., Stefanescu, C. y Cartes, J. E. 1992. Diets and bathymetric distributions of two bathyal sharks of the Catalan sea (western Mediterranean). Mar. Ecol. Prog. Ser. 82: 21-30.
- Cartes, J. 1991. Análisis de las comunidades y estructura trófica de los crustáceos decápodos batiales del mar catalán. Tesis Doctoral. Universidad Politécnica de Cataluña. 627 pp.
- Casanova, B. 1974. Les euphausiacés de la Méditerranée (Systématique et développement larvaire. Biogéographie et Biologie) These. 380 pp.
- Caswell, H. 1978. Predator-mediated coexistence: a nonequilibrium model. Amer. Net. 112: 127-154.
- Cerro, del Ll. y Portas, F. 1984. Contribució al coneixement de la ictiofauna de la comarca del Garraf (Catalunya). Butll. Soc. Cat. Ictio. Herp. 6:4-25.
- Clarke, M.R., Merrett, N.R. 1972. The significance of squid, whale and other remains from the stomachs of bottom living deep-sea fish. J. mar. biol. Ass. U.K. 52: 599-603.

- Cocker, J.E. 1978. Adaptations of deep sea fishes. *Env. Biol. Fish.* Vol.3, No 4: 389-399.
- Cohen, D. M. 1986. Family Moridae: 713-723. *In: Fishes of the North-eastern Atlantic and the Mediterranean. Vol 2.* Whitehead, P.J.P., Bauchot, M.-L, Hureau, J.-C, Nielsen, J., Tortonese, E. (eds.). UNESCO. Bungay.
- Collet, R. 1896. Poissons provenant des campagnes du yacht l'"Hirondelle" (1885-1888). *Result. Camp. scient. Prince Albert I,* 10: VIII + 198 pp.
- Colom, G. 1974. Foraminíferos ibéricos. *Investigación Pesquera,* Tomo 38 (1): 1-245.
- Compagno, L.J.V. 1984. *FAO species catalogue. Vol. 4. Sharks of the world. An annotated and illustrated catalogue of sharks species Known to date. Part 1. Hexanchiformes to Lamniformes.* *FAO Fish. Synop. (125) Vol.4, Pt. 1:* 249 pp.
- Crabtree, R.E., Sulak, K.J., Musick, J.A. 1985. Biology and distribution of species of *Polyacanthonotus* (Pisces: Notacanthiformes) in the western North Atlantic. *Bull. mar. Sci.* 36: 235-248
- Chevreux, E., Fage, L. 1925. *faune de France. 9. Amphipodes.* Académie des Sciences. Paris.
- Childress, J.J, Gluck, D.L., Carney, R.S., Gowing, M.M. 1989. Benthopelagic biomass distribution and oxygen consumption in a deep-sea benthic boundary layer dominated by gelatinous organisms. *Limnol. Oceanogr.,* 34(5): 913-930.
- Daniels, R.A. 1982. Feeding ecology of some fishes of the antarctic peninsula. *Fish. Bull. H.S.* 80: 575-588
- Dayton, P. K. y Hessler, R. R. 1972. Role of biological disturbance in maintaining diversity in the deep-sea. *Deep-Sea Res.* 19: 199-208.

- De Groot, S.J. 1971. On the interrelationships between morphology of the alimentary tract, food and feeding behaviour in flatfishes (Pisces: Pleuronectiformes). *Neth. J. Sea Res.* 5: 121-196
- Della Croce, N., Drago, N. y Flocchini, G. 1988. Ricerche biologiche e geofisiche campagna oceanografica N/R "Minerva" (14-31/8/1987). Università di Genova. Istituto di Science Ambientali Marine, *Rapp. Tecnico*, 24:12 pp.
- Du Buit, M.-H. 1978. The diet of some teleost fishes in the Wyville Thomson Ridge area. *Oceanol. Acta* 1: 129-134
- Estrada, M., Vives, F. y Alcaraz, M. 1985. Life and the productivity of the open sea: 148-197. *In: Key environments. Western Mediterranean.* Margalef, R. (ed.). Pergamon Press Ltd. Oxford.
- Fage, L. 1951. *Faune de France*. 54. Cumacés. Paris. 136 pp.
- Fänge, R. y Grove, D. 1979. Digestion. *In Fish physiology*. Vol VIII: 161-204. Hoar, W. S., Randall, D. J. y Brett, J. R. (eds.). Academic press. New York.
- Fishelson, L. 1980. Partitioning and sharing of space and food resources by fishes. P. 415-445. *In: Fish behavior and its use in the capture and culture of fishes.* ICLARM Conf. Proc. S. J. E. Bardach, J. J.; Magnuson, J. J.; May, R. C. y Reinhart, J. M. (eds.). Int. Conf. Living Aquat. Resour., Manila.
- Flos, J. 1985. The driving machine: 60-99. *In: Key environments. Western Mediterranean.* Margalef, R. (ed.). Pergamon Press Ltd. Oxford.
- Foxton, P. 1970. The vertical distribution of pelagic decapods (Crustacea: Natantia) collected on the SOND Cruise 1965. II. The Penaeidea and general discussion. *J. Mar. Biol. Ass. U. K.* 50: 961-1000.

- Foxton, P. 1971/72. Observations on the Vertical Distribution of the Genus Acantheephyra (Crustacea: Decapoda) in the eastern North Atlantic, with particular reference to Species of the "purpurea" Group. Proc. R. Soc. Edinb. (B) 73(30): 301-313.
- Franqueville, C. 1971. Macroplancton profond (Invertébrés) de la Méditerranée nord-occidentale. Tethys, 3(1): 11-56.
- Gage, J. D., Tyler, P. A. 1991. Deep sea biology. A natural history of organism at the deep-sea floor. Cambridge University Press, Cambridge, England. 504 pp.
- Gascon, D. y Legget, W. C. 1977. Distribution, abundance and resource utilization of littoral zone fishes in response to a nutrient/production gradient in Lake Memphremagog. J. Fish. Res. Bd Can. 34: 1105-1117.
- Gause, G. F. 1935. Experimental demonstration of Volterra's periodic oscillation in the numbers of animals. J. exper. Biol. 12:44-48.
- Geevarghese, C. 1983. Morphology of the alimentary tract in relation to diet among gobioid fishes. J. nat. Hist. 17: 731-741
- Geistdoerfer, P. 1978. Ecology alimentaire des Macrouridae. Rev. Trav. Inst. scient. tech. Pêch. marit. 42: 177-260
- Geistdoerfer, P. 1978-79. Connaissances s sur la biologie de la famille des Macrouridae (Téléostéens Gadiformes). Vie Milieu (2): 323-351.
- Geistdoerfer, P. 1981. Morphologie et histologie de l'appareil digestif des Macrouridae (téléostéens). I: Morphologie de l'appareil digestif. Cybium 5: 3-44
- Geistdoerfer, P. 1986. Family Macrouridae: 644-676. In: Fishes of the north-eastern Atlantic and the Mediterranean. Vol. 2. Whitehead, P. J. P., Bauchot, M. L., Hureau, J. C., Nielsen, J. y Tortonese, E. (eds.). UNESCO. Bungay.

- Geistdoerfer, P. y Rannou, M. 1972. Poissons benthiques récoltés en Méditerranée occidentale par le N. O. "Jean Charcot" (campagne Polymède). Bull. Mus. natn. Hist. nat., 3ème sér. 25: 101-110.
- Giglioli, E. H. 1881. La scoperta di una fauna abissale nel Mediterraneo. Nature 24: 381.
- Giglioli, E. H. 1882. New deep-sea fish from the Mediterranean. Nature 27: 198-199.
- Giglioli, E. H. 1893. Di una nuova specie di Macrourid appartenente alla fauna abissale del Mediterraneo. Zool. Anz. 16: 343-345.
- Golani, D. 1986/87. On deep-water sharks caught off the Mediterranean coast of Israel. Isr. J. Zool. 34(1-2): 23-31.
- Goldstein, M.E. 1986. Características morfológicas del sistema digestivo y hábitos alimentarios de la brótola (Urophycis brasiliensis) (Pisces: Gadidae). Publ. Com. Tec. Mix. Fr. Mar. 1: 351-368
- Golovan, G.A., Pakhorukov, N.P. 1980. New data on the ecology and morphometry of Alepocephalus rostratus (Alepocephalidae). J. Ichthyol. 20: 77-84
- Goodyear, R.H., Zahuranec, B.J., Pugh, L.W., Gibbs, R.H.Jr. 1972. Ecology and vertical distribution of Mediterranean midwater fishes. In: Mediterranean Biological Studies, Final Report, Part 3. Smithsonian Institution, Washington.
- Gordon, J. D. M. 1986. The fish populations of the Rockall Trough. Proceedings of the Royal Society of Edimburg, 88 B: 191-204.
- Gordon, J. D. M., Duncan, J. A. R. 1985. The ecology of the deep-sea benthic and benthopelagic fish of the slopes of the Rockall Trough, Northeastern Atlantic. Progress in Oceanography, 15: 37-69.
- Gordon, J. D. M., Duncan, J. A. R. 1987. Deep-sea bottom-living

- fishes at two repeat stations at 2200 and 2900 m in the Rockall Trough, northeastern Atlantic Ocean. *Mar. Biol.* 96: 309-325
- Grey, M. 1953. Fishes found below a depth of 2000 meters. XIV Intern. Zool. Congr. Copenhagen.
  - Grey, M. 1956. The distribution of fishes found below a depth of 2000 m. *Fieldiana, Zool.* 36(2): 75-183.
  - Haedrich, R.L., Rowe, G.T., Polloni, P.T. 1980. The megabenthic fauna in the deep-sea south of New England, USA *Mar. Biol.*, 57: 165-179.
  - Harder, W. 1975. Anatomy of fishes. Part I. E. Schweizer-bartsche Verlagsbuchhandlung. Nägele u. Obermiller (ed.). Stuttgart. 612 pp.
  - Hardin, G. 1960. The competitive exclusion principle. *Science*, 131: 1292-1297.
  - Hargreaves, P.M., Ellis, C.J., Angel, M.V. 1984. An assessment of biological processes close to the sea bed in a slope region and its significance to the assessment of sea-bed disposal of radioactive waste. I.O.S. Report, 1985: 121 pp.
  - Hayward, P.J., Ryland, J.S. 1990. The marine fauna of the British isles and North-West Europe. Vol 1 y 2. Oxford Science Publications. Oxford.
  - Hopkins, T. S. 1985. Physics of the sea: 100-125. *In: Key environments. Western Mediterranean.* Margalef, R. (ed.). Pergamon Press Ltd. Oxford.
  - Hureau, J.-C. 1970. Biologie comparée de quelques poissons antarctiques (Nototheniidae). *Bull. Inst. océanogr.* 68(1391): 1-250.
  - Hureau, J.C., P. Geistdoerfer & M. Rannou. 1979. The ecology of deep-sea benthic fishes. *Sarsia* 64: 103-108.

- Hutchinson, G. E. 1959. Homage to Santa Rosalia, or why are there so many kinds of animals?. *Amer. Natur.* 93: 145-159.
- Hynes, H. B. N. 1950. The food of freshwater sticklebacks (*Gasterosteus aculeatus* and *Pygosteus pungitius*) with a review of methods used in studies of the food of fishes. *J. an. Ecol.* 19: 36-58.
- Hyslop, E.J. 1980. Stomach contents analysis- a review of methods and their application. *J. Fish Biol.*, 17: 411-429.
- Ibañez, M. 1975. Notas ictiológicas II: sobre la especie Lepidion eques Gunther, 1887 (Moridae) en aguas del Golfo de Vizcaya. *Munibe*, 3-4: 207-213.
- Ivlev, V.S. 1961. *Experimental Ecology of the Feeding of Fishes.* Yale Univ. Press, New Haven, Conn. USA. 302 p.
- Jacobshagen, E. 1913. Untersuchungen über das Darmsystem der Fische und Dipnoer II. *Jena Z. Naturw.* 49: 373- 810
- Jones, N. S. 1976. *British Cumaceans.* Academic Press London and New York. 86 pp.
- Jumars, P. A. y Gallagher, E. 1982. Deep-sea community structure: three plays on the benthic proscenium: 217-255. *In: The environment of the deep-sea.* Ernst, W. G. y Morin, J. G. (eds.). Prentice-Hall. New jersey.
- Kaim-Malka, R. A. 1969(70). Biologie y écologie de quelques *Ampelisca* (Crustacea. Amphipoda) de la région de Marseille. *Tethys.* 1(4): 977-1022.
- Kapoor, B. G. 1953. The anatomy and histology of the alimentary canal in relation to the feeding habits of a siluroid fish. *Wallago attu.* *J. Zool. Soc. India.* 5: 191-210.
- Kapoor, B.G., Smith, H., Verghina, I.A. 1975. The alimentary canal and digestion in teleosts. *Adv. mar. Biol.* 13: 109-239

- Kawaguchi, K. y Mauchline, J. 1982. Biology of myctophid fishes (Family Myctophidae) in the Rockall Trough, northeastern Atlantic. *Biol. Ocean.* 1: 337-373.
- Keast, A. 1978. Feeding interrelations between age-groups of pumpkinseed (*Leponis gibbosus*) and comparisons with bluegill (*L. macrochirus*). *J. Fish. Res. Bd Can.* 35: 12-27.
- Klausewitz, W. 1989. Deep-sea and deep water fish of the eastern Mediterranean, collected during the METEOR expedition 1987. *Senckenbergiana marit.* 20 (5/6):251-263.
- Lagarde, J. 1983. *Initiation à l'analyse des donnés.* Dunod. Paris. 157 pp.
- Lagardère, J. P. 1971. Les crevettes des côtes du Maroc. *Trav. Inst. Sci. Cher. Fac. Sci.* 36: 140 pp.
- Lagardère, J.-P 1977. Recherches sur la distribution verticale et sur l'alimentation des crustacés décapodes benthiques de la pente continentale du golfe de Gascogne. Analyse des groupements carcinologiques. *Bull. Cent. Etud. Rech. sci., Biarritz* 11(4): 367-440.
- Laurens, J. Menzies, R.J., Bacescu, M.C. 1962. *Abyssal crustacea.* N 1. Columbia University Press, New York & London.
- Ledoyer, M. 1977. Contribution à l'étude de l'écologie de la faune vagile profonde de la Méditerranée nord occidentale. I. Les Gammariens (Crustacea, Amphipoda). *Boll. Mus. Civ. St. Nat. Verona*, IV: 321-421.
- Ledoyer, M. 1983. Contribution à l'étude de l'écologie de la faune vagile profonde de la Méditerranée nord occidentale. II. Les cumacés (Crustacea). *Tethys.* 11(1): 67-81.
- Lozano Cabo, F. 1952. Estudio preliminar sobre la biometría, la biología y la anatomía general de Notacanthus bonapartei. *Bol. Inst.*