

Prog. Ocean. 13: 353-382.

- Roe, H. S. J., James, P. T. y Thurston, M. H. 1984. The diel migrations and distributions within a mesopelagic community in the North East Atlantic. 6. Medusae, Ctenophores, Amphipodes and Euphausiids. Prog. Oceanog. 13: 425-460.
- Roper, C.F.E. 1972. Ecology and vertical distribution of Mediterranean pelagic cephalopods. In: Mediterranean Biological Studies, Final Report, Part 5. Smithsonian Institution, Washington.
- Roper, C. F. E. y Young, R. E. 1975. Vertical distribution of pelagic cephalopods. Smithson. Contr. Zool. 209: 1-51.
- Rosecchi, E., Nouaze, Y. 1987. Comparaison de cinq indices alimentaires utilisés dans l'analyse des contenus stomachaux. Rev. Trav. Inst. Pech. mar. 49(3-4): 111-123.
- Roughgarden, J. 1974. Species packing and the competition function with illustrations from coral reef fish. Theor. Populat. Biol. 5: 163-186.
- Roughgarden, J. 1976. Resource partitioning among coexisting species - a coevolutionary approach. Theoret. Pop. Biol. 9: 388-424.
- Roule, L. 1919. Poissons provenant des campagnes du yacht "Princesse Alice" (1891-1913) et du yacht l'"Hirondelle II" (1914). Résult. Camp. scient. Prince Albert I, 52: 191 pp.
- Salat, J. y Font, J. 1985. Masses d'aigua i circulació a la Mediterrània. Quad. Ecol. Apl. 8: 61-98.
- Saldanha, L. 1986. Family Nettastomatidae: 562-566. In: Fishes of the North-eastern Atlantic and the Mediterranean. Vol 2. Whitehead, P.J.P., Bauchot, M.-L, Hureau, J.-C, Nielsen, J., Tortonese, E. (eds.). UNESCO. Bungay.
- Saldanha, L. 1988. Notes sur le régime alimentaire de deux poissons

de profondeur: *Alepocephalus bairdii* et *Bathypterois dubius*. Cybium, 12(2): 169-171.

- Sale, P. G. 1979. Habitat partitioning and competition in fish communities. p. 323-331. In: Predator-prey systems in fisheries management. H. E. Clepper (ed.). Sport Fishing Institute, Washington, D. C.
- Schmidt, J. 1912. Introduction: 1-51. In: Rep. Dan. Oceanogr. Exped. Mediterr. 1: 1-149.
- Schoener, T.W. 1970. Non-synchronous spatial overlap of lizards in patchy habitats. Ecology, 51: 408-418.
- Schoener, T.W. 1974. Resource partitioning in ecological communities. Science 185: 27-39.
- Sedberry, G.R., Musick, J.A. 1978. Feeding strategies of some demersal fishes of the continental slope and rise off the mid-Atlantic coast of the USA. Mar. Biol. 44: 357-375.
- Serra, J. 1981. Els canyons submarins del marge continental català. Treb. Inst. Cat. Hist. Nat. 9: 63-67.
- Shannon, C.E., Weaver, W. 1963. The Mathematical Theory of Communication. University of Illinois Press, Urbana.
- Silverberg, N., Edenborn, H.M., Ouellet, G., Bland, P. 1987. Direct evidence of a mesopelagic fish, Melanostigma atlanticum (Zoarcidae), spawning within bottom sediments. Envir. Biol. Fish. 20: 195-202.
- Smith, D. G., Bohlke, J. E. y Castle, P. H. J. 1981. A revision of the Nettastomid eel genera *Nettastoma* and *Nettenchelis* (Pisces: Anguilliformes), with description of six new species. Proc. Biol. Soc. Wash. 94(2): 535-560.
- Smith, K.L. 1982. Zooplankton of a bathyal benthic boundary layer: In situ rates of oxygen consumption and ammonium excretion.

- Limnol. Oceanogr. 27: 461-h471.
- Sokal, R.R., Rohlf, F.F. 1969. Biometry. The principles and practice of statistics in biological research. W.H. Freeman & Co., San Francisco.
 - Somero, G. N., Siebenaller, J. F. y Hochachka, P. W. 1983. Physiological adaptations of deep-sea animals: 261-330. In: Deep-sea biology, The Sea. Vol 8. Rowe, G. T. (ed.). John Wiley y Sons. New York.
 - Sorbe, J.C. 1972. Ecologie et ethologie alimentaire de l'ichthyofaune chalutable du plateau continental Sud Gascogne. Tesis. Université d'Aix- Marseille.
 - Sorbe, J.C. 1977. Régime alimentaire de *Phycis blennoides* (Brünnich, 1768) dans le sud du golfe de Gascogne. Rev. Trav. Inst. Pêches mar. 41(3): 271-289.
 - Sorbe, J. C. 1987. Structural evolution of two suprabenthic softbottom communities of the South Gascogne continental shelf. 22nd European Marine Biology Symposium, Barcelona: 1-20.
 - Springer, S. 1967. Social organization of sharks populations. In: Gilbert, R.F., Matheuson, R.F., Rall, D.P. (ed.). Sharks, Skates and Rays. John Hopkins Press, Standford.
 - Stefanescu, C. 1991. Comunidades ictiológicas demersales del mar catalán (Mediterráneo noroccidental) por debajo de los 1000 m de profundidad. Tesis Doctoral. Universidad de Barcelona. 490 pp.
 - Sulak, K. J. 1977. The systematics and biology of Bathypterois (Pisces, Chlorophthalmidae). Galathea Rep. 14: 49-108.
 - Sulak, K. J. 1984a. A comparative ecological analysis of temperate and tropical demersal deep-sea fish faunas in the Western North Atlantic. 211 pp., Adaptation of Ph. D. Dissertation, University of Miami, Miami.

"Princesse Alice". Résult. Camp. scient. Prince Albert I, 35: 174 pp.

Washington, D. C.

- Windell, J. T. y Bowen, S. H. 1978. Methods for study of fish diets based on analysis of stomach contents: 227-254. In: Methods for assessment of fish production in fresh waters. Bagena, t. (ed.). Blackwell Scientific Publications. Oxford.
- Wishner, K.F. 1980. The biomass of the deep-sea benthopelagic plankton. Deep-Sea Res. 27A: 203-216.
- Wurtz, M., Vacchi, M. 1978. Ricerca di cicli nittemerali nell'alimentazione di Selaci bartiali. Quad. Lab. Tecnol. Pesca 3(suppl.): 155-164).
- Yano, K. y Tanaka, S. 1984. Some biological aspects of the deep-sea squaloid shark *Centroscymnus* from Suruga Bay, Japan. Bull. Jap. Soc. Sci. Fish. 50(2): 249-256.
- Yano, K. y Tanaka, S. 1988. Size at Maturity. Reproductive Cycle, Fecundity, and Depth Segregation of the Deep Sea Squaloid Sharks *Centroscymnus owstoni* and *C. coelolepis* in Suruga Bay, Japan. Nippon Suisan Gakkaishi 54(2): 167-174.
- Yasuda, F. 1960. The types of food habits of fishes assured by stomach contents examination. Bull. Jap. Soc. Sci. Fish. 26(7): 653-662.
- Zaret, T. M. y Rand, A. S. 1971. Competition in tropical stream fishes: support for the competitive exclusion principle. Ecolgy. 52: 336-342.
- Zarquiey, R. 1968. Crustáceos decápodos ibéricos. Investigación pesquera, tomo 32: 1-510.
- Zenkevich, L.A. & J.A. Birstein. 1956.- Studies of the deep water fauna and related problems. Deep-sea Research, Vol 4, pp. 54-64.
- Zugmayer, E. 1911. Poissons provenant des campagnes du yacht

- Templeman, W. 1970. A review of the morid fish genus Lepidion of the North Atlantic with first records of Lepidion eques from the Western North Atlantic. *J. Fish. Res. Board Can.* 27: 457-498.
- Thiel, H. 1983. Meiobenthos and nanobenthos of the deep-sea: 167-230. In: Deep-sea biology, The sea. Vol 8. Rowe, G. T. (ed.). John Wiley y Sons. New York.
- Tiganus, V. 1986. Distribution qualitative et quantitative des peuplements d'Amphipodes du substrat meuble sur le plateau continental roumain. *Cercetari Marine. Rech. Mar.* 19: 91-108.
- Tilman, D., Mattson, M. y Lauger, S. 1981. Competition and nutrient kinetics along a temperature gradient: an experimental test of a mechanistic approach to niche theory. *Limnol. Oceanogr.* 26: 1020-1033.
- Trégouboff, G., Rose, M. 1957. Manuel de planctonologie Méditerranéenne. Tome I. Tome II. Centre National de la Recherche Scientifique. Paris.
- Vegas-Vélez, M. 1972. Contribution à l'étude de l'appareil digestif des téléostéens. Rapports avec l'éthologie alimentaire. Thèse Troisième Cycle. Université Aix, Marseille.
- Venec-Peyré, M. T. 1985. Etude préliminaire des foraminifères benthiques des sédiments profonds. In: Peuplements profonds du Golfe du Gascogne. IFREMER: 255-262.
- Voss, G. L. 1967. The biology and bathymetric distribution of deep-sea cephalopods. *Stud. trop. Oceanogr. Miami.* 5: 511-535.
- Wallace, R. K. 1981. An assessment of diet overlap indexes. *Trans. Am. Fish. Soc.* 110: 72-76.
- Werner, E. E. 1979. Niche partitioning by food size in fish communities. P. 311-322. In: Predator-prey systems in fisheries management. H. E. Clepper (ed.). Sport Fishing Institute,

- Sulak, K. J. 1984b. Family: Chlorophthalmidae (including Bathypteroideae, Benthosauridae, Ipnopidae): 412-420. In: Fishes of the North-eastern Atlantic and the Mediterranean. Vol 1. Whitehead, P.J.P., Bauchot, M.-L, Hureau, J.-C, Nielsen, J., Tortonese, E. (eds.). UNESCO. Bungay.
- Sulak, K. J. 1986. Family: Notacanthidae: 599-603. In: Fishes of the North-eastern Atlantic and the Mediterranean. Vol 2. Whitehead, P.J.P., Bauchot, M.-L, Hureau, J.-C, Nielsen, J., Tortonese, E. (eds.). UNESCO. Bungay.
- Suyehiro, Y. 1942. A study on the digestive system and feeding habits of fish. Jap. J. Zool. 10: 1-303
- Svetovidov, A. 1934. On the correlation between the character of food and the number of pyloric caeca in fishes. Dokl. Akad. Nauk SSSR 3: 70-72.
- Svetovidov, A. N. 1953. Corrélations entre le nombre de caecums pyloriques et le comportement alimentaire des poissons. In: Essais sur les grands problèmes d'ichthyologie. Acad. Sienc. U.R.S.S.
- Svetovidov, A. N. 1986. Family Gadidae: 680-710. In: Fishes of the North-eastern Atlantic and the Mediterranean. Vol 2. Whitehead, P.J.P., Bauchot, M.-L, Hureau, J.-C, Nielsen, J., Tortonese, E. (eds.). UNESCO. Bungay.
- Syvitski, J.P.M., Silverberg, N., Ouellet, G., Asprey, K.W. 1983. First observations of benthos and seston from a submersible in the lower St.Lawrence estuary. Geogr. Phys. Quater. 37: 227-240.
- Targett, T.E. 1981. Trophic ecology and the structure of coastal antarctic fish communities. Mar. Ecol. Prog. Ser. 4: 243-246.
- Tattersall; W.M., Tattersall, B.S. 1951. The British Mysidacea. Adlar y Sm, Ltd. London.