

VIII.- Bibliografía

VIII.- Bibliografía.

- Almouzni, G. and Méchali, M. (1988). Assembly of spaced chromatin by DNA synthesis in extracts from *Xenopus* eggs. *EMBO J.*, **7**: 664-672.
- Amano, T.; Okita, Y. and Hoshi, M. (1992). Treatment of starfish sperm with egg jelly induces the degradation of histones. *Devel. Growth Differ.*, **34**: 99-106.
- Annunziato, A.T. (1989). Inhibitors of topoisomerases I and II arrest DNA replication, but do not prevent nucleosome assembly *in vivo*. *J. Cell. Sci.*, **93**: 593-603.
- Ausió, J. (1995). Histone H1 and the evolution of the nuclear sperm-specific proteins. In: Advances in spermatozoal phylogeny and taxonomy. B.G.M. Jamieson; J. Ausió and J.L. Justine (Eds.). Mém. Mus. natn. Hist. nat., París, **166**: 447-462.
- Ausió, J. and van Holde. (1987). A dual chromatin organization in the sperm of the bivalve mollusc *Spisula solidissima*. *Eur. J. Biochem.*, **165**: 363-371.
- Azorín, F.; Olivares, C.; Jordán, A.; Pérez-Grau, L.; Cornudella, L.; and Subirana, J.A. (1983). Heterogeneity of the Histone-containing Chromatin in Sea Cucumber Spermatozoa. Distribution of the Basic Protein ϕ_0 and Absence of Non-histone Proteins. *Exp. Cell Res.*, **148**: 331-344.
- Azorín, F.; Rocha, E.; Cornudella, L.; and Subirana, J.A. (1985). Anomalous Nuclease Digestion of *Holothuria* Sperm Chromatin Containing Histone H1 Variants. *Eur. J. Biochem.*, **148**: 529-532.
- Banerjee, S. and Cantor, C.R. (1990). Nucleosome assembly of simian virus 40 DNA in a mammalian cell extract. *Mol. Cell. Biol.*, **10**: 2863-2873.
- Balakier, H. And Tarkowski, A.K. (1980). The role of germinal vesicle karyoplasm in the development of male pronucleus in the mouse. *Exper. Cell Res.*, **128**: 79-85.
- Barnes, R.D. (1980). Invertebrate Zoology. 4 th. Edition. Saunders College, Philadelphia, USA.
- Barone, J.R.; De Lara, J.; Cummings, K.B. and Ward, W.S. (1994). DNA organization in human spermatozoa. *J. Androl.*, **15**: 139-144.
- Barry, J.M.; and Merriam, R.W. (1972). Swelling of hen erythrocyte nuclei in cytoplasm from *Xenopus* eggs. *Experimental Cell Research*, **71**: 90-96.
- Becker, P.B. and Wu, C. (1992). Cell-free system for assembly of transcriptionally repressed chromatin from *Drosophila* embryos. *Mol. Cell. Biol.*, **12**: 2241-2249.
- Bellvé, A.R.; Chaudrika, R.C.; Martinova, Y. and Barth, A. (1992). The perinuclear matrix as a structural element of the mouse sperm nucleus. *Devel. Biol.*, **47**: 451-465.
- Benavente, R. and Krohne, G. (1985). Change of karyoskeleton during spermatogenesis of *Xenopus*: Expression of lamin IV, a nuclear lamina protein specific for the male germ line. *Proc. Natl. Acad. Sci., USA*, **82**: 6176-6180.
- Berrios, M. and Bedford, J.M. (1976). Oocyte maturation: Aberrant postfusion responses of the rabbit primary oocyte to penetrating spermatozoa. *J. Cell Sci.*, **39**: 1-12.
- Berrios, M. and Avilón, A.A. (1990). Nuclear formation in a *Drosophila* cell-free system. *Exp. Cell Res.*, **191**: 64-70.

- Bloch, D.P. (1969). A catalog of sperm histones. *Genetics Suppl.*, **61**: 93-111.
- Bloch, D.P. and Hew, H.Y.C. (1960). Changes in nuclear histones during fertilization and early embryonic development in the pulmonate snail, *Helix aspersa*. *J. Biophys. Biochem. Cytol.*, **8**: 69-81.
- Blow, J.J. and Laskey, R.A. (1988). A role for the nuclear envelope in controlling DNA replication within the cell cycle. *Nature*, **332**: 546-548.
- Blum, H.; Beier, H; and Gross, H.J. (1987). Improved silver staining of plant proteins, RNA and DNA in polyacrylamide gels. *Electrophoresis*, **8**: 93-99.
- Bols, N.C. and Kasinsky, H.E. (1973). An electrophoretic comparison of histones in anuran testes. *Can. J. Zool.*, **51**: 203-208.
- Bonne-Andrea, C.; Harper, F.; Sobczak, J. and DeRecondo, A.-M. (1984). Rat liver HMG1: a physiological nucleosome assembly factor. *EMBO J.*, **3**: 1193-1199.
- Bonne-Andrea, C.; Wong, M.L. and Alberts, B. (1990). In vitro replication through nucleosomes without histone displacement. *Nature*, **343**: 719-726.
- Braunstein, M.; Rose, A.B.; Holmes, S.G.; Allis, C.D. and Broach, J.R. (1993). Transcriptional silencing in yeast is associated with reduced histone acetylation. *Genes Dev.*, **7**: 592-603.
- Brown, D.B.; Blake, E.J.; Wolgemuth, D.J.; Gordon, K. And Ruddle, F.H. (1987). Chromatin decondensation and DNA synthesis in human sperm activated *in vitro* by using *Xenopus laevis* egg extracts. *J. Exper. Zool.*, **242**: 215-231.
- Brown, D.B.; Miskimins, W.K. and Ruddle, F.H. (1991). Partial purification of *Xenopus laevis* egg extract factor(s) that induce swelling in permeabilized human sperm. *J. Exp. Zool.*, **258**: 263-272.
- Brown, S.A.; Imbalzano, A.N. and Kingston, R.E. (1996). Activator-dependent regulation of transcriptional pausing on nucleosomal templates. *Genes Dev.*, **10**: 1479-1490.
- Brownell, J.E. and Allis, C.D. (1996). Special HATs for special occasions: linking histone acetylation to chromatin assembly and gene activation. *Curr. Opin. Genet. Dev.*, **6**: 176-184.
- Buesa, C.; del Valle, L.; Saperas, N.; Goethals, M.; Lloris, D. and Chiva, M. (1998). Primary structure of scombrine α : Two different species with an identical protamine. *Comp. Biochem. Physiol.*, **119B**: 145-149.
- Bulger, M.; Ito, T.; Kamakaka, T.T. and Kadonaga, J.T. (1995). Assembly of regularly spaced nucleosome arrays by *Drosophila* chromatin assembly factor 1 and a 56 kDa histone-binding protein. *Proc. Natl. Acad. Sci. USA*, **92**: 11726-11730.
- Buratowski, S. (1995). Mechanisms of gene activation. *Science*, **270**: 1773-1774
- Bürglin, T.R.; Mattaj, I.W.; Newmeyer, D.D.; Zeller, R. and De Robertis, E.M. (1987). Cloning of nucleoplasmin from *Xenopus laevis* oocytes and analysis of its developmental expression. *Genes Dev.*, **1**: 97-107.
- Burns, L.G. and Peterson, C.L. (1997). The yeast SWI-SNF complex facilitates binding of a transcriptional activator to nucleosomal sites *in vivo*. *Mol. Cell. Biol.*, **17**: 4811-4819.
- Busslinger, M.; and Barberis, A. (1985). Synthesis of sperm and late histone cDNA's of the sea urchin with a primer complementary to the conserved 3' terminal palindrome: Evidence for tissue-specific and more general histone gene variants. *Proc. Natl. Acad. Sci., USA*, **82**: 5676-5680.

- Calvin, H.I. and Bedford, J.M. (1971). Formation of disulphide bonds in the nucleus and accessory structures of mammalian spermatozoa during maturation in the epididymis. *J. Reprod. Fertil. Suppl.*, **13**: 65-75.
- Cameron, L.A. and Poccia, D.L. (1994). *In vitro* development of the sea urchin male pronucleus. *Devel. Biol.*, **162**: 568-578.
- Carrol, A.G. and Ozaki, H. (1979). Changes in the histones of the sea urchin *Strongylocentrotus purpuratus* at fertilization. *Cell Res.*, **119**: 307-315.
- Carron, C.P. and Longo, F.J. (1982). Relation of cytoplasmic alkanization to microvillar elongation and microfilament formation in the sea urchin egg. *Devel. Biol.*, **89**: 128-137.
- Cairns, B.R. (1998). Chromatin remodeling machines: similar motors, ulterior motives. *Trends Biochem. Sci.*, **23**: 20-25.
- Casas, M.T.; Mura, C.V.; Subirana, J.A.; and Cornudella, L. (1989). Purification and Immunocytolocalization of Protein ϕ_0 from Sperm Cells of the Echinoderm *Holothuria tubulosa*. *Exp. Cell Res.*, **182**: 14-25.
- Chaeveau, J.; Moule, Y. and Rouiller, C. (1956). Isolation of pure and unaltered liver nuclei morphology and biochemical composition. *Exp. Cell Res.*, **11**: 317-321.
- Chaudhary, N. and Courvalin, J.-C. (1993). Stepwise reassembly of the nuclear envelope at the end of mitosis. *J. Cell Biol.*, **122**: 295-306.
- Chiva, M.; Saperas, N.; Cáceres, C. and Ausió, J. (1995). Nuclear basic proteins from sperm of tunicates, cephalochordates, agnathans and fish. In: Advances in spermatozoal phylogeny and taxonomy. B.G.M. Jamieson; J. Ausió and J.L. Justine (Eds.). Mém. Mus. natn. Hist. nat., París, **166**: 501-514.
- Collas, P. and Poccia, D. (1995a). Formation of the sea urchin male pronucleus *in vitro*: Membrane-independent chromatin decondensation and nuclear envelope-dependent nuclear swelling. *Mol. Reprod. Devel.*, **42**: 106-113.
- Collas, P. And Poccia, D. (1995b). Lipophilic organizing structures of sperm nuclei target membrane vesicle binding and are incorporated into the nuclear envelope. *Devel. Biol.*, **169**: 123-125.
- Collas, P.; Pinto-Correia, C. and Poccia, D. (1995). Lamin dynamics during sea urchin male pronucleus formation *in vitro*. *Exper. Cell Res.*, **219**: 687-698.
- Conaway, R.C. and Conaway, J.W. (1993). General initiation factors for RNA polymerase II. *Ann. Rev. Biochem.*, **62**: 161-190.
- Cornudella, L.; and Rocha, E. (1979). Nucleosome Organization during Germ Cell Development in the Sea Cucumber *Holothuria tubulosa*. *Biochemistry*, **18**: 3724-3732.
- Cotten, M.; Sealy, L. and Chalkley, R. (1986). Massive phosphorylation distinguishes *Xenopus laevis* nucleoplasmin isolated from oocytes or unfertilized eggs. *Biochemistry*, **25**: 5063-5069.
- Cotten, M. and Chalkey, R. (1987). Purification of a novel, nucleoplasmin-like protein from somatic nuclei. *EMBO J.*, **6**: 3945-3954.
- Cothren, C.C. and Poccia, D.L. (1993). Two steps required for male pronucleus formation in the sea urchin egg. *Exp. Cell Res.*, **205**: 126-133.
- Crevel, G. and Cotterill, S. (1995). DF-31, a sperm decondensation factor from *Drosophila melanogaster*: purification and characterization. *EMBO J.*, **14**: 1711-1717.

- Crevel, G.; Huikeshoven, H.; Cotterill, S.; Simon, M.; Wall, J.; Philpott, A.; Laskey, R.A.; McConnell, M.; Fisher, P.A. and Berrios, M. (1997). Molecular and cellular characterization of CRP1, a *Drosophila* chromatin decondensation protein. *J. Structural Biology*, **118**: 9-22.
- Das, N.K.; Micou-Eastwood, J. and Alfert, M. (1975). Cytochemical studies on the protamine-type protein transition in sperm nuclei after fertilization and the early embryonic histones of *Urechis caupo*. *Devel. Biol.*, **43**: 333-339.
- Dasso, M.; Dimitrov, S.; and Wolffe, A.P. (1994). Nuclear assembly is independent of linker histones. *Proc. Natl. Acad. Sci. USA*, **91**: 12477-12481.
- Deguchi, R. and Osanai, K. (1994a). Repetitive intracellular Ca^{2+} increases at fertilization and the role of Ca^{2+} in meiosis reinitiation from the first metaphase in oocytes of marine bivalves. *Devel. Biol.*, **163**: 162-174.
- Deguchi, R. and Osanai, K. (1994b). Meiosis reinitiation from first prophase is dependent on the levels of intracellular Ca^{2+} and pH in oocytes of the bivalves, *Mactra chinensis* and *Limaria hakodatensis*. *Devel. Biol.*, **166**: 587-599.
- Deguchi, R. and Osanai, K. (1995). Serotonin-induce meiosis reinitiation from first prophase and from first metaphase in oocytes of the marine bivalve *Hiatella flaccida*: Respective changes in intracellular Ca^{2+} and pH. *Devel. Biol.*, **171**: 483-496.
- Del Valle, L.; Giménez, P.; Saperas, N. and Chiva, M. (1997). Empaquetament del DNA en els espermatozoides. En: Dinàmica estructural de macromolècules. Ed. F. Artigas. *Treballs de la SCB*, **48**: In Press
- Dimitrov, S.; Almouzni, G.; Dasso, M. and Wolffe, A.P. (1993). Chromatin transitions during early *Xenopus* embryogenesis: changes in histone H4 acetylation and in linker histone type. *Develop. Biol.*, **160**: 214-227.
- Dimitrov, S.; Dasso, M.C. and Wolffe, A.P. (1994). Remodeling sperm chromatin in *Xenopus laevis* egg extracts: the role of core histone phosphorylation and linker histone B4 in chromatin assembly. *J. Cell Biol.*, **126**: 591-601.
- Dimitrov, S.; and Wolffe, A.P. (1996). Remodeling somatic nuclei in *Xenopus laevis* egg extracts: molecular mechanisms for the selective release of histone H1 and H1 α from chromatin and the acquisition of transcriptional competence. *The EMBO Journal*, **15**: 5897-5906.
- Dingwall, C.; Sharnick, S.V. and Laskey, R.A. (1982). A polypeptide domain that specifies migration of nucleoplasmin into the nucleus. *Cell*, **30**: 449-458.
- Dingwall, C. and Allan, J. (1984). Accumulation of the isolated carboxy-terminal domain of histone H1 in the *Xenopus* oocyte nucleus. *EMBO J.*, **3**: 1933-1937.
- Dingwall, C.; Dilworth, S.M.; Black, S.J.; Kearsey, S.E.; Cox, L.S. and Laskey, R.A. (1987). Nucleoplasmin cDNA sequence reveals polyglutamic acid tracts and a cluster of sequences homologous to putative nuclear localization signals. *EMBO J.*, **6**: 69-74.
- Dingwall, C.; Robbins, J.; Dilworth, S.M.; Roberts, B. and Richardson, W.D. (1988). The nucleoplasmin nuclear location sequence is larger and more complex than that of SV40 large T antigen. *J. Cell Biol.*, **107**: 841-849.
- Dingwall, C.; Robbins, J. and Dilworth, S.M. (1989). Characterisation of the nuclear location sequence of *Xenopus* nucleoplasmin. *J. Cell Sci. (Suppl.)*, **11**: 243-248.
- Doonan, S. (1996). Chromatography on hydroxyapatite. In: Protein Purification Protocols. Ed. S. Doonan. Humana Press Inc., Totowa, NJ, USA. Methods in Molecular Biology, Vol. 59, Chapter 21: 211-215.

- Drabent, B.; Louroutziatis, A.; Prats, E.; Cornudella, L. and Doenecke, D. (1995). Structure of histone H2B and H4 genes of the sea cucumber *Holothuria tubulosa*. *The Journal of Sequencing and Mapping*, **6**: 41-45.
- Dreyer, C.; Stick, R. and Hausen, P. (1986). Uptake of oocyte nuclear proteins by nuclei of *Xenopus* embryos. In: Proceeding of the workshop on nucleocytoplasmic transport. Eds. R. Peters and M.F. Trendelenburg. Springer-Verlag, Berlin. pp. 143-157.
- Dubé, F.T.; Schmidt, T.; Johnson, C.H. and Epel, D. (1985). The hierarchy of requirements for an elevated intracellular pH during early development of sea urchin embryos. *Cell*, **40**:657-666
- Dubé, F. (1988). The relationship between early ionic event, the pattern synthesis, and oocyte activation in the surf clam, *Spisula solidissima*. *Devel. Biol.*, **126**: 233-241.
- Dubé, F. and Couto, L. (1990). The sodium requirement for activation of surf clam oocytes . *Cell Biol. Int. Rep.*, **14**: 463-471.
- Earnshaw, W.C.; Honda, B.M. and Laskey, R.A. (1980). Assembly of nucleosomes: The reaction involving *Xenopus laevis* nucleoplasmin. *Cell*, **21**: 373-383.
- Ecklund, P.S. and Levine, L. (1975). Mouse sperm basic nuclear protein, electrophoretic characterization and fate after fertilization. *J. Cell Biol.*, **66**: 251-262.
- Edmondson, D.G. and Roth, S.Y. (1996). Chromatin and transcription. *Faseb J.*, **10**: 1173-1182.
- Eng, L.A. and Metz, C.B. (1980). Sperm head decondensation by a high molecular weight fraction of sea urchin egg homogenates. *J. Exper. Zool.*, **212**: 159-167.
- Epel, D. (1978). Mechanism of activation of sperm and egg during fertilization of sea urchin gametes. *Curr. Topics Devel. Biol.*, **12**: 185-246.
- Farrants, O.; Blomquist, P.; Kwon, H. and Wrangle, O. (1997). Glucocorticoid receptor-glucocorticoid response element binding stimulates nucleosome disruption by the SWI/SNF complex. *Mol. Cell. Biol.*, **17**: 895-905.
- Franke, W.W. (1974). Structure, biochemistry and functions of the nuclear envelope. *Int. Rev. Cytol. Suppl.*, **4**: 72-236.
- Garagna, S. and Redi, C.A. (1988). Chromatin topology during the transformation of the mouse sperm nucleus into pronucleus *in vivo*. *J. Exper. Zool.*, **246**: 187-193.
- Garbers, D.L.; Ramarao, C.S.; Shimomura, H. and Bently, J.K. (1986). The regulation of spermatozoa by egg peptides. *Develop. Growth and Differ.* **28(suppl)**: 15-16.
- Georgel, P.T.; Tsukiyama, T. and Wu, C. (1997). Role of histone tails in nucleosome remodeling by *Drosophila* NURF. *EMBO J.*, **16**: 4717-4726.
- Gerace, L.; Comeau, C. and Benson, M. (1984). Organization and modification of nuclear lamina structure. *J. Cell Sci. Suppl.*, **1**: 137-160.
- Gerace, L. and Burke, B. (1988). Functional organization of the nuclear envelope. *Ann. Rev. Cell Biol.*, **4**: 335-374.
- Girard, J.; Payan, P. and Sardet, C. (1982). Changes in intracellular cations following fertilization of sea urchin eggs. Na^+/H^+ and Na^+/K^+ exchanges. *Exp. Cell Res.*, **142**: 215-221.
- Glass, J.R. and Gerace, L. (1990). Lamins A and C bind and assemble at the surface of mitotic chromosomes. *J. Cell Biol.*, **111**: 1047-1057.

- Glikin, G.C.; Ruberti, I. and Worcel, A. (1984). Chromatin assembly in *Xenopus* oocytes: *in vitro* studies. *Cell*, **37**: 33-41.
- Goodwin, G.H.; Woodhead, L. and Johns, E.W. (1977). The presence of high mobility group non-histone proteins in isolated nucleosomes. *FEBS (Fed. Eur. Biochem. Soc.) Lett.*, **73**: 85
- Goodwin, G.H.; Walker, J.M. and Johns, E.W. (1978). The high mobility group (HMG) non-histone chromosomal proteins. In: The Cell Nucleus, Vol. VI, Chromatin, part C. Ed. H. Busch. Academic Press, New York, USA. pp 181-219.
- Green, G.R. and Poccia, D.L. (1985). Phosphorylation of sea urchin sperm H1 and H2B histones precedes chromatin decondensation and H1 exchange during pronuclear formation. *Devel. Biol.*, **108**: 235-245.
- Green, G.R. and Poccia, D.L. (1988). Interaction of sperm histone variants and linker DNA during spermiogenesis in the sea urchin. *Biochemistry*, **27**: 619-625.
- Green, G.R. and Poccia, D.L. (1989). Phosphorylation of sea urchin histone CSH2A. *Dev. Biol.*, **134**: 413-419.
- Green, G.R.; Collas, P.; Burrell, A. and Poccia, D.L. (1995). Histone phosphorylation during sea urchin development. *Seminars in Cell Biology*, **6**: 219-227.
- Groudine, M. and Conkin, K.F. (1985). Chromatin structure and *de-novo* methylation of sperm DNA: Implications for activation of the paternal genome. *Science*, **228**: 1061-1068.
- Gruss,C.; Wu, W.; Koller, T. and Sogo, J.M. (1993). Disruption of nucleosomes at the replication fork. *EMBO J.*, **12**: 4533-4545.
- Guarente, L. (1995). Transcriptional coactivators in yeast and beyond. *Trends Biochem. Sci.*, **20**: 516-521.
- Guerrier, P.; Brassart, M.; David, C. and Moreau, M. (1986). Sequential control of meiosis reinitiation by pH and Ca²⁺ in oocytes of the *Prosobranquio* mollusk, *Patella vulgata*. *Devel. Biol.*, **114**: 315-324.
- Gurdon, J.B. (1976). Injected nuclei in frog eggs: Fate, enlargement and chromatin dispersal. *J. Embryol. Exper. Morphol.*, **36**: 523-540.
- Gurdon, J.B. (1968). Changes in somatic cell nuclei inserted into growing and maturing amphibian oocytes. *J. Embryol. Exp. Morphol.*, **20**: 401-414.
- Gurdon, J.B. and Woodland, H.R. (1968). The cytoplasmic control of nuclear activity in animal development. *Biol. Rev.*, **43**: 233-267.
- Hansen, S.K. and Tjian,R. (1995). TAFs and TFIIA mediate differential utilization of the tandem *Adh* promoters. *Cell*, **82**: 1-20.
- Hamel,J.F.; Himmelman, J.H.; and Dufresne, L. (1993). Gametogenesis and Spawning of the Sea Cucumber *Psolus fabricii* (Duben and Korn). *Biol. Bull.*, **184**: 125-143.
- Heald, R. and McKeon, F. (1990). Mutations of phosphorylation sites in lamin A that prevent nuclear lamina disassembly in mitosis. *Cell*, **61**: 579-589.
- Hecht, A.; Laroche, T.; Strahl-Bolsinger,S.; Gasser, S.M. and Grunstein, M. (1995). Histone H3 and H4 N-termini interact with SIR3 and SIR4 proteins: A molecular model for the formation of heterochromatin in yeast. *Cell*, **80**: 583-592.

- Hecht, N.B. (1989). Mammalian protamines and their expression. In: Histones and other basic proteins. L. Hnilica, G. Stein and J. Stein (Eds), CRC Press, Boca Ratón, Florida, USA. pp 347-374.
- Herlands, L.; Allfrey, V.G. and Poccia, D. (1982). Translational regulation of histone synthesis in the sea urchin *Strongylocentrotus purpuratus*. *J. Cell Biol.*, **94**: 219-223.
- Herlands, L. and Maul, G.G. (1994). Characterization of a major nucleoplasmin-like germinal vesicle protein which is rapidly phosphorylated before germinal vesicle breakdown in *Spisula solidissima*. *Devel. Biol.*, **161**: 530-537.
- Hill, C.S.; Packman, L.C. and Thomas, J.O. (1990). Phosphorylation of clustered -Ser-Pro-X-Lys-Arg- motifs in sperm histones H1 and H2B. *EMBO J.*, **9**: 805-813.
- Hoffman, P. and Chalkley, R. (1978). Procedures for minimizing protease activity during isolation of nuclei, chromatin and the histones. In: Methods in Cell Biology, Vol. XVII, Chromatin and chromosomal protein research. Ed. G. Stein; J. Stein; and L.J. Kleinsmith. Academic Press, New York, USA. pp. 1-12
- Hoffman, A.; Chiang, C.M.; Oelgeschlager, T.; Xie, X.; Burley, S.K.; Nakatani, Y. and Roeder, R. (1996). A histone octamer-like structure within TFIID. *Nature*, **380**: 356-359.
- Höger, T.H.; Krohne, G. and Kleinschmidt, J.A. (1991). Interaction of *Xenopus* lamins A and LII with chromatin *in vitro* mediated by a sequence element in the carboxyterminal domain. *Exper. Cell Res.*, **197**: 280-289.
- Hozumi, K.; Iwata, K.; Fukue, K.; Itoh, T.; Sakairi, N.; Nishimura, S-I.; Tokura, S.; Katagiri, Ch.; and Nishi, N. (1996). Structure-Function Relationship of Molecular Chaperone Protein-Nucleoplasmin-. N. Nishi (Ed.). *Peptide Chemistry 1995*. Protein Research Foundation, Osaka-Japan. pp. 341-344.
- Ikadai, H. and Hoshi, M. (1981). Biochemical studies on the acrosome reaction of the starfish, *Asterias amurensis*: Purification and characterization of the acrosome reaction-inducing substance. *Devel. Growth Differ.*, **23**: 81-88.
- Ito, T.; Tyler, J.K.; Bulger, M.; Kobayashi, R. and Kadonaga, J.T. (1996a). ATP-facilitated chromatin assembly with a nucleoplasmin-like protein from *Drosophila melanogaster*. *J. Biological Chem.*, **271**: 25041-25048.
- Ito, T.; Bulger, M.; Kobayashi, R. and Kadonaga, J.T. (1996b). *Drosophila* NAP-1 is a core histone chaperone that functions in ATP-facilitated assembly of regularly spaced nucleosomal arrays. *Mol. Cell. Biol.*, **16**: 3112-3124.
- Itoh, T.; Ohsumi, K. and Katagiri, Ch. (1993). Remodeling of human sperm chromatin mediated by nucleoplasmin from amphibian eggs. *Develop. Growth Differ.*, **35**: 59-66.
- Itoh, T.; Ausio, J.; and Katagiri, CH. (1997). Histone H1 Variants as Sperm-Specific Nuclear Proteins of *Rana catesbeiana*, and Their Role in Maintaining a Unique Condensed State of Sperm Chromatin. *Molecular Reproduction and Development*, **47**: 181-190.
- Isackson, P.J.; Debold, W.A. and Reeck, G.R. (1980). Isolation and separation of chicken erythrocyte high mobility group non-histone chromatin proteins by chromatography on phosphocellulose. *FEBS Lett.*, **119**: 337.
- Iwata, K.; Hozumi, K.; Tokura, S.; and Nishi, N. (1997). Interaction of Molecular Chaperone Protein, Nucleoplasmin, with DNA-Protamine Complex as Fish Sperm Nuclei Model. C. Kitada (Ed.). *Peptide Chemistry 1996*. Protein Research Foundation, Osaka-Japan. pp. 161-164.
- Iwata, K.; Hozumi, K.; Itoh, T.; Sakairi, N.; Tokura, S.; Katagiri, CH.; and Nishi, N. (1997). Conformation of nucleoplasmin and its interaction with DNA-protamine complex as a simple model of fish sperm nuclei. *Int. J. Biol. Macromol.*, **20**: 171-178.

- Iwata, K.; Hozumi, K.; Iihara, A.; Nomuzi, M.; Sakairi, N. and Nishi, N. (1999). Mechanism of salmon sperm decondensation by nucleoplasmin. *Int. J. Biol. Macromol.*, **26**: 95-101.
- Jackson, V.; Shires, A.; Tanphaichitr, N. and Chalkley, R. (1976). Modification to histones immediately after synthesis. *J. Mol. Biol.*, **104**: 471-483.
- Jordan, A. (1982). Tesis Doctoral. Facultad de Química, Universidad de Barcelona, Barcelona-España.
- Johnson, R.T. and Rao, P.N. (1971). Nucleo-cytoplasmic interactions in the achievement of nuclear synchrony in DNA synthesis and mitosis in multinucleate cells. *Biol. Rev.*, **46**: 97-155.
- Johnson, J.D.; Epel, D. and Paul, M. (1976). Intracellular cations following fertilization of sea urchin eggs after fertilization. *Nature*, **262**: 661-664.
- Kamakaka, R.T.; Bulger, M. and Kadonaga, J.T. (1993). Potentiation of RNA polymerase II transcription by Gal4-VP16 during but not after DNA replication and chromatin assembly. *Genes Dev.*, **7**: 1779-1795.
- Kasinsky, H.E. (1989). Specificity and distribution of sperm basic proteins. In: Histones and other nuclear basic proteins. L.S. Hnilica, G. Stein and J. Stein (Eds.), CRC Press, Boca Ratón, Florida, USA, pp. 73-163.
- Kasinsky, H.E. (1995). Evolution and origins of sperm nuclear basic proteins. In: Advances in spermatozoal phylogeny and taxonomy. B.G.M. Jamieson, J. Ausió, and J.L. Justine (Eds.), Mém. natn. Hist. nat., Paris, France, **166**: 463-473.
- Kasinsky, H.E.; Huang, S.Y.; Kwaik, S.; Mann, M.; Sweeney, J. and Yee, B. (1978). On the diversity of sperm histones in the vertebrates: Electrophoretic variability of testis-specific histone patterns in Anura contrasts with relative constancy in Squamata. *J. Exp. Zool.*, **203**: 109-126.
- Kasinsky, H.E.; Huang, S.Y.; Mann, M.; Roca, J.; and Subirana, J.A. (1985). On the diversity of sperm histones in the vertebrates: IV. Cytochemical and amino acid analysis in Anura. *J. Exp. Zool.*, **234**: 33-46.
- Katagiri, C. and Moriya, M. (1976). Spermatozoon response to the toad egg matured after removal of germinal vesicle. *Devel. Biol.*, **50**: 235-241.
- Katagiri, C. and Ohsumi, K. (1994). Remodeling of sperm chromatin induced in egg extracts of amphibians. *Int. J. Dev. Biol.*, **38**: 209-216.
- Kaufman, P.D. (1996). Nucleosome assembly: the CAF and the HAT. *Curr. Opin. Cell Biol.*, **8**: 369-373.
- Kaufman, P.D. and Botchan, M.R. (1994). Assembly of nucleosomes: do multiple assembly factors mean multiple mechanisms?. *Curr. Opin. Genet. Dev.*, **4**: 229-235.
- Kaufman, P.D.; Kobayashi, R.; Kessler, N. and Stillman, B. (1995). The p150 and p60 subunits of chromatin assembly factors 1: a molecular link between newly synthesized histones and DNA replication. *Cell*, **81**: 1105-1114.
- Kawasaki, K.; Philpott, A.; Avilion, A.A.; Berrios, M. and Fisher, P.A. (1994). Chromatin decondensation in *Drosophila* embryo extracts. *J. Biol. Chem.*, **269**: 10169-10176.
- Kellogg, D.R.; Kikuchi, A.; Fujii-Nakata, T.; Turck, C.W. and Murray, A.W. (1995). Members of the NAP / SET family of proteins interact specifically with B-type cyclins. *J. Cell Biol.*, **130**: 661-673.
- Kim, J.L.; Nikolov, D.B. and Burley, S.K. (1993). Co-crystal structure of TBP recognizing the minor groove of a TATA element. *Nature*, **365**: 520-527.

- Kingston, R.E.; Bunker, C.A. and Imbalzano, A.N. (1996). Repression and activation by mulyiprotein complexes that alter chromatin structure. *Genes Dev.*, **10**: 905-920.
- Kishimoto, T.; and Kanatami, H. (1973). Induction of starfish oocyte maturation by disulfide reducing agents. *Exp. Cell. Res.*, **82**: 296-302.
- Kishimoto, T.; Cayer, M.L.; and Kanatami, H. (1976). Starfish oocyte maturation and reduction of disulfide-bond on oocyte surface. *Exp. Cell. Res.*, **101**: 104-110.
- Kleinschmidt, J.A. and Franke, W.W. (1982). Soluble acidic complexes containing histones H3 and H4 in nuclei of *Xenopus laevis* oocytes. *Cell*, **29**: 799-809.
- Kleinschmidt, J.A.; Dingwall, C.; Maier, G. and Franke, W.W. (1986). Molecular characterization of a karyophilic, histone-binding protein: cDNA cloning, amino acid sequence and expression of nuclear protein N1/N2 of *Xenopus laevis*. *EMBO J.*, **5**: 3547-3552.
- Kleinschmidt, J.A. and Seiter, A. (1988). Identification of domains involved in nuclear uptake and histone binding of protein N1 of *Xenopus laevis*. *EMBO J.*, **7**: 1605-1614.
- Kline, D. (1988). Calcium dependent events at fertilization of the frog egg: Injection of a calcium buffer blocks ion channel opening, exocytosis and formation of pronuclei. *Devel. Biol.*, **149**: 80-89.
- Koehler, J.K.; Wurschmidt, U. and Larsen, M.P. (1983). Nuclear and chromatin structure in rat spermatozoa. *Gamete Res.*, **8**: 357-370.
- Kokubo, T.; Gong,D.-W.; Wooton, J.C.; Horikoshi, M. and Roeder, R.G. (1993). Molecular cloning, structural relationships and interactions of *Drosophila* TFIID subunits. *Nature*, **367**: 484-487.
- Kornberg, R.D. (1974). Chromatin structure: A repeating unit of histones and DNA. *Science*, **184**: 868-871.
- Krohne, G. and Franke, W.W. (1980a). Immunological identification and localization of the predominant nuclear protein of the amphibian oocyte nucleus. *Proc. Natl. Acad. Sci. USA*, **77**: 1034-1038.
- Krohne, G. and Franke, W.W. (1980b). A mayor soluble acidic protein located in nuclei of diverse vertebrate species. *Exp. Cell Res.*, **129**: 167-189.
- Krude, T. (1995). Nucleosome assembly during DNA replication. *Current Biology*, **5**: 1232-1234.
- Krude, T. and Knippers, R. (1991). Transfer of nucleosomes from parental to replicated chromatin. *Mol. Cell. Biol.*, **11**: 6257-6267.
- Krude, T. and Knippers, R. (1993). Nucleosome assembly during complementary DNA strand synthesis in extracts from mammalian cells. *J. Biol. Chem.*, **268**: 14432-14442.
- Kunkle, M.; Longo, F.J. and Magun, B.E. (1978a). Nuclear protein changes in the maternally and paternally derived chromatin at fertilization. *J. Exper. Zool.*, **203**: 381-390.
- Kunkle, M.; Magun, B.E. and Longo, F.J. (1978b). Analysis of isolated sea urchin nuclei incubated in egg cytosol. *J. Exper. Zool.*, **203**: 381-390.
- Laemmli, U.K. (1970). Cleavage of structural proteins during the assembly of the head of the bacteriophage T4. *Nature*, **227**: 680-685.

- Lanford, R.E.; Kanda, P. and Kennedy, R.C. (1986). Induction of nuclear transport with a synthetic peptide homologous to the SV40 T antigen transport signal. *Cell*, **46**: 575-582.
- Lanzendorf, S.E.; Slusser, J.; Malloney, M.K. et al. (1988). A preclinical evaluation of pronuclear formation by microinjection of human spermatozoa into human oocytes. *Fertil. Steril.*, **49**: 835-842.
- Laskey, R.A. (1983). Phosphorylation of nuclear proteins. *Philos. Trans. R. Soc. Lond. B Biol. Sci.*, **302**: 143-150.
- Laskey, R.A.; Mills, A.D. and Morris, N.R. (1977). Assembly of SV40 chromatin in a cell-free system from *Xenopus* eggs. *Cell*, **10**: 237-243.
- Laskey, R.A.; Honda, B.M.; Mills, A.D. and Finch, J.T. (1978). Nucleosomes are assembled by an acidic protein which binds histones and transfers them to DNA. *Nature*, **275**: 416-420.
- Leno, G.H. and Laskey, R.A. (1991). The nuclear membrane determines the timing of DNA replication in *Xenopus* egg extracts. *J. Cell Biol.*, **112**: 557-566.
- Leno, G.H.; Downes, C.S. and Laskey, R.A. (1992). The nuclear membrane prevents replication of human G2 nuclei but not G1 nuclei in *Xenopus* egg extract. *Cell*, **69**: 151-158.
- Leno, G.H.; Mills, A.D.; Philpott, A. and Laskey, R.A. (1996). Hyperphosphorylation of nucleoplasmin facilitates *Xenopus* sperm decondensation at fertilization. *J. Biol. Chem.*, **271**: 7253-7256.
- Lohka, M.J. (1988). The reconstitution of nuclear envelopes in cell-free extracts. *Cell Biol. Internat. Rep.*, **12**: 833-848.
- Lohka, M.J. and Maller, J.L. (1985). Induction of nuclear envelope breakdown, chromosome condensation and spindle formation in cell-free extracts. *J. Cell Biol.*, **101**: 518-526.
- Lohka, M.J. and Masui, Y. (1983a). Formation *in vitro* of sperm pronuclei and mitotic chromosomes induced by amphibian ooplasmic components. *Science*, **220**: 719-721.
- Lohka, M.J. and Masui, Y. (1983b). The germinal vesicle material required for sperm pronuclear formation is located in the soluble fraction of egg cytoplasm. *Exper. Cell Res.*, **148**: 481-491.
- Lohka, M.J. and Masui, Y. (1984). Roles of cytoplasmic particles in nuclear envelope assembly and sperm pronuclear formation in cell-free preparations from amphibian eggs. *J. Cell Biol.*, **98**: 1222-1230.
- Longo, F.J. (1973). Fertilization: A comparative ultrastructural review. *Biol. Reprod.*, **9**: 149-215.
- Longo, F.J. (1976). Derivation of the membrane comprising the male pronuclear envelope in inseminated sea urchin eggs. *Devel. Biol.*, **49**: 347-368.
- Longo, F.J. (1978). Insemination of immature sea urchin (*Arbacia punctulata*) eggs. *Devel. Biol.*, **62**: 271-291.
- Longo, F.J. (1984). Transformation of sperm nuclei incorporated into sea urchin (*Arbacia punctulata*) embryos at different stages of the cell cycle. *Devel. Biol.*, **103**: 168-181.
- Longo, F.J. (1990). Dynamics of sperm nuclear transformation at fertilization. In: Fertilization in Mammals. Eds: B.D. Bavister; J. Cummins and E.R.S. Roldan. Serono Symposia, Norwell, MA, USA. Pp. 297-307.
- Longo, F.J. (1997). Fertilization. Second Edition. Chapman & Hall. London, UK.
- Longo, F.J.; Krohne, G. and Franke, W.W. (1987). Basic proteins of the perinuclear theca of mammalian spermatozoa and spermatids: A novel class of cytoskeletal elements. *J. Cell Biol.*, **105**: 1105-1120.

- Longo, F.J.; Mathews, L. and Palazzo, R.E. (1994). Sperm nuclear transformations in cytoplasmic extracts from surf clam, *Spisula solidissima* oocytes. *Devel. Biol.*, **162**: 245-258.
- Longo, F.J.; Ushiyama, A.; Chiba, K. and Hoshi, M. (1995). Ultrastructural localization of acrosomal reaction-inducing substance (ARIS) on sperm of the starfish, *Asterias amurensis*. *Mol. Reprod. Devel.*, **41**: 91-99.
- Luttmer, S.J. and Longo, F.J. (1987). Rates of male pronuclear enlargement in sea urchin zygotes. *J. Exper. Zool.*, **243**: 289-298.
- Maleszewski, M. (1992). Behavior of sperm nuclei incorporated into parthenogenetic mouse eggs prior to the first cleavage division. *Mol. Reprod. Devel.*, **33**: 215-221.
- Mantle, T.J. and Noone, P. (1996). Chromatofocusing. In: Protein Purification Protocols. Ed. S. Doonan. Humana Press Inc., Totowa, NJ, USA. Methods in Molecular Biology, Vol. 59, Chapter 24: 249-254.
- Maruyama, Y.K. (1980). Artificial induction of oocyte Maturation and Development in the sea cucumbers *Holothuria leucospilota* and *Holothuria pardalis*. *Biol. Bull.*, **158**: 339-348.
- Maruyama, Y.K. (1985). Holothurian Oocyte Maturation Induced by Radial Nerve. *Biol. Bull.*, **168**: 249-262.
- Maruyama, Y.K. (1986). Induction of Sea Cucumber Oocyte Maturation by Starfish Radial Nerve Extracts. *The Journal of Experimental Zoology*, **238**: 241-248.
- Maruyama, Y.K. (1990). Roles of the Polar Cytoplasmic Region in Meiotic Divisions in Oocytes of the Sea Cucumber, *Holothuria leucospilota*. *Biol. Bull.*, **179**: 264-271.
- Masui, Y. (1985). Meiotic arrest in animal oocytes. In: Biology of Fertilization. Eds: C.B. Metz and A. Monroy. Vol 1, Academic Press, Orlando, FL, USA. Pp. 189-219.
- Masui, Y.; Meyerhof, P.G.; Miller, M.A. and Wasserman, W.J. (1977). Roles of bivalent cations in maturation and activation of vertebrate oocytes. *Differentiation*, **9**: 49-57.
- Masui, Y. and Clarke, H. (1979). Oocyte maturation. *Int. Rev. Cytol.*, **57**: 185-282.
- Matsui, T.; Nishiyama, I.; Hino, A. and Hoshi, M. (1986). Induction of the acrosoma reaction in starfish. *Develop. Growth and Differ.*, **28**: 339-348.
- Maul, G.G. and Schatten, G. (1986). Nuclear lamins during gametogenesis, fertilization and early development. In: Nucleocytoplasmic Transport. Eds: R. Peters and M. Trendelenburg. Springer-Verlag, New York, USA. Pp. 123-134.
- Maul, G.G.; French, B.T. and Bechtol, K.B. (1986). Identification and redistribution of lamins during nuclear differentiation in mouse spermatogenesis. *Devel. Biol.*, **115**: 68-77.
- McPherson, S. and Longo, F.J. (1993). Chromatin structure-function alterations during mammalian spermatogenesis: DNA nicking and repair in elongating spermatids. *Eur. J. Histochem.*, **37**: 109-128.
- Meijer, L.; and Guerrier, P. (1984). Maturation and Fertilization in Starfish Oocytes. *International Review of Cytology*, **86**: 129-196.
- Mengod, L. (1976). Tesis Doctoral. Universidad de Barcelona, Barcelona, España.
- Merriam, R.W. (1969). Movement of cytoplasmic proteins into nuclei induced to enlarge and initiate DNA or RNA synthesis. *J. Cell Sci.*, **5**: 333-359.

- Moore, S. and Stein, W.H. (1963). Chromatographic determination of amino acids by the use of automatic recording equipment. *Methods Enzymol.*, **6**: 819-831.
- Moss, S.B.; Donovan, M.J. and Bellv , A.R. (1987). The occurrence and distribution of lamin proteins during mammalian spermatogenesis and early embryonic development. *Ann. NY. Acad. Sci.*, **513**: 74-89.
- Motlik, J.; Kopecny, V.; Pivko, J. and Fulka, J. (1980). Distribution of proteins labeled during meiotic maturation in rabbit and pig eggs at fertilization. *J. Reprod. Fert.*, **58**: 415-419.
- Mu oz-Guerra, S.; Azor n, F.; Casas, M.T.; Marcket, X.; Maristany, M.A.; Roca, J. and Subirana, J.A. (1982). Structural organization of sperm chromatin from the fish *Carassius auratus*. *Exp. Cell Res.*, **137**: 47-53.
- Murray, A. and Hunt, T. (1993). The cell cycle. W.H. Freeman, New York.
- Naish, S.J.; Perreault, S.D.; Foehner, A.L. and Zirkin, B.R. (1987a). DNA synthesis in the fertilizing hamster sperm nucleus: Sperm template availability and egg cytoplasmic control. *Biol. Reprod.*, **36**: 245-253.
- Naish, S.J.; Perreault, S.D. and Zirkin, B.R. (1987b). DNA synthesis following microinjection of heterologous sperm and somatic cell nuclei into hamster oocytes. *Gamete Res.*, **18**: 109-120.
- Neigeborn, L. and Carlson, M. (1994). Genes affecting the regulation of *SUC2* gene expression by glucose repression in *Saccharomyces cerevisiae*. *Genetics*, **108**: 845-858.
- Ner, S.S. and Travers, A.A. (1994). HMG-D, the *Drosophila melanogaster* homologue of HMG1 protein, is associated with early embryonic chromatin in the place of histone H1. *EMBO J.*, **13**: 1817-1822.
- Newport, J. (1987). Nuclear reconstitution *in vitro*: Stages of assembly around protein-free DNA. *Cell*, **48**: 205-217.
- Newrock, K.M.; Alfageme, C.R.; Nardi, R.V. and Cohen, L.H. (1977). Histone changes during chromatin remodeling in embryogenesis. *Cold Spring Harbor Symp. Quant. Biol.*, **42**: 421-431.
- Nightingale, K.; Dimitrov, S.; Reeves, R. and Wolffe, A. (1996). Evidence for a shared structural role for HMG1 and linker histones B4 and H1 organizing chromatin. *EMBO J.*, **15**: 548-561.
- Nikolov, D.B. and Burley, S.K. (1994). 2.1A resolution refined structure of a TATA box-binding protein (TBP). *Nature Structural Biology*, **1**: 621-637.
- Nikolov, D.B.; Chen, H.; Halay, E.D.; Usheva, A.A.; Hisatake, K.; Lee, D.K.; Roeder, R.G. and Burley, S.K. (1995). Crystal structure of a TFIIIB-TBP-TATAelement ternary complex. *Nature*, **377**: 121-128.
- Noll, M. and Kornberg, R.D. (1977). Action of micrococcal nuclease on chromatin and the location of histone H1. *J. Mol. Biol.*, **109**: 393-404.
- Nonchev, S. and Tsanev, R. (1990). Protamine-histone replacement and DNA replication in the male mouse pronucleus. *Mol. Reprod. Devel.*, **25**: 72-76.
- Ohshima, H. (1925). Notes on the development of the sea cucumber, *Thyone briareus*. *Science*, **61**: 402-422.
- Ohsumi, K. and Katagiri, Ch. (1991a). Characterization of the ooplasmic factor inducing decondensation of and protamine removal from toad sperm nuclei: Involvement of nucleoplasmin. *Dev. Biol.*, **148**: 295-305.
- Ohsumi, K. and Katagiri, Ch. (1991b). Ocurrence of H1 subtypes specific to pronuclei and cleavage-stage cell nuclei of anuran amphibians. *Dev. Biol.*, **147**: 110-120.

- Ohsumi, K.; Shimada, A.; Okumura, E.; Kishimoto, T. and Katagiri, C. (1995). Dependence of removal of sperm-specific proteins from *Xenopus* sperm nuclei on the phosphorylation state of nucleoplasmin. *Develop. Growth Differ.*, **37**: 329-336.
- Olivares, C; Azorin, F; Subirana, J.A. and Cornudella, L. (1987). The Interaction of the Histone H1-Related Protein ϕ_0 with Chromatin. *Biophysical Chemistry*, **28**: 51-57.
- Orphanides, O.; LeRoy, G.; Chang, C.-H.; Luse, D.S. and Reinberg, D.(1998). FACT, a factor that facilitates transcript elongation through nucleosomes. *Cell*, **92**: 105-116.
- Owen-Hughes, T.; Utley, R.T.; Cole, J.; Peterson, C.L. and Workman, J.L.(1996). Persistent site -specific remodeling of a nucleosome arrays by transientaction of the SWI / SNF complex. *Science*, **273**: 513-516.
- Palermo, G.; Joris, H.; Devroey, P. and VanSteirteghem, A.C. (1992). Pregnancies after intracytoplasmic injection of single spermatozoon into an oocyte. *Lancet*, **340**: 17-18.
- Palermo, G.; Joris, H.; Derde, M-P. et al. (1993). Sperm characteristics and outcome of human assisted fertilization by subzonal insemination and intracytoplasmic sperm injection. *Fertil. Steril.*, **59**: 826-835.
- Panyim, S. and Chalkley, R. (1969). High resolution acrylamide gel electrophoresis of histones. *Arch. Biochem. Biophys.*, **130**: 337-346.
- Parthun, M.R.; Widom, J. and Gottschling, D.E. (1996). The major cytoplasmic histone acetyltransferase in yeast links to chromatin replication and histone metabolism. *Cell*, **87**: 85-94.
- Payan, P.; Girard, J-P; and Ciapa, B. (1983). Mechanisms regulating intracellular pH in sea urchin egg. *Devel. Biol.*, **100**:29-38.
- Perreault, S.D. (1990). Regulation of sperm nuclear reaction during fertilization. In: Fertilization in Mammals. Eds: B.D. Bavister, J. Cummins and E.R.S. Roldan. Serono Symposia, Norwell, MA, USA. pp. 285-296.
- Perreault, S.D. (1992). Chromatin remodeling in mammalian zygotes. *Mutation Res.*, **296**: 43-55.
- Perreault, S.D.; Naish, S.J. and Zirkin,B.R. (1987). The timing of hamster sperm nuclear decondensation and male pronuclear formation is related to sperm nuclear disulfide bondcontent. *Biol. Reprod.*, **36**: 239-244.
- Perreault, S.D.; Barbee, R.R. and Slott, V.L. (1988). Importance of glutathione in the acquisition and maintenance of sperm nuclear decondensing activity in maturing hamster oocytes. *Devel. Biol.*, **125**: 181-186.
- Peterson, C.L. and Tamkun, J.W. (1995). The SWI-SNF complex: a chromatin remodeling machine?.*Trends Biochem. Sci.(TIBS)*, **20**: 143-146.
- Phelan, J.; Subirana, J.A.; and Cole, R.D. (1972). An Unusual Group of Lysine-Rich Histones from Gonads of a Sea Cucumber, *Holothuria tubulosa*. *Eur. J. Biochem.*, **31**: 63-68.
- Philpott, A.; Leno, G.H. and Laskey, R.A. (1991). Sperm decondensation in *Xenopus* egg cytoplasm is mediated by nucleoplasmin. *Cell*, **65**: 569-578.
- Philpott, A. and Leno, G.H. (1992). Nucleoplasmin remodels sperm chromatin in *Xenopus* egg extracts. *Cell*, **69**: 759-767.
- Poccia, D. (1986). Remodeling of nucleoproteins during gametogenesis, fertilization and early development. *Int. Rev. Cytol.*, **105**: 1-65.

- Poccia, D. (1989). Reactivation and remodeling of the sperm nucleus following fertilization. In: The Molecular Biology of Fertilization. H. Schatten and G. Schatten (Eds). Academic Press, New York, USA. Pp. 115-135.
- Poccia, D. (1995). Male germ line specific histones of sea urchin and sea stars. In: Advances in spermatozoal phylogeny and taxonomy. B.G.M. Jamieson, J. Ausió, and J.L. Justine (Eds). Mém. Mus. natn. Hist. nat., París, France, vol. 166, pp. 475-489.
- Poccia, D.; Krystal, G.; Nishioka, D. and Salik, J. (1978). Control of sperm chromatin structure by egg cytoplasm in the sea urchin. In: ICN-UCLA Symposium on Molecular and Cellular Biology: Cell Reproduction, XII. E.R. Dirksen, D.M. Prescott and C.F. Fox (Eds.). Academic Press, New York, USA. Pp. 197-206
- Poccia, D.; Salik, J. and Krystal, G. (1981). Transitions in histone variants of the male pronucleus following fertilization and evidence for a maternal store of cleavage-stage histones in the sea urchin egg. *Devel. Biol.*, **82**: 287-296.
- Poccia, D.; Wolff, R.; Kragh, S. and Williamson, P. (1985). RNA synthesis in male pronuclei of the sea urchin. *Biochim. Biophys. Acta*, **824**: 349-356.
- Poccia, D.L.; Simpson, M.V.; and Green, G.R. (1987). Histone variant transitions during sea urchin spermatogenesis. *Dev. Biol.*, **121**: 445-453.
- Poccia, D.; Pavon, W. And Green, G.R. (1990). 6DMAP inhibits chromatin decondensation but not sperm histone kinase in sea urchin male pronuclei. *Exper. Cell Res.*, **188**: 226-234.
- Poccia, D.L. and Green, G.R. (1992). Packaging and unpackaging the sea urchin sperm genome. *Trends Biochem. Sci. (TIBS)*, **17**: 223-227.
- Porter, D.C. and Vacquier, V.D. (1986). Phosphorylation of sperm histone H1 is induced by the egg jelly layer in the sea urchin *Strongylocentrotus purpuratus*. *Devel. Biol.*, **116**: 203-212.
- Prather, R.S. and Schatten, G. (1992). Construction of the nuclear matrix at the transition from maternal to zygotic control of development in the mouse: An immunocytochemical study. *Mol. Reprod. Devel.*, **32**: 203-208.
- Prats, E. and Cornudella, L. (1995). The gene encoding the sperm-specific basic nuclear protein ϕ_0 from sea cucumber. In: Jamieson, B.G.M.; Ausio, J. and Justine, J.-L. (Eds). Advances in Spermatozoal Phylogeny and Taxonomy. Mém. Mus. natn. Hist. nat. (Paris), **166**: 491-500.
- Pruss, D.; Hayes, J.J. and Wolffe, A.P. (1995). Nucleosomal anatomy-where are the histone?. *BioEssays*, **17**: 161-170.
- Randall, S.K. and Kelly, T.J. (1992). The fate of parental nucleosomes during SV40 DNA replication. *J. Biol. Chem.*, **267**: 14259-14265.
- Raven, C.P. (1961). Oogenesis: The storage of developmental information. Pergamon Press Ltd., London, UK.
- Rice, P.; Garduño, R.; Itoh, T.; Katagiri, C. and Ausio, J. (1995). Nucleoplasmin-mediated decondensation of *Mytilus* sperm chromatin. Identification and partial characterization of a nucleoplasmin-like protein with sperm-nuclei decondensing activity in *Mytilus californianus*. *Biochemistry*, **34**: 7563-7568.
- Risley, M.S.; Einheber, S. and Bruncrot, D.A. (1986). Changes in DNA topology during spermatogenesis. *Chromosoma*, **94**: 217-227.
- Risley, M.S. (1990). Chromatin organization in sperm. In, Chromosoma: Eukaryotic, Prokaryotic and Viral. K.W. Adolph (Ed.), CRC Press, Boca Raton, FL. pp.61-85.

- Robbins, J.; Dilworth, S.M.; Laskey, R.A. and Dingwall, C. (1991). Two interdependent basic domains in nucleoplasmin nuclear targeting sequence: identification of a class of bipartite nuclear targeting sequence. *Cell*, **64**: 615-623.
- Rodman, T.C.; Pruslin, F.H.; Hoffmann, H.P. and Allfrey, V.G. (1981). Turnover of basic chromosomal proteins in fertilized eggs: A cytoimmunochemical study of events *in vivo*. *J. Cell Biol.*, **90**: 351-361.
- Roth, S.Y. and Allis, C.D. (1996a). Histone acetylation and chromatin assembly: A single escort, multiple dances?. *Cell*, **87**: 5-8.
- Roth, S.Y. and Allis, C.D. (1996b). The subunit-exchange model of histone acetylation. *Trends in Cell Biology*, **6**: 371-375.
- Ruiz-Carrillo, A.; Wangh, L.J. and Allfrey, V.G. (1975). Processing of newly synthesized histone molecules. *Science*, **190**: 117-128.
- Ruiz-Lara, S.A.; Cornudella, L.; and Rodriguez-Campos, A. (1996). Dissociation of protamine-DNA complexes by *Xenopus* nucleoplasmin and minichromosome assembly *in vitro*. *Eur. J. Biochem.*, **240**: 186-194.
- Salik, J.; Herlands, L.; Hoffmann, H.P. and Poccia, D. (1981). Electrophoretic analysis of the stored histone pool in unfertilized sea urchin eggs: Quantification and identification by antibody binding. *J. of Cell Biology*, **90**: 385-395.
- Saperas, N.; Ribes, E.; Buesa, C.; García-Hegart, F. and Chiva, M. (1993). Differences in chromatin condensation during spermiogenesis in two species of fish with distinct protamines. *J. Exp. Zool.*, **265**: 185-194.
- Saperas, N.; Ausió, J.; Lloris, D. and Chiva, M. (1994). On the evolution of protamines in bony fish: Alternatives to the "retroviral horizontal transmission" hypothesis. *J. Mol. Evol.*, **39**: 282-295.
- Saperas, N.; Buesa, C.; Abian, J.; Vandekerckhove, J.; Kasinsky, H.E. and Chiva, M. (1996). The primary structure of a chondrichyan protamine: a new apparent contradiction in protamine evolution. *J. Mol. Evol.*, **43**: 528-535.
- Saperas, N.; Chiva, M.; Pfeiffer, D.C.; Kasinsky, H.E. and Ausio, J. (1997). Sperm nuclear basic proteins (SNBPs) of Agnathans and Chondrichthyans: Variability and evolution of sperm proteins in fish. *J. Mol. Evol.*, **44**: 422-431.
- Sauer, F.; Hansen S.K. and Tjian, R. (1995a). Multiple TAF_{II}s directing synergistic activation of transcription. *Science*, **270**: 1783-1788.
- Sauer, F.; Hansen, S.K. and Tjian, R. (1995b). DNA template and activator-coactivator requirements for transcriptional synergism by *Drosophila* bicoid. *Science*, **270**: 1825-1828.
- Savic, A.; Richman, P.; Williamson, P. and Poccia, D. (1981). Alterations in Chromatin structure during early sea urchin embryogenesis. *Proc. Natl. Acad. Sci. USA*, **78**: 3706-3710.
- Schatten, G.; Maul, G.G.; Schatten, H. et al. (1985a). Nuclear lamins and peripheral nuclear antigens during fertilization and embryogenesis in mice and sea urchin. *Proc. Natl. Acad. Sci. USA*, **82**: 4727-4731.
- Schmidt-Zachmann, M.S.; Hügle-Dörr, B. and Franke, W.W. (1987). A constitutive nucleolar protein identified as a member of the nucleoplasmin family. *EMBO J.*, **6**: 1881-1890.
- Sealy, L.; Cotten, M. and Chalkley, R. (1986). *Xenopus* nucleoplasmin: egg vs oocyte. *Biochemistry*, **25**: 3064-3072.
- Sewell, M.A.; Tyler, P.A.; Young, C.M.; and Conard, C. (1997). Ovarian Development in the Class Holothuroidea: a Reassessment of the "Tubule Recruitment Model". *Biol. Bull.*, **192**: 17-26.

- Sheehan, M.A.; Mills, A.D.; Sleeman, A.M. et al. (1988). Steps in the assembly of replication-competent nuclei in a cell-free system from *Xenopus* eggs. *J. Cell Biol.*, **106**: 1-12.
- Shen, S.S. and Steinhardt, R.A. (1978). Direct measurement of intracellular pH during metabolic derepression of the sea urchin egg. *Nature*, **272**: 253-254.
- Skobolina, M. (1976). Role of karyoplasm in the emergence of capacity of egg cytoplasm to induce DNA synthesis in transplanted sperm nuclei. *J. Embryol. Exper. Morphol.*, **36**: 67-72.
- Smiley, S.; and Cloney, R.A. (1985). Ovulation and the fine structure of the *Stichopus californicus* (Echinodermata: Holothuroidea) fecund ovarian tubules. *Biol. Bull.*, **169**: 342-364.
- Smith, A.E.; Kalderon, D.; Roberts, B.L.; Colledge, W.H.; Edge, M.; Gillett, P.; Markham, A.; Pancha, E. and Richardson, W.D. (1985). *Proc. R. Soc. London, Ser. B*, **221**: 43-53.
- Smith, S. and Stillman, B. (1989). Purification and characterization of CAF-1, a human cell factor required for chromatin assembly during DNA replication in vitro. *Cell*, **58**: 15-25.
- Smith, S. and Stillman, B. (1991a). Stepwise assembly of chromatin during DNA replication in vitro. *EMBO J.*, **10**: 971-980.
- Smith, S. and Stillman, B. (1991b). Immunological characterization of chromatin assembly factor 1, a human cell factor required for chromatin assembly during DNA replication in vitro. *J. Biol. Chem.*, **266**: 12041-12047.
- Sobel, R.E.; Cook, R.G.; Perry, C.A.; Annunziato, A.T. and Allis, C.D. (1995). Conservation of the position-related acetylation sites in newly synthesized histones H3 and H4. *Proc. Natl. Acad. Sci. USA*, **92**: 1237-1241.
- Stern, M.J.; Jensen, R.E. and Herskowitz, I. (1984). Five SWI genes are required for expression of the HO genes in yeast. *J. Mol. Biol.*, **178**: 853-868.
- Stick, R. and Hausen, P. (1985). Changes in the nuclear lamina composition during early development of *Xenopus laevis*. *Cell*, **41**: 191-200.
- Stricker, S.; Prather, R.; Simerly, C. et al. (1989). Nuclear architectural changes during fertilization and development. In: The Cell Biology of Fertilization. H. Schatten and G. Schatten (Eds.). Academic Press, New York, USA. pp. 225-250.
- Strickland, M.; Strickland, W.; Brandt, W.; and Von Holt, C. (1977). The complete amino-acid sequence of histone H2B(1) from sperm of the sea urchin *Parechinus angulosus*. *Eur. J. Biochem.*, **77**: 263-275.
- Strickland, M.; Strickland, W.; Brandt, W.; Von Holt, C.; Wittmann-Liebold, B.; and Lehmann, A. (1978). The complete aminoacid sequence of histone H2B(3) from sperm of the sea urchin *Parechinus angulosus*. *Eur. J. Biochem.*, **89**: 443-452.
- Strickland, W.; Strickland, M.; Brandt, W.; and Von Holt, C. (1977). The complete amino-acid sequence of histone H2B(2) from sperm of the sea urchin *Parechinus angulosus*. *Eur. J. Biochem.*, **77**: 277-286.
- Strickland, W.N.; Strickland, M.; Brandt, W.F.; Von Holt, C.; Lehmann, A. and Wittmann-Liebold, B. (1980). The primary structure of histone H1 from the sea urchin *Parechinus angulosus*. Sequence of the c-terminal CNBr peptide and the entire primary structure. *Eur. J. Biochem.*, **104**: 567-578.
- Struhl, K. (1996). Chromatin structure and RNA polymerase II connection: implications for transcription. *Cell*, **84**: 179-182.

- Subirana, J.A. (1970). Nuclear Proteins from a Somatic and a Germinal Tissue of the Echinoderm *Holothuria tubulosa*. *Exp. Cell Res.*, **63**: 253-260.
- Subirana, J.A. (1990a). Analysis of the charge distribution in the C-terminal region of histone H1 as related to its interaction with DNA. *Biopolymers*, **29**: 1351-1357.
- Subirana, J.A. (1990b). Proteins as counterions of DNA: A new model of nucleoprotamine structure. In: Water and ions in biomolecular systems. Ed. Vasilescu, D.; Jaz, J.; Packer, L. & Pullman, B. Proceedings of the 5th UNESCO international conference. Birkhäuser Verlag Basel. pp. 63-70.
- Subirana, J.A. (1991). Protein-DNA interactions in spermatozoa. In: Comparative spermatology 20 years after. Ed. Baccetti, B. Serono symposia publications from Raven Press. New York. Vol. **75**: 89-92.
- Subirana, J.A. (1992). Order and disorder in 30 nm chromatin fibers. *FEBS*, **302**: 105-107.
- Subirana, J.A.; Cozculluela, C.; Palau, J. and Unzeta, M. (1973). Protamines and other basic proteins from spermatozoa of molluscs. *Biochem. Biophys. Acta*, **317**: 364-379.
- Subirana, J.A.; Muñoz-Guerra, S.; Aymami, J.; Radermacher, M. and Frank, J. (1985). The layered organization of nucleosomes in 30 nm chromatin fibers. *Chromosoma*, **91**: 377-390.
- Subirana, J.A. and Colom, J. (1987). Comparison of protamines from freshwater and marine bivalve mollusc: evolutionary implications. *FEBS Lett.*, **220**: 193-196.
- Sugasawa, K.; Ishimi, Y.; Eki, T.; Hurwitz, J.; Kikuchi, A. and Hanaoaka, F. (1992). Nonconservative segregation of parental nucleosomes during simian virus 40 chromosome replication in vitro. *Proc. Natl. Acad. Sci. USA*, **89**: 1055-1059.
- Suzuki, M.; Sugiura, M. and Ebashi, S. (1990). Sea urchin Protease Specific to the SPKK Motif in histone. *J. Biochem.*, **108**: 347-355.
- Szöllösi, D. and Ris, H. (1961). Observations on sperm penetration in the rat. *J. Biophys. Biochem. Cytol.*, **10**: 275-283.
- Szöllösi, D.; Szöllösi, M.S.; Czolowska, R. and Tarkowski, A.K. (1990). Sperm penetration into immature mouse oocytes and nuclear changes during maturation: an EM study. *Biol. Cell.*, **69**: 53-64.
- Takamune, K.; Nishida, K.; Takai, H. and Katagiri, Ch. (1991). Primary structure of toad sperm protamines and nucleotide sequence of their cDNAs. *Eur. J. Biochem.*, **196**: 401-406.
- Tanaka, Y. (1958). Seasonal changes occurring in the gonad of *Stichopus japonicus*. *Bull. Fac. Fish. Hokkaido Univ.*, **9**: 29-36.
- Taunton, J.; Hassig, C.H. and Schreiber, S.L. (1996). A mammalian histone deacetylase related to the yeast transcriptional regulator Rpd3p. *Science*, **272**: 408-411.
- Thadani, V.M. (1979). Injection of sperm heads into immature rat oocyte. *J. Exper. Zool.*, **210**: 161-168.
- Thomas, J.O. and Kornberg, R.D. (1978). The study of histone association. In: Methods in Cell Biology. G. Stein, J. Stein, L.K. Stein (Eds.). Academic Press, New York, Vol. 18: 429-440.
- Tsukiyama, T.; Becker, P.B. and Wu, C. (1994). ATP-dependent nucleosome disruption at a heat-shock promoter mediated by binding of GAGA transcription factor. *Nature*, **367**: 525-532.

- Ulitzer, N. and Gruenbaum, Y. (1989). Nuclear envelope assembly around sperm chromatin in cell-free preparation from *Drosophila* embryos. *FEBS Lett.*, **259**: 113-116.
- Vacquier, V.D.; Brandriff, B. and Glabe, C.G. (1979). The effect of soluble egg jelly on the fertilizability of acid-dejellied sea urchin egg. *Develop. Growth and Differ.*, **21**: 49-60.
- Vacquier, V.D. (1986). Activation of sea urchin spermatozoa during fertilization. *Trends Biochem. Sci. (TIBS)*, **11**: 78-81.
- Vacquier, V.D.; Porter, D.C.; Keller, S.H. and Aukermann, M. (1989). Egg jelly induces the phosphorylation of histone H3 in spermatozoa of the sea urchin *Arbacia punctulata*. *Devel. Biol.*, **133**: 111-118.
- Vancurova, I.; Paine, T.M.; Lou, W. and Paine, P.L. (1995). Nucleoplasmin associates with and is phosphorylated by casein kinase II. *J. Cell Sci.*, **108**: 779-787.
- Varga-Weisz, P.D.; Wilm, M.; Bonte, E.; Dumas, K.; Mann, M. and Becker, P.B. (1997). Chromatin-remodelling factor CHRAC contains the ATPases ISWI and topoisomerase II. *Nature*, **388**: 598-602.
- Varga-Weisz, P.D.; Blank, T.A. and Becker, P.B. (1995). Energy-dependent chromatin accessibility and nucleosome mobility in a cell-free system. *EMBO J.*, **14**: 2209-2216.
- Verdaguer, N.; Perelló, M.; Palau, J. and Subirana, J.A. (1993). Helical structure of basic proteins from spermatozoa comparison with model peptides. *Eur. J. Biochem.*, **214**: 879-887.
- Verreault, A.; Kaufman, P.D.; Kobayashi, R. and Stillman, B. (1996). Nucleosome assembly by a complex of CAF-1 and acetylated histones H3 / H4. *Cell*, **87**: 95-104.
- Verrijzer, C.P.; Chen, J.-L.; Yokomori, K. and Tjian, R. (1995). Binding of TAFs to core elements directs promoter selectivity by RNA polymerase II. *Cell*, **81**: 1115-1125.
- Vigers, G.P.A. and Lohka, M.J. (1991). A distinct vesicle population targets membrane and pore complexes to the nuclear envelope in *Xenopus* eggs. *J. Cell Biol.*, **112**: 545-556.
- von Lindern, M.; van Baal, S.; Wiegant, J.; Raap, A.; Hagemeijer, A. and Grosveld, G. (1992). *can*, a putative oncogene associated with myeloid leukemogenesis, may be activated by fusion of its 3' half to different genes: Characterization of the setgene. *Mol. Cell. Biol.*, **12**: 3346-3355.
- Wakefield, L. and Gurdon, J.B. (1983). Cytoplasmic regulation of 5S RNA genes in nuclear-transplant embryos. *EMBO J.*, **2**: 1613-1619.
- Walker, J.M.; Gooderham, K.; Hastings, J.R.; Mayes, E. and Johns, E.W. (1980). The primary structures of non-histone chromosomal proteins HMG 1 and 2. *FEBS Lett.*, **122**: 264-270.
- Wallrath, L.L.; Lu, Q.; Granok, H. and Elgin, S.C.R. (1994). Architectural variations of inducible eukaryotic promoters: present and remodeling chromatin structure. *BioEssays*, **16**: 165-170.
- Ward, W.S. (1993). Deoxyribonucleic acid loop domain tertiary structure in mammalian spermatozoa. *Biol. Reprod.*, **48**: 1193-1201.
- Ward, W.S.; Partin, A.W. and Coffey, D.S. (1989). DNA loop domains in mammalian spermatozoa. *Chromosoma*, **98**: 153-159.
- Ward, W.S.; McNeil, J.; deLarca, J. and Lawrence, J. (1996). Localization of three genes in the hook-shaped sperm nucleus by fluorescent *in situ* hybridization. *Biol. Reprod.*, **54**: 1271-1278.

- Wasserman, W.J. and Smith, L.D. (1978). The cyclic behavior of a cytoplasmic factor controlling nuclear membrane breakdown. *J. Cell Biol.*, **78**: R15-R22.
- Wiesel, S. and Schultz, G.A. (1981). Factors which may affect removal of protamine from sperm DNA during fertilization in the rabbit. *Gamete Res.*, **4**: 25-34.
- Wilding, M. (1996). Calcium and cell cycle control in early embryos. *Zygote*, **4**: 1-6.
- Winston, F. and Carlson, M. (1992). Yeast SWI / SNF transcriptional activators and the SPT / SIN chromatin connection. *Trends Genet.*, **8**: 387-391.
- Wolffe, A.P. and Pruss, D. (1996). Deviant nucleosomes: the functional specialization of chromatin. *Trends in Genetics (TIG)*, **12**: 58-62.
- Workman, J.L. and Kingston, R.E. (1992). Nucleosome core displacement *in vitro* via a metastable transcription factor/nucleosome complex. *Science*, **258**: 1780-1784.
- Wright, S.J. and Schatten, G. (1990). Teniposide, a topoisomerase II inhibitor, prevents chromosome condensation and separation but not decondensation in fertilized surf clam, *Spisula solidissima* oocytes. *Devel. Biol.*, **142**: 224-232.
- Wu, C. (1980). The 5' ends of *Drosophila* heat shock genes in chromatin are hypersensitive to DNase I. *Nature*, **286**: 854-860.
- Xie, X.; Kokubo, T.; Cohen, S.L.; Mirza, U.A.; Hoffman, A.; Chait, B.T.; Roeder, R.G.; Nakatani, Y. and Burley, S.K. (1996). Structural similarity between TAFS and the heterotetrameric core of the histone octamer. *Nature*, **380**: 316-322.
- Yamashita, M.; Onozato, H.; Nakanishi, T. and Nagahama, Y. (1990). Breakdown of the sperm nuclear envelope is a prerequisite for male pronucleus formation: Direct evidence from the gynogenetic crucian carp, *Carassius auratus langsdorffii*. *Devel. Biol.*, **137**: 155-160.
- Yanagimachi, R. and Noda, Y.D. (1970a) Ultrastructural changes in the hamster sperm head during fertilization. *J. Ultrastruct. Res.*, **31**: 465-485.
- Yanagimachi, R. (1994). Mammalian Fertilization. In: The Physiology of Reproduction. 2nd. Ed., E. Knobil & J.D. Neill (Eds.), Raven Press, New York, pp. 189-317.
- Young, R.J. (1979). Rabbit sperm chromatin is decondensed by thiol-induced proteolytic activity not endogenous to its nucleus. *Biol. Reprod.*, **20**: 1001-1004.
- Young, R.J. and Sweeney, K. (1978). Mammalian ova and one-cell embryos do not incorporate phosphate into nucleic acids. *Eur. J. Biochem.*, **91**: 111-117.
- Zawel, L. and Reinberg, D. (1992). Advances in RNA polymerase II transcription. *Curr. Opin. Cell Biol.*, **4**: 488-495.
- Zawel, L. and Reinberg, D. (1993). Initiation of transcription by RNA polymerase II. A multi step process. *Prog. Nucl. Acids Res. Mol. Biol.*, **44**: 67-108.
- Zirkin, B.R. and Chang, T.S.K. (1977). Involvement of endogenous proteolytic activity in thiol-induced release of DNA template restrictions in rabbit sperm nuclei. *Biol. Reprod.*, **17**: 131-137.
- Zirkin, B.R.; Chang, T.S.K. and Heaps, J. (1980). Involvement of an acrosin-like proteinase in the sulphydryl-induced degradation of rabbit sperm nuclear protamine. *J. Cell Biol.*, **85**: 116-121.

VIII.- Bibliografía

- Zirkin, B.R.; Soucek, D.A.; Chang, T.S.K. and Perreault, S.D. (1985). *In vitro* and *in vivo* studies of mammalian sperm nuclear decondensation. *Gamete Res.*, **11**: 349-365.
- Zirkin, B.R.; Perreault, S.D. and Naish, S.J. (1989). Formation and function of the male pronucleus during mammalian fertilization. In: *The Molecular Biology of Fertilization*. Eds. H. Schatten and G. Schatten. Academic Press, New York, pp. 91-114.
- Zweidler, A. (1978). Resolution of histones by polyacrylamide gel electrophoresis in presence of nonionic detergents. In: *Methods in Cell Biology*, Vol. XVII. Ed. G. Stein, J. Stein, and L.J. Kleinsmith. Academic Press, New York, USA. pp 235-251.