

Evolución geoquímica de cuencas evaporíticas terciarias: implicaciones en la composición isotópica disuelto en el océano durante el terciario

Dionisio Ignacio Cendón Sevilla



Aquesta tesi doctoral està subjecta a la llicència **Reconeixement- NoComercial – SenseObraDerivada 3.0. Espanya de Creative Commons.**

Esta tesis doctoral está sujeta a la licencia **Reconocimiento - NoComercial – SinObraDerivada 3.0. España de Creative Commons.**

This doctoral thesis is licensed under the **Creative Commons Attribution-NonCommercial-NoDerivs 3.0. Spain License.**



Evolución geoquímica de cuencas evaporíticas terciarias: implicaciones en la composición isotópica disuelto en el océano durante el terciario

Dionisio Ignacio Cendón Sevilla

FACULTAD DE GEOLOGIA
Departamento de Cristalografía, Mineralogía
y Depósitos Minerales

INSTITUTO DE CIENCIAS DE LA TIERRA
"JAUME ALMERA" - CSIC

DIRECTORES DE LA TESIS:

Dr. Juan José Pueyo Mur
Dr. Carlos Ayora Ibáñez.

Barcelona, 1999

9.- BIBLIOGRAFÍA

- Anati, D. A. y Stiller, M. (1991). The post-1979 thermohaline structure of the Dead Sea and the role of double-diffusive mixing. *Limnology and Oceanography* 36(2), 342-354.
- Anati, D. A., Stiller, M., Shasha, S. y Gat, J. R. (1987). Changes in the thermo-haline structure of the Dead Sea: 1979-1984. *Earth and Planetary Science Letters* 84, 109-121.
- Anderson, A. J., Clark, A. H., Ma, X.-P., Palmer, G. R., MacArthur, J. D. y Roedder, E. (1989). Proton-induced x-ray and gamma-ray emission analysis of unopened fluid inclusions. *Economic Geology* 84, 924-939.
- Arthurton, R. S. (1973). Experimentally produced halite compared with Triassic layered halite-rock from Cheshire, England. *Sedimentology* 20, 145-160.
- Artyushkov, E. V., Baer, M. A. y Mörner, N.-A. (1996). The east Carpathians: Indications of the phase transitions, lithospheric failure and decoupled evolution of thrust belt and its foreland. *Tectonophysics* 262, 101-132.
- Ault, W. U. y Kulp, J. L. (1959). Isotopic geochemistry of sulphur. *Geochimica et Cosmochimica Acta* 16, 201-235.
- Ayora, C., Cendón, D. I., García-Veigas, J., Pueyo, J. J. y Taberner, C. (1997). Claves sobre la génesis de yacimientos de silvita. *Boletín de la Sociedad Española de Mineralogía* 20-A, 79-80.
- Ayora, C., Cendón, D. I., Taberner, C., Fanlo, I., García-Veigas, J. y Pueyo, J. J. (1998). Water-rock reactions in evaporite basins: their role in the formation of potash deposits. En: *Water-Rock Interaction* (ed. A. Hulston), pp. 289-292. Balkema.
- Ayora, C. y Fontarnau, R. (1990). X-ray microanalysis of frozen fluid inclusions. *Chemical Geology* 89, 135-148.
- Ayora, C., García-Veigas, J. y Pueyo, J. J. (1994a). The chemical and hydrological evolution of an ancient potash-forming evaporite basin as constrained by mineral sequence, fluid inclusion composition, and numerical simulation. *Geochimica et Cosmochimica Acta* 58(16), 3379-3394.
- Ayora, C., García-Veigas, J. y Pueyo, J. J. (1994b). X-ray microanalysis of fluid inclusions and its application to the geochemical modeling of evaporite basins. *Geochimica et Cosmochimica Acta* 58, 43-55.
- Ayora, C., Taberner, C., Pierre, C. y Pueyo, J. J. (1994c). Refining the $\delta^{34}\text{S}$ and $\delta^{18}\text{O}$ values of sulphate in ancient oceans. *Mineralogical Magazine* 58A, 32-33.
- Ayora, C., Taberner, C., Pierre, C. y Pueyo, J. J. (1995). Modeling the sulfur and oxygen isotopic composition of sulfates through a halite-potash sequence: Implications for the hydrological evolution of the Upper Eocene Southpyrenean Basin. *Geochimica et Cosmochimica Acta* 59(9), 1799-1808.

- Baar, A. y Kühn, R. (1962). Der werdegang der kalisalzlagerstätten am oberrhein. *Neues Jahrbuch für Mineralogie, Abhandlungen* 97(3), 289-336.
- Banks, D. A., Davies, G. R., Yardley, B. W. D., McCaig, A. M. y Grant, N. T. (1991). The chemistry of brines from an Alpine thrust system in the Central Pyrenees: an application of fluid inclusion analysis to the study of fluid behaviour in orogenesis. *Geochimica et Cosmochimica Acta* 55, 1021-1030.
- Bein, A., Hovorka, S. D. y Fisher, R. S. (1991). Fluid inclusions in bedded permian halite, Palo Duro basin, Texas: Evidence for modification of seawater in evaporite brine-pools and subsequent early diagenesis. *Journal of Sedimentary Petrology* 61(1), 1-14.
- Benali, S., Schreiber, B. C., Helman, M. L. y Philp, R. P. (1995). Characterization of organic matter from a restricted/evaporative sedimentary environment: Late Miocene of Lorca basin, Southeastern Spain. *AAPG Bulletin* 79(6), 816-830.
- Bentor, Y. K. (1961). Some geochemical aspects of the Dead Sea and the question of its age. *Geochimica et Cosmochimica Acta* 25, 239-260.
- Bentor, Y. K. (1969). On the evolution of subsurface brines in Israel. *Chemical Geology* 4, 83-110.
- Bergerat, F. y Geysant, J. (1980). La fracturation tertiaire de l'Europe du Nord: résultat de la collision Afrique-Europe. *Comptes Rendus de l'Académie des Sciences de Paris (D)*, 290, 1521-1524.
- Bitzer, K., Travé, A., Carmona, J. M. y Calvet, F. (1998). Fluid flow in foreland basins during emplacement of thrust sheets: modelling the south-Pyrenean Ainsa basin. *Bulletin de la Société Géologique de France* 169(5), 627-634.
- Blanc-Valleron, M. M. (1991). Les formations paléogènes évaporitiques du bassin potassique de Mulhouse et des bassins plus septentrionaux d'Alsace. Tesis doctoral, Université Louis Pasteur.
- Blanc-Valleron, M. M. y Gannat, E. (1985). Cartographie de subsurface du Salifère supérieur du bassin potassique de Mulhouse (Oligocène, Alsace). *Bulletin de la Société Géologique de France* 8(1), 823-836.
- Boillot, G. y Capdevilla, R. (1977). The Pyrenees: Subduction and collision? *Earth and Planetary Science Letters* 35, 151-160.
- Boiron, M. C., Dubessy, J., Andre, N., Briand, A., Lacour, J. L., Mauchien, P. y Mermet, J. M. (1991). Analysis of mono-atomic ions in individual fluid inclusions by laser-produced plasma emission spectroscopy. *Geochimica et Cosmochimica Acta* 55, 917-923.
- Boiron, M. C., Dubessy, J., Moissette, A., Geertsen, C., Banks, D. A., Prieto, A. C., Lacour, J. L. y Mauchien, P. (1995). Elemental analysis of individual aqueous inclusions. Part I: New developments using micro laser ablation optical emission spectroscopy (MLA-OES). *Boletín de la Sociedad Española de Mineralogía* 18-1, 28-29.

- Bottrell, S. H., Yardley, B. y Buckley, F. (1988). A modified crush-leach method for analysis of fluid inclusion electrolytes. *Bulletin de Minéralogie* 111, 279-290.
- Braitsch, O. (1971). *Salt Deposits. Their Origin and Composition*. Springer-Verlag.
- Brantley, S. L., Crerar, D. A., Moller, N. E. y Weare, J. H. (1984). Geochemistry of a modern marine evaporite: Bocana de Virrilá, Perú. *Journal of Sedimentary Petrology* 54(2), 447-462.
- Bryant, R. G., Sellwood, B. W., Millington, A. C. y Drake, N. A. (1994). Marine-like potash evaporite formation on a continental playa: case study from Chott el Djerid, southern Tunisia. *Sedimentary Geology* 90, 269-291.
- Burke, W. H., Denison, R. E., Hetherington, E. A., Koepnick, R. B., Nelson, H. F. y Otto, J. B. (1982). Variation of seawater $^{87}\text{Sr}/^{86}\text{Sr}$ throughout Phanerozoic time. *Geology* 10, 516-519.
- Busquets, P., Ortí, F., Pueyo, J. J., Riba, O., Rosell, L., Saez, A., Salas, R. y Taberner, C. (1985). Evaporite Deposition and Diagenesis in the Saline (Potash) Catalan Basin, Upper Eocene. Institut d'Estudis Ilerdencs (Diputació Provincial de Lleida).
- Butler, R. W. H., Lickorish, W. H., Grasso, M., Pedley, H. M. y Ramberti, L. (1995). Tectonics and sequence stratigraphy in Messinian basins, Sicily: Constraints on the initiation and termination of the Mediterranean salinity crisis. *Geological Society of America Bulletin* 107(4), 425-439.
- Canals, A. (1992). Microanalyses des inclusions primaires dans la halite: contraintes pour la modélisation de la sédimentation évaporitique. Exemple du bassin de Mulhouse. En: IFP. Division de Géologie-Géochimie.
- Canals, A., Carpentier, B., Huc, A. Y., Guilhaumou, N. y Ramsey, M. H. (1993). Microanalysis of primary fluid inclusions in halite: constraints for an evaporitic sedimentation modeling. Application to the Mulhouse basin (France). *Organic Geochemistry* 20(8), 1139-1151.
- Cann, J. H. y De Deckker, P. (1981). Fossil Quaternary and living foraminifera from athalassic (non-marine) saline lakes, Southern Australia. *Journal of Paleontology* 55(3), 660-670.
- Casas, E. y Lowenstein, T. K. (1989). Diagenesis of saline pan halite: comparison of petrographic features of modern, quaternary and permian halites. *Journal of Sedimentary Petrology* 59(5), 724-739.
- Casas, E., Lowenstein, T. K., Spencer, R. J. y Pengxi, Z. (1992). Carnallite mineralization in the nonmarine, Qaidam basin, China: Evidence for the early diagenetic origin of potash evaporites. *Journal of Sedimentary Petrology* 62(5), 881-898.
- Cendón, D. I., Ayora, C., García-Veigas, J., Pierre, C., Pueyo, J. J. y Taberner, C. (1997a). The evolution of the Lorca basin and the composition of the Messinian ocean. *Terra Nova* 9, 412.

- Cendón, D. I., Ayora, C. y Pueyo, J. J. (1997b). The origin of barren bodies in the Subiza potash deposit, Navarra, Spain. *XIV-ECROFI, Abstracts Volume*, 62-63.
- Cendón, D. I., Ayora, C. y Pueyo, J. J. (1998a). The origin of barren bodies in the Subiza potash deposit, Navarra, Spain: Implications for sylvite formation. *Journal of Sedimentary Research* 68(1), 43-52.
- Cendón, D. I., Pueyo, J. J., Ayora, C., Taberner, C. y Pierre, C. (1998b). Geochemical evolution of Suria evaporite sequence: Implications for the chemistry of the Eocene ocean. *Mineralogical Magazine* 62A, 294-295.
- Cendón, D. I., Taberner, C., Ayora, C., García-Veigas, J., Pueyo, J. J. y Pierre, C. (1997c). The formation of the Lorca halite sequence: Marine versus continental. *Neogene Mediterranean Paleoceanography. Abstract Volume*, 17-18.
- Claypool, G. E., Holser, W. T., Kaplan, I. R., Sakai, H. y Zak, I. (1980). The age curves of sulfur and oxygen isotopes in marine sulfate and their mutual interpretation. *Chemical Geology* 28, 199-260.
- Courtot, C., Gannat, E. y Wendling, E. (1972). Le bassin potassique de Mulhouse et ses environs. Étude du Tertiaire. *Sciences Géologiques Bulletin, Strasbourg* 25(2-3), 69-91.
- Craig, H. (1961). Standard for reporting concentrations of deuterium and oxygen-18 in natural waters. *Science* 133, 1833-1834.
- Chiba, H. y Sakai, H. (1985). Oxygen isotope exchange rate between dissolved sulfate and water at hydrothermal temperatures. *Geochimica et Cosmochimica Acta* 49, 993-1000.
- Das, N., Horita, J. y Holland, H. D. (1990). Chemistry of fluid inclusions in halite from the Salina Group of the Michigan Basin: Implications for Late Silurian seawater and the origin of sedimentary brines. *Geochimica et Cosmochimica Acta* 54, 319-327.
- Decima, A., McKenzie, J. A. y Schreiber, B. C. (1988). The origin of "evaporative" limestones: an example from the Messinian of Sicily (Italy). *Journal of Sedimentary Petrology* 58, 256-272.
- Decima, A. y Wezel, F. C. (1973). Osservazioni sulle evaporiti messiniane della Sicilia centro-meridionale. *Riv. Min. Siciliana* 22, 171-187.
- del Valle. (1929). Investigación de las sales potásicas de Navarra. *Boletín del Instituto Geológico de España* 51, 89-107.
- Dellwig, L. F. (1955). Origin of the salina salt of Michigan. *Journal of Sedimentary Petrology* 25(2), 83-110.
- DePaolo, D. J. e Ingram, B. L. (1985). High-resolution stratigraphy with Strontium isotopes. *Science* 227, 938-941.
- Diamond, L. W., Marshall, D. D., Jackman, J. A. y Skippen, G. B. (1990). Elemental analysis of individual fluid inclusions in minerals by Secondary Ion Mass Spectrometry

- (SIMS): Application to cation ratios of fluid inclusions in an Archaean mesothermal gold-quartz vein. *Geochimica et Cosmochimica Acta* 54, 545-552.
- Dubinina, V. N. y Yarzhemskii, Y. Y. (1953). Facies transitions in Verkhnekamsk deposit salt strata. *Doklage Akademia Nauk. SSSR.* 90(6), 1131-1134.
- Edmond, J. M., Measures, C., Mangum, B., Grant, B., Slater, F. R., Collier, R., Hudson, A., Gordon, L. I. y Corliss, J. B. (1979). On the formation of metal-rich deposits at ridge crests. *Earth Planetary Science Letters* 46, 1-18.
- El Tabakh, M., Schreiber, B. C., Utha-Aroon, C., Coshell, L. y Warren, J. K. (1998). Diagenetic origin of basal anhydrite in the Cretaceous Maha Sarakham salt: Khorat Plateau, NE Thailand. *Sedimentology* 45, 579-594.
- Elderfield, H. (1986). Strontium isotope stratigraphy. *Palaeogeography, Palaeoclimatology, Palaeoecology* 57, 71-90.
- Eugster, H. P., Harvie, C. E. y Weare, J. H. (1980). Mineral equilibria in a six-component seawater system, Na-K-Mg-Ca-SO₄-Cl-H₂O, at 25°C. *Geochimica et Cosmochimica Acta* 44, 1335-1347.
- Fanlo, I. y Ayora, C. (1998). The evolution of the Lorraine evaporite basin: implications for the chemical and isotope composition of the Triassic ocean. *Chemical Geology* 146, 135-154.
- Faura i Sans, M. y Marín, A. (1926). La cuenca potásica de Cataluña y Pirineo Central. *XIV Congreso Geológico Internacional, Madrid C-3*, 213.
- Field, C. y Lombardi, G. (1972). Sulfur isotopic evidence for the supergene origin of alunite deposits, Tolfa district, Italy. *Mineralium Deposita* 7(2), 113-125.
- Filly, A., Letolle, R. y Puset, M. (1975). L'Analyse isotopique du soufre. Aspects techniques. *Analysis* 3, 197-200.
- Fontes, J., Filly, A., Gaudant, J. y Düringer, P. (1991). Origine continentale des évaporites paléogènes de Haute Alsace: arguments paléocéologiques, sédimentologiques et isotopiques. *Bulletin de la Société Géologique de France* 162(4), 725-737.
- Frantz, J. D., Mao, H. K., Zhang, Y.-G., Wu, Y., Thompson, A. C., Underwood, J. H., Giauque, R. D., Jones, K. W. y Rivers, M. L. (1988). Analysis of fluid inclusions by X-ray fluorescence using synchrotron radiation. *Chemical Geology* 69, 235-244.
- Friedman, I. y O'Neil, J. R. (1977). Compilation of stable isotope fractionation factors of geochemical interest. En: *Data of Geochemistry*, Vol. Prof. Pap. (ed. M. Fleischer), pp. 440-444. U.S. Geological Survey.
- Friedrichsen, H. (1981). Geothermal systems in the upper Rhine Graben and northern Black Forest: a chemical and stable isotope study. *Tectonophysics* 73, 125-140.
- Fritz, B. (1981). Étude Thermodynamique et Modélisation des Réactions Hydrothermales et Diagénetiques. Tesis doctoral, Université Louis Pasteur de Strasbourg, Institut de Géologie.

- Fritz, P., Basharmal, G. M., Drimmie, R. J., Ibsen, J. y Qureshi, R. M. (1989). Oxygen isotope exchange between sulphate and water during bacterial reduction of sulphate. *Chemical Geology (Isotope Geoscience Section)* 79(2), 99-105.
- Galamay, A. R., Bukowski, K. y Przybylo, J. (1997). Chemical composition and origin of brines in the Badenian evaporite basin of the Carpathian Foredeep: fluid inclusion data from Wieliczka (Poland). *Slovak Geological Magazine* 3(2), 165-171.
- Galamay, A. R. y Karoli, S. (1997). Geochemistry of the Badenian salts from the East Slovakian Basin, Slovakia. *Slovak Geological Magazine* 3(3), 187-192.
- Ganor, J. y Katz, A. (1989). The geochemical evolution of halite structures in hypersaline lakes: The Dead Sea, Israel. *Limnology and Oceanography* 34(7), 1214-1223.
- García, D. (1986). Étude isotopique et géochimique des eaux thermales des Vosges méridionales. Application géothermique. Tesis doctoral, Université des Sciences et Techniques du Languedoc.
- García-Veigas, J. (1993). Geoquímica de inclusiones fluidas en formaciones salinas, Microanálisis Cryo-SEM-EDS. Tesis doctoral, Universitat de Barcelona.
- García-Veigas, J., Ortí, F., Rosell, L., Ayora, C., Rouchy, J.-M. y Lugli, S. (1995). The Messinian salt of the Mediterranean: geochemical study of the salt from the Central Sicily Basin and comparison with the Lorca Basin (Spain). *Bulletin de la Société Géologique de France* 166(6), 699-710.
- García-Veigas, J., Ortí, F., Rosell, L. y Inglès, M. (1994). Caracterización petrológica y geoquímica de la Unidad Salina messiniense de la cuenca de Lorca (sondeos S4 y S5). *Geogaceta* 15, 78-81.
- García-Veigas, J., Rosell, L. y Garlicki, A. (1997). Petrology and geochemistry (fluid inclusions) of Miocene halite rock salts (Badenian, Poland). *Slovak Geology Magazine* 3(3), 181-186.
- Garlicki, A. (1979). Sedymentacja soli miocenskich w Polsce. *Prace Geol. Komis. Nauk Geol. PAN Kraków* 119.
- Garlicki, A. (1994). Formal lithostratigraphic units of the Miocene: Wieliczka formation. *Przegląd Geologiczny* 42(1), 26-28.
- Garlicki, A. y Wiewiórka, J. (1981). The distribution of bromine in some halite rock salts of the Wieliczka salt deposit (Poland). *Annales Societatis Geologorum Poloniae* 51(3/4), 353-359.
- Garven, G. (1995). Continental-scale groundwater flow and geologic processes. *Annual Review of Earth and Planetary Science Letters* 23, 89-117.
- Geel, T. (1976). Messinian gypsiferous deposits of the Lorca Basin (Province of Murcia, SE Spain). *Memorie della Società Geologica Italiana* 16, 369-385.

- Gerler, J. (1987). Trace elements in hydrothermal quartzes and their relation to the amount and salinity of fluid inclusions. *Fortschritte der Mineralogie* 65 (Bhf. 1), 56.
- Gil Rincón, J. A. (1998). Modelización geodinámica y numérica de estructuras evaporíticas (Cuencas Surpirenaica y Cantábrica). Tesis doctoral, Universitat de Barcelona.
- Gizaw, B. (1996). The origin of high bicarbonate and fluoride concentrations in waters of the Main Ethiopian Rift Valley, East African Rift system. *Journal of African Earth Sciences* 22(4), 391-402.
- Grant, N. T., Banks, D. A., McCaig, A. M. y Yardley, B. W. D. (1990). Chemistry, source, and behavior of fluids involved in Alpine thrusting of the Central Pyrenees. *Journal of Geophysical Research* 95(B6), 9123-9131.
- Guillén-Mondéjar, F., Rodríguez Estrella, T., Arana, R. y López Aguayo, F. (1995a). Historia geológica de la cuenca de Lorca (Murcia): influencia de la tectónica en la sedimentación. *Geogaceta* 18, 30-33.
- Guillén-Mondéjar, F., Rodríguez Estrella, T., Arana, R. y López Aguayo, F. (1995b). Unidades tectosedimentarias y rupturas en la Cuenca de Lorca (Murcia). *Geogaceta* 17, 39-42.
- Günther, D., Audetat, A., Frischknecht, R. y Heinrich, A. (1997). A 193 nm excimer laser ablation-inductively coupled plasma-mass spectrometer system for fluid inclusion analysis. *XIV-ECROFI, Abstracts Volume*, 144-145.
- Gutiérrez, J. R. (1985). Étude des filons à Pb-Zn-Cu comportant une gangue sulfatée silicifiée de la région de Tahnn-Watwiller (Vosges du Sud). Tesis doctoral, Inst. nat. polytechn. Lorraine, Nancy.
- Hackett, J. P., Bischoff, J. y Bischoff, J. L. (1973). New data on the stratigraphy, extent, and geologic history of the Red Sea geothermal deposits. *Economic Geology* 68, 553-564.
- Halas, S., Jasionowski, M. y Peryt, T. M. (1996). Anomalia izotopowa w badenskich gipsach Poniżnia. *Przegląd Geologiczny* 44(10), 1054-1056.
- Halas, S. y Kurpiewski, A. (1982). Skład izotopowy tlenu i siarki w siarczanach rudy siarkowej ze złóż tarnobrzeskich. *Przegląd Geologiczny* 30, 69-73.
- Hall, T. A. (1971). *The Microprobe Assay of Chemical Elements*. Academic Press.
- Hardie, L. A. (1987). Perspectives dolomitization: A critical view of some current views. *Journal of Sedimentary Petrology* 57(1), 166-183.
- Hardie, L. A. (1990). The roles of rifting and hydrothermal CaCl₂ brines in the origin of potash evaporites: An hypothesis. *American Journal of Science* 290, 43-106.
- Hardie, L. A. (1996). Secular variation in seawater chemistry: An explanation for the coupled secular variation in the mineralogies of marine limestones and potash evaporates over the past 600 m.y. *Geology* 24(3), 279-283.

- Hardie, L. A. y Eugster, H. P. (1971). The depositional environment of marine evaporites: a case for shallow, clastic accumulation. *Sedimentology* 16, 187-220.
- Harvie, C. E., Moller, N. y Weare, J. H. (1984). The prediction of mineral solubilities in natural waters: The Na-K-Mg-Ca-H-Cl-SO₄-OH-HCO₃-CO₃CO-CO₂-H₂O system to high ionic strengths at 25^o C. *Geochimica et Cosmochimica Acta* 48, 723-751.
- Harvie, C. E., Weare, J. H., Hardie, L. A. y Eugster, H. P. (1980). Evaporation of Seawater: Calculated Mineral Sequences. *Science* 208, 498-500.
- Hauber, L. (1982). Geology of the salt field Rheinfelden-Riburg, Switzerland. En: *5th International Symposium on Salt*, Vol. 1 (ed. A. H. Coogan y L. Hauber), pp. 83-90. The Northern Ohio Geological Society INC.
- Haynes, F. M., Sterner, S. M. y Bodnar, R. J. (1988). Synthetic fluid inclusions in natural quartz. IV. Chemical analyses of fluid inclusions by SEM/EDA: Evaluation of method. *Geochimica et Cosmochimica Acta* 52, 969-977.
- Heinrich, C. A. y Cousens, D. R. (1989). Semi-quantitative electron microprobe analysis of fluid inclusion salts from the Mount Isa copper deposit (Queensland, Australia). *Geochimica et Cosmochimica Acta* 53, 21-28.
- Hendry, J. P., Taberner, C., Marshall, J. D., Pierre, C. y Carey, P. F. (en prensa). Coral reef diagenesis records pore fluid evolution and palaeohydrology of a siliciclastic basin margin succession (Eocene south Pyrenean foreland basin, NE Spain). *Geological Society of America Bulletin*.
- Herrmann, A. G. y Knipping, B. (1993). *Waste Disposal and Evaporites*. Springer-Verlag.
- Hite, R. J. (1983). The sulfate problem in marine evaporites. En: *6th International Symposium on Salt*, Vol. 1 (ed. B. C. Schreiber), pp. 217-230. Salt Institute.
- Hite, R. J. y Japakasetr, T. (1979). Potash deposits of the Khorat Plateau, Thailand and Laos. *Economic Geology* 74, 448-458.
- Hofmann, P., Huc, A. Y., Carpentier, B., Schaeffer, P., Albrecht, P., Keely, B., Maxwell, J. R., Sinninghe Damsté, J. S., De Leeuw, J. W. y Leythaeuser, D. (1993a). Organic matter of the Mulhouse Basin, France: a synthesis. *Organic Geochemistry* 20(8), 1105-1123.
- Hofmann, P., Leythaeuser, D. y Carpentier, B. (1993b). Paleoclimate controlled accumulation of organic matter in Oligocene evaporite sediments of Mulhouse basin. *Organic Geochemistry* 20(8), 1125-1138.
- Holser, W. T. (1963). Chemistry of brine inclusions in Permian salt from Hutchinson, Kansas. En: *Symposium on Salt* (ed. N. O. G. Society), pp. 86-95.
- Holser, W. T. (1966). Bromide geochemistry of salt rocks. En: *Marine Evaporites. Origin, Diagenesis and Geochemistry*, Vol. Benchmark papers in geology (ed. D. W. Kirkland), pp. 333-360. Dowden, Hutchinson & Ross, Inc.

- Holser, W. T. (1968). Chemistry of brine inclusions in Permian salt from Hutchison, Kansas (abstr.). *Geological Society of America. Special Paper* 88, 537.
- Holser, W. T. y Kaplan, I. R. (1966). Isotope geochemistry of sedimentary sulfates. *Chemical Geology* 1, 93-135.
- Holser, W. T., Kaplan, I. R., Sakai, H. y Zak, I. (1979). Isotope geochemistry of oxygen in the sedimentary sulfate cycle. *Chemical Geology* 25(1-2), 1-17.
- Holwerda, J. G. y Hutchinson, R. W. (1968). Potash-bearing evaporites in the Danakil area, Ethiopia. *Economic Geology* 63, 124-150.
- Holland, H. D. (1972). The geologic history of sea water- an attempt to solve the problem. *Geochimica et Cosmochimica Acta* 36, 637-651.
- Holland, H. D. (1973). Systematics of the isotopic composition of sulfur in the oceans during the Phanerozoic and its implications for atmospheric oxygen. *Geochimica et Cosmochimica Acta* 37, 2605-2616.
- Holland, H. D., Horita, J. y Seyfried, W. E. (1996). On the secular variations in the composition of phanerozoic marine potash evaporites. *Geology* 24(11), 993-996.
- Holland, H. D., Lazar, B. y McCaffrey, M. (1986). Evolution of the atmosphere and oceans. *Nature* 320, 27-33.
- Horita, J., Friedman, T. J., Lazar, B. y Holland, H. D. (1991). The composition of Permian seawater. *Geochimica et Cosmochimica Acta* 55, 417-432.
- Horita, J., Weinberg, A., Das, N. y Holland, H. D. (1996). Brine inclusions in Halite and the origin of the middle Devonian Prairie Evaporites of Western Canada. *Journal of Sedimentary Research* 66(5), 956-964.
- Horn, E. E. (1983). Analysis of fluid inclusions using the Heidelberg proton microprobe (abstr.). *Symposium, ECROFI, Orleans, 6-8 April 1983, Program* 34.
- Horn, E. E. y Traxel, K. (1987). Investigations of individual fluid inclusions with the Heidelberg proton microprobe- a nondestructive analytical method. *Chemical Geology* 61, 29-35.
- Houk, R. S., Fassel, V. A., Flesh, G. D., Svec, H. J., Gray, A. L. y Taylor, C. E. (1980). Inductively coupled argon plasma as an ion source for mass spectrometric. Determination of trace elements. *Anal. Chem.* 52, 2283-2289.
- Hovorka, S. D. (1992). Halite pseudomorphs after gypsum in bedded anhydrite - clue to gypsum-anhydrite relationships. *Journal of Sedimentary Petrology* 62(6), 1098-1111.
- Hsü, K. J., Ryan, W. B. F. y Cita, M. B. (1973). Late Miocene desiccation of the Mediterranean. *Nature* 242, 240-244.

- Hudec, P. P. y Sonnenfeld, P. (1980). Comparison of Caribbean solar ponds with inland solar lakes of British Columbia. En: *Hypersaline Brines and Evaporite Environments*, Vol. 28 (ed. A. Nissenbaum), pp. 101-114. Elsevier.
- IGME. (1977). Velez-Blanco. En: Mapa Geológico de España (1:50.000).
- IGME. (1982). *Aplicación de la Investigación de Pizarras Bituminosas en la Zona de Lorca (Murcia) (Fase II)*. IGME.
- Jarvis, K. E., Gray, A. L. y Houk, R. S. (1992). *Handbook of Inductively Coupled Plasma Mass Spectrometry*. Chapman and Hall.
- Jean-Baptiste, P., Charlou, J. L. y Stievenard, M. (1997). Oxygen isotope study of mid-ocean ridge hydrothermal fluids: Implication for the oxygen-18 budget of the oceans. *Geochimica et Cosmochimica Acta* 61(13), 2669-2677.
- Jensen, M. L. y Nakai, N. (1962). Sulfur isotope meteorite standards, results and recommendations. En: *Biogeography of Sulfur Isotopes* (ed. M. L. Jensen), pp. 30-35. Natl. Sci. Found Symp.
- Jones, B. F., Eugster, H. P. y Rettig, S. L. (1977). Hydrochemistry of the Lake Magadi basin, Kenya. *Geochimica et Cosmochimica Acta* 41, 53-72.
- Kaever, M. y Meiburg, P. (1976). Stratigraphy of the Tertiary and its depositional sedimentary environments of the Diemel-Weser area (Hessen Graben, W-Germany). *Bulletin de la Société Géologique de France* 18(5), 1277-1282.
- Kasprzyk, A. (1993). Gypsum facies in the Badenian (Middle Miocene) of southern Poland. *Canadian Journal of Earth Science* 30, 1799-1814.
- Kasprzyk, A. (1997). Oxygen and sulphur isotope composition of Badenian (Middle Miocene) gypsum deposits in southern Poland: a preliminary study. *Geological Quarterly* 41(1), 53-60.
- Kastens, K. y Mascle, J. (1990). The geological evolution of the Tyrrhenian Sea: An introduction to the scientific results of ODP Leg 107. *Proceedings of the Ocean Drilling Program, Scientific Results* 107, 3-26.
- Kelly, W. C. y Burgio, P. A. (1983). Cryogenic scanning electron microscopy of fluid inclusions in ore and gangue minerals. *Economic Geology* 78, 1262-1267.
- Kendall, A. C. (1989). Brine mixing in the Middle Devonian of western Canada and its possible significance to regional dolomitization. *Sedimentary Geology* 64, 271-285.
- Korenevskii, S. M. (1989). Substitution and depletion zones of potash and potassium-magnesium salt deposits. *All-Union Scientific-Research Institute of Geology, Leningrad, (Translated from Litologiya i Poleznye Iskopaemye)* 1, 66-82.
- Kováč, M., Nagymarosy, A., Soták, J. y Sutovská, K. (1993). Late Tertiary paleogeographic evolution of the Western Carpathians. *Tectonophysics* 226, 401-415.

- Kovalevich, V. M. (1990). Phanerozoic evolution of ocean water composition. *Geochemistry International* 25(6), 20-27.
- Kovalevich, V. M. y Petrichenko, O. I. (1997). Chemical composition of brines in Miocene evaporite basins of the Carpathian region. *Slovak Geological Magazine* 3(3), 173-180.
- Kozłowski, A. (1978). Pneumatolytic and hydrothermal activity in the Karkonosze-Izera block. *Acta Geologica Polonica* 28, 171-222.
- Kramer, J. R. (1965). History of sea water. Constant temperature-pressure equilibrium models compared to liquid inclusion analyses. *Geochimica et Cosmochimica Acta* 29, 921-945.
- Kramm, U. y Wedepohl, K. H. (1991). The isotopic composition of strontium and sulfur in seawater of Late Permian (Zechstein) age. *Chemical Geology* 90, 253-262.
- Kroopnick, P. y Craig, H. (1976). Oxygen isotope fractionation in dissolved oxygen in the deep sea. *Earth and Planetary Science Letters* 32, 375-388.
- Lazar, B. y Holland, H. D. (1988). The analysis of fluid inclusions in halite. *Geochimica et Cosmochimica Acta* 52, 485-490.
- Le Carlier, C., Royer, J.-J. y Flores, L. (1994). Convective heat transfer at the Soultz-sous-Forêts Geothermal site: implications for oil potential. *First Break* 12(11), 553-560.
- Linn, K. O. y Adams, S. S. (1966). Barren halite zones in potash deposits Carlsbad, New Mexico. En: *Second Symposium on Salt*, Vol. 1, pp. 59-69.
- Longinelli, A. (1980). Isotope geochemistry of some Messinian evaporites: Paleoenvironmental implications. *Paleogeography, Paleoclimatology, Paleoecology* 29, 95-123.
- Longinelli, A. (1989). Oxygen-18 and sulphur-34 in dissolved oceanic sulphate and phosphate. En: *Handbook of Environmental Isotope Geochemistry*, Vol. 3 (ed. P. F. y J. C. Fontes), pp. 219-255. Elsevier.
- Longinelli, A. y Craig, H. (1967). Oxygen-18 variations in sulfate ions in sea water and saline lakes. *Science* 156, 56-59.
- Lowenstein, T. K. y Spencer, R. J. (1990). Syndepositional origin of potash evaporites: Petrographic and fluid inclusion evidence. *American Journal of Science* 290, 1-42.
- Lowenstein, T. K., Spencer, R. J. y Zhang, P. (1989). Origin of ancient potash evaporites; clues from the modern nonmarine Qaidam Basin of western China. *Science*. 245(4922), 1090-1092.
- Lutz, F. (1975). Die verteilung von bromid in Gezüchteten und natürlichen chloridischen Salzmineralen. Tesis doctoral, Eberhard-Karls-University.
- Lloyd, R. M. (1967). Oxygen-18 composition of oceanic sulfate. *Science* 156, 1228-1231.
- Lloyd, R. M. (1968). Oxygen isotope behavior in the sulphate-water system. *Journal of Geophysical Research* 73(18), 6099-6110.

- Maiklem, W. R. (1971). Evaporative drawdown - a mechanism for water-level lowering and diagenesis in the Elk Point basin. *Bulletin of Canadian Petroleum Geology* 19(2), 487-503.
- Marín, A. (1929). Plan de investigación de la cuenca potásica del noreste de España. *Boletín del Instituto Geológico y Minero de España* 11 (3ª Ser.), 73-86.
- Marín, A. (1932). Sondeos de investigación de sales potásicas. *Boletín de Sondeos* 3, 29-99.
- Marín, A. (1945). La depresión del Ebro, la tectónica y los yacimientos minerales. *Boletín del Instituto Geológico y Minero de España* 57, 9-57.
- Marín, A. (1950). "Minería de la potasa". *Potasas Españolas S.A*, 206.
- Marshall, A. T. (1977). Iso-atomic droplets as models for the investigation of parameters affecting X-ray microanalysis of biological specimens. *Micron*. 8, 193-200.
- Marshall, A. T. (1988). Progress in quantitative X-ray microanalysis of frozen-hydrated bulk biological samples. *Jour. Electron Microsc. Tech.* 9, 254-266.
- Martínez Díaz, J. J. y Hernández Enrile, J. L. (1992). Fracturación y control tectosedimentario neógeno en el borde sureste de la Cuenca de Lorca. *Boletín Geológico y Minero* 103(6), 971-983.
- Masachs, V. (1952). La edad, el origen y los movimientos de las sales paleógenas de la Cuenca del Ebro. *Memorias y Comunicaciones del Instituto Geológico Prov.* 9, 51-65.
- Mavrogenes, J. A., Bodnar, R. J., Anderson, A. J., Bajt, S., Sutton, S. R. y Rivers, M. L. (1995). Assessment of the uncertainties and limitations of quantitative elemental analysis of individual fluid inclusions using synchrotron X-ray fluorescence (SXRF). *Geochimica et Cosmochimica Acta* 59(19), 3987-3995.
- Mazor, F. (1962). Radon and radium content of some Israeli water sources and a hypothesis on underground reservoirs of brines, oils and gases in the Rift Valley. *Geochimica et Cosmochimica Acta* 26, 765-786.
- McCaffrey, M. A., Lazar, B. y Holland, H. D. (1987). The evaporation path of seawater and the coprecipitation of Br- and K+ with halite. *Journal of Sedimentary Petrology* 57(5), 928-937.
- McCaig, A. M., Wayne, D. M., Marshall, J. D., Banks, D. y Henderson, I. (1995). Isotopic and fluid inclusion studies of fluid movement along the Gavarnie thrust, Central Pyrenees: reaction fronts in carbonate mylonites. *American Journal of Sciences* 295, 309-343.
- McIntosh, R. A. y Wardlaw, N. C. (1968). Barren halite bodies in the sylvinitic mining zone at Esterhazy, Saskatchewan. *Canadian Journal of Earth Sciences* 5, 1221-1235.
- Meiburg, P. y Kaever, M. (1986). Das Tertiär zwischen Egge und Weser. En: *Nordwestdeutschland im Tertiär* (ed. H. Tobien), pp. 42-86. Borntraeger.

- Menez, B., Philippot, P., Chevallier, P., Gibert, F., Legrand, F. y Populus, P. (1997). Synchrotron X-ray microfluorescence analysis of individual fluid inclusions: $(K\alpha/K\beta)_i$ used as an absorption correction term for concentration estimates. *XIV-ECROFI, Abstracts Volume*, 207-208.
- Metzger, F. W., Kelly, W. C., Nesbitt, B. E. y Essene, E. J. (1977). Scanning electron microscopy of daughter minerals in fluid inclusions. *Economical Geology* 72, 141-152.
- Meybeck, M. (1979). Concentration des eaux fluviales en elements majeurs et apports en solution aux oceans. *Révista de Géologie et de Géographie Physique* 21, 215-246.
- Miller, M. F. y Shepherd, T. J. (1984). The determination of lead in fluid inclusions using voltammetric trace analysis: an exploratory investigation. *Chemical Geology* 42, 249-259.
- Miralles, L. y Sans, M. (1996). Fábrica de rocas salinas en una zona de cizalla (anticlinal de Súria, Barcelona). *Geogaceta* 20(4), 763-766.
- Mizutani, Y. y Rafter, T. A. (1973). Isotopic behavior of sulphate oxygen in the bacterial reduction of sulphate. *Geochemical Journal* 6, 183-191.
- Moissette, A., Dubessy, J., Boiron, M. C., Fabre, C., Mauchien, P. y Lacour, J. L. (1997). Laser ablation OES and its application to individual fluid inclusion analysis: state of the art. *XIV-ECROFI, Abstracts Volume*, 211-212.
- Montenat, C., Ott d'Estevou, P. y Delort, T. (1990). Le bassin de Lorca. En: *Les Bassins Néogènes du Domaine Bétique Oriental, Espagne.*, Vol. 12-13 (ed. C. Montenat), pp. 261-280. Doc. et Trav. IGAL.
- Montenat, C., Ott d'Estevou, P., Larouzière, F. D. y Bedu, P. (1987). Originalité Géodynamique des bassins Néogènes du domaine Bétique Oriental (España). *Notes et Memoires; TOTAL, Compagnie Francaise des Pétroles*. 21, 11-49.
- Moore, G. R. (1980). A chemical and isotopic study of fluid inclusions from the Northern Pennines orefield. Tesis doctoral, University of Durham (UK).
- MOPU. (1990). Análisis de calidad de aguas. En: MOPU (Dirección general de obras hidráulicas).
- Moreton, R. B., Echlin, P., Grupta, B. L. y Hall, T. A. (1974). Preparation of frozen-hydrated tissue sections for X-ray microanalysis in the scanning electron microscope. *Nature* 247, 113-114.
- Morgan, A. J. y Davies, T. W. (1982). An electron microprobe study of the influence of beam current density on the stability of detectable elements in mixed-salts (isoatomic) microdroplets. *Journal of Microscopy* 125, 103-116.
- Müller, D. W. (1987). Strontium 87/86 as fluid tracer in late Miocene evaporites of the Tyrrhenian Sea (western Mediterranean sea). *Eos, Transactions, American Geophysical Union* 68(44), 1320.

- Müller, D. W. y Mueller, P. A. (1991). Origin and age of the Mediterranean messinian evaporites: implications from Sr isotopes. *Earth and Planetary Science Letters* 107, 1-12.
- Muñoz, J. A. (1992). Evolution of a continental collision belt: ECORS-Pyrenees crustal balanced cross-section. En: *Thrust Tectonics* (ed. K. R. McClay), pp. 235-246. Chapman & Hall.
- Nakai, N. y Jensen, M. L. (1964). The kinetic isotope effect in the bacterial reduction and oxidation of sulfur. *Geochimica et Cosmochimica Acta* 28, 1893-1912.
- Nambu, M. y Sato, T. (1981). The analysis of fluid inclusions in the microgram range with an ion microanalyzer. *Bulletin de Minéralogie* 104, 827-833.
- Nambu, M., Sato, T., Ohmori, Y. y Hayakawa, N. (1977). A trial on the microanalysis of fluid inclusions. *Minería y Geología* 58, 40.
- Neev, D. y Emery, K. O. (1967). The Dead Sea, depositional processes and environments of evaporites. *Israel Geological Survey Bulletin* 41, 1-147.
- Nei, T. (1975). Application of cryo scanning electron microscopy in biology. *Cryo-biology* 12, 588.
- Nesteroff, W. D. y Glaçon, G. (1977). Le caractère rythmique des évaporites messiniennes en Méditerranée orientale (coupe d'Eraclea Minoa, Sicile). *Bulletin de la Société Géologique de France* 19, 489-500.
- Nielsen, H. (1967). Sulfur isotopes in the Rhinegraben evaporite sulfates. *Abhandlungen des Geologischen Landesamtes in Baden-Württemberg* 6, 27-29.
- Nielsen, H. y Ricke, W. (1964). Schwefel-Isotopenverhältnisse von Evaporiten aus Deutschland; Ein Beitrag zur Kenntnis von $\delta^{34}\text{S}$ im Meerwasser-sulfat. *Geochimica et Cosmochimica Acta* 28, 577-591.
- Nissenbaum, A. (1975). The microbiology and biogeochemistry of the Dead Sea. *Microbial Ecology* 2, 139-161.
- Nissenbaum, A. (1977). Minor and trace elements in Dead Sea water. *Chemical Geology* 19, 99-111.
- Ortí, F. (1989). Evaporitas marinas. En: *Sedimentología*, Vol. 2 (ed. CSIC), pp. 89-177. CSIC.
- Ortí, F., Pueyo, J. J. y Rosell, L. (1985). La halite du bassin potassique sud-pyrénéen (Eocène supérieur, Espagne). *Bulletin de la Société Géologique de France* 8(6), 863-872.
- Ortí, F. y Salvany, J. M. (1990). Formaciones evaporíticas de la Cuenca del Ebro y cadenas periféricas, y de la zona de levante. Nuevas aportaciones y guía de superficie. En: ENRESA y GPPG, 366 pp.

- Parafiniuk, J. y Halas, S. (1997). Sulfur- and oxygen-isotope composition as the genetic indicator for celestine from the Miocene evaporites of the Carpathian Foredeep. *Slovak Geological Magazine* 3(2), 131-134.
- Parafiniuk, J., Kowalski, W. y Halas, S. (1994). Stable isotope geochemistry and the genesis of the Polish native sulphur deposits- a review. *Geological Quarterly* 38(3), 473-496.
- Pauwels, H., Fouillac, C. y Fouillac, A.-M. (1993). Chemistry and isotopes of deep geothermal saline fluids in the Upper Rhine Graben: Origin of compounds and water-rock interactions. *Geochimica et Cosmochimica Acta* 57, 2737-2749.
- Pérez Lorente, F., Rodríguez Estrella, T., Serrano, F., Mancheño, M. A., Guillén Mondéjar, F., Arana, R., López Aguayo, F. y Fernández Tapia, M. T. (1992). Contribución a la bioestratigrafía Neógena de la cuenca de Lorca (Murcia). *III Congreso Geológico de España y VIII Congreso Latinoamericano de Geología* 1, 185-190.
- Peryt, T. (1997). En: Informe no publicado del Panstwowy Instytut Geologiczny.
- Peryt, T. M., Halas, S., Karoli, S. y Peryt, D. (1997a). Zapis izotopowy zmian środowiskowych podczas depozycji gipsów badenskich w Kobericach koło Opawy. *Przegląd Geologiczny* 45(8), 807-810.
- Peryt, T. M. y Jasionowski, M. (1994). In situ formed and redeposited gypsum breccias in the Middle Miocene Badenian of southern Poland. *Sedimentary Geology* 94, 153-163.
- Peryt, T. M., Karoli, S., Peryt, D., Petrichenko, O. I., Gedl, P., Narkiewicz, W., Durkovicová, J. y Dobieszynska, Z. (1997b). Westernmost occurrence of the middle Miocene Badenian gypsum in central Paratethys (Koberice, Moravia, Czech Republic). *Slovak Geological Magazine* 3(2), 105-120.
- Peryt, T. M. y Kasprzyk, A. (1992). Carbonate-evaporite sedimentary transitions in the Badenian (middle Miocene) basin of southern Poland. *Sedimentary Geology* 76, 257-271.
- Peryt, T. M. y Kovalevich, V. M. (1997). Association on redeposited salt breccias and potash evaporites in the lower Miocene of Stebnyk (Carpathian foredeep, West Ukraine). *Journal of Sedimentary Research* 67(5), 913-922.
- Peryt, T. M., Peryt, D., Szaran, J., Halas, S. y Jasionowski, M. (1998a). O poziomie anhydrytowym Badenu w otworze wiertniczym Ryszkowa wola 7 K. Jarosławia (SE Polska). *Biuletyn Panstwowego Instytutu Geologicznego* 379, 61-80.
- Peryt, T. M., Pierre, C. y Gryniv, S. P. (1998b). Origin of polyhalite deposits in Zechstein (Upper Permian) Zdrada platform (northern Poland). *Sedimentology* 45, 565-578.
- Petrichenko, O. I. (1973). Methods of study of inclusions in minerals of saline deposits. En: *Naukova Dumka*, Vol. Fluid Inclusion Res. Proc. of COFFI 12 (1979) (ed. E. R. y. A. Koslowski), pp. 90.
- Petrichenko, O. I., Peryt, T. M. y Poberegsky, A. V. (1997). Peculiarities of gypsum sedimentation in the Middle Miocene Badenian evaporite basin of Carpathian Foredeep. *Slovak Geological Magazine* 3(2), 91-104.

- Pierre, C. (1982). Teneurs en isotopes stables (^{18}O , ^2H , ^{13}C , ^{34}S) et conditions de genèse des évaporites marines: Application à quelques milieux actuels et au Messinien de la Méditerranée. Tesis doctoral, Université de Paris-Sud Centre d'Orsay.
- Pierre, C. (1983). Polyhalite replacement after gypsum at Ojo de Liebre lagoon (Baja California, Mexico): An early diagenesis by mixing of marine brines and continental waters. En: *6th International Symposium on Salt*, Vol. 1 (ed. B. C. Schreiber), pp. 257-265. Salt Institute.
- Pierre, C. (1985). Isotopic Evidence for the dynamic Redox cycle of dissolved sulphur compounds between free and interstitial solutions in marine salt pans. *Chemical Geology* 53, 191-196.
- Pierre, C. y Rouchy, J. M. (1990). Sedimentary and diagenetic evolution of Messinian evaporites in the Tyrrhenian sea (ODP LEG 107, sites 652, 653, and 654): Petrographic, mineralogical, and stable isotope records. En: *Proceedings of the Ocean Drilling Program, Scientific Results*, Vol. 107 (ed. K. A. Kastens, J. Mascle y e. al.), pp. 187-201.
- Pierre, C., Taberner, C., Urquiola, M. M. y Pueyo, J. J. (1994). Sulphur and oxygen isotopic composition of sulphates in hypersaline environments, as markers of redox depositional versus diagenetic changes. *Mineralogical Magazine* 58A, 724-725.
- Pierre, C., Utrilla, R., Ortí, F. y Pueyo, J. J. (1984). Preliminary stable isotope investigation in carbonates and gypsum from the coastal Salina of Bonmatí (Santa Pola, Alicante, Spain). *Revista d'Investigacions Geològiques* 38-39, 229-235.
- Platt, J. P. y Vissers, R. L. M. (1989). Extensional collapse of thickened continental lithosphere: A working hypothesis for the Alboran Sea and Gibraltar arc. *Geology* 17, 540-543.
- Playà, E. (1998). Les evaporites de les conques bètiques marginals (Fortuna-Lorca, Miocè Superior): Comparació amb altres conques mediterrànies. Tesis doctoral, Universitat de Barcelona.
- Playà, E., Ortí, F. y Rosell, L. (1995). Composición isotópica de los yesos Messinienses de la cuenca de Fortuna (Murcia): implicaciones sedimentológicas. *Geogaceta* 18, 63-66.
- Playà, E., Rosell, L. y Ortí, F. (1997). Geoquímica Isotópica ($\delta^{34}\text{S}$, $^{87}\text{Sr}/^{86}\text{Sr}$) y contenidos en estroncio de las evaporitas messinienses de la cuenca de Sorbas (Almería). En: *Avances en el conocimiento del Terciario Ibérico* (ed. J. P. C. y J. Morales), pp. 161-164.
- Poutoukis, D. (1991). Hydrochimie, teneurs isotopiques et origine des saumures associées aux gisements de potasse d'Alsace. Tesis doctoral, Université de Paris-Sud Centre d'Orsay.
- Pueyo, J. J. (1975a). Estudio petrológico y geoquímico de los yacimientos potásicos de Cardona, Suria, Sallent y Balsareny (Barcelona, España). Tesis doctoral, Universidad de Barcelona.

- Pueyo, J. J. (1975b). Sedimentary structures observed in salt of the Catalanian potash basin. *IX Congr. Int. de Sedimentologie, Nice Livret guide Exc. 20, Le bassin Tertiaire et les gisements de potasse*, 59-64.
- Pueyo, J. J. (1977). El bromo y el rubidio como indicadores genéticos en las evaporitas de la cuenca potásica catalana. *Revista del Instituto de Investigaciones Geológicas* 32, 77-86.
- Pueyo, J. J. y San Miguel, A. (1974). Características petrológicas de las sales sódicas, potásicas y magnésicas de la Cuenca Potásica Catalana. *Instituto de Investigaciones Geológicas Diputación Provincial, Universidad de Barcelona* 29, 23-49.
- Pueyo, J. J., Vaquer, R. y Traveria, A. (1974). Distribución de sulfatos en los yacimientos de la cuenca potásica catalana. *Instituto de Investigaciones Geológicas Diputación Provincial Universidad de Barcelona* 29, 51-65.
- Puigdefàbregas, C. (1975). La sedimentación molásica en la cuenca de Jaca. *Revista Pirineos CSIC monografías del Instituto de Estudios Pirenaicos*(104), 188.
- Quinton, P. (1978). Ultramicroanalysis of biological fluids with Energy Dispersive X-ray Spectrometry. *Micron*. 9, 57-69.
- Ramsey, M. H., Coles, B. J., Wilkinson, J. J. y Rankin, A. H. (1992). Single fluid inclusion analysis by laser ablation inductively coupled plasma atomic emission spectrometry: quantification and validation. *Journal of Analytical Atomic Spectrometry* 7, 587-593.
- Rankin, A. H., Ramsey, M. H., Coles, B., Van Langevelde, F. y Thomas, C. R. (1992). The composition of hypersaline, iron-rich granitic fluids based on laser-ICP and Synchrotron-XRF microprobe analysis of individual fluid inclusions in topaz, Mole granite, eastern Australia. *Geochimica et Cosmochimica Acta* 56, 67-79.
- Rebelle, M. (1986). Sédimentologie, géochimie et palynologie du bassin évaporitique du Zechstein à partir de données de sub-surface (Mer du Nord, Hesse- R.F.A.). Nature et genèse des faciès confinés. Tesis doctoral, Document BRGM N°8.
- Rees, C. E., Jenkins, W. J. y Monster, J. (1978). The sulphur isotopic composition of ocean water sulphate. *Geochimica et Cosmochimica Acta* 42, 377-381.
- Riba, O. (1967). Resultados de un estudio sobre el Terciario continental de la parte Este de la Depresión Central Catalana. *Acta Geológica Hispánica* 2, 3-8.
- Ricchiuto, T. E. y McKenzie, J. A. (1978). Stable isotopic investigation of Messinian sulfate samples from DSDP Leg 42A, Eastern Mediterranean sea. *Initial Reports of the Deep Sea Drilling Project* 42, 657-660.
- Rick, B. (1990). Sulphur and oxygen isotopic composition of Swiss Gipskeuper (Upper Triassic). *Chemical Geology* 80, 243-250.
- Ríos, J. M. (1963). Materiales salinos del suelo español. *Memorias del Instituto Geológico y Minero de España* 64, 166.

- Rodríguez-Estrella, T., Mancheño, M. A., Guillén Mondéjar, F., López Aguayo, F., Arana, R., Fernández Tapia, M. T. y Serrano, F. (1992). Tectónica y sedimentación Neógena en la Cuenca de Lorca (Murcia). *III Congreso Geológico de España y VIII Congreso Latinoamericano de Geología* 1, 201-206.
- Roedder, E. (1962). Ancient fluids in crystals. *Scientific American* 207(4), 38-47.
- Roedder, E. (1984a). *Fluid Inclusions*. Mineralogical Society of America.
- Roedder, E. (1984b). The fluids in salt. *American Mineralogist* 69, 413-439.
- Roedder, E. (1990). Fluid inclusion analysis - Prologue and epilogue. *Geochimica et Cosmochimica Acta* 54, 495-507.
- Roedder, E., d'Angelo, W. M., Dorrzapf, A. F. y Aruscavage, J. a. P. J. (1987). Composition of fluid inclusions in permian salt beds, Palo Duro basin, Texas, U.S.A. *Chemical Geology* 61, 79-90.
- Rögl, F., Steininger, F. F. y Müller, C. (1978). Middle Miocene salinity crisis and paleogeography of the Paratethys (Middle and Eastern Europe). *Initial Reports of DSDP* 42(Part 1), 985-990.
- Rosell, L. (1983). Estudi petrològic, sedimentològic i geoquímic de la formació de sals potàssiques de Navarra (Eocè superior), Universitat de Barcelona.
- Rosell, L., Ortí, F., Kasprzyk, A., Playà, E. y Peryt, T. M. (1998). Strontium geochemistry of miocene primary gypsum: Messinian of southeastern Spain and Sicily and Badenian of Poland. *Journal of Sedimentary Research* 68(1), 63-79.
- Rosell, L. y Pueyo, J. J. (1984). Características geoquímicas de la formación de sales potásicas de Navarra (Eoceno superior). Comparación con la cuenca potásica catalana. *Acta Geológica Hispánica* 19(2), 81-95.
- Rosell, L. y Pueyo, J. J. (1997). Second marine evaporitic phase in the south Pyrenean Foredeep: The Priabonian potash basin (Late Eocene; autochthonous-allochthonous Zone). En: *Sedimentary Deposition in Rift and Foreland Basins in France and Spain (Paleogene and Lower Neogene)* (ed. G. B. y. C. Schreiber), pp. 358-387. Columbia University Press.
- Ross, D. A. (1972). Red Sea hot brine area: revisited. *Science* 175, 1455-1456.
- Rouchy, J. M. (1982). *La Genèse des Evaporites Messiniennes de Méditerrané*. Compagnie Française des Pétroles et de la Société Nationale ELF-Aquitaine.
- Rouchy, J. M. y Saint Martin, J. P. (1992). Late Miocene events in the Mediterranean as recorded by carbonate-evaporite relations. *Geology* 20, 629-632.
- Rouchy, J. M., Taberner, C., Blanc-Valleron, M. M., Sprovieri, R., Russell, M., Pierre, C., Di Stefano, E., Pueyo, J. J., Caruso, A., Dinarés-Turell, J., Gomis-Coll, E., Wolff, G., Cespuglio, G., Ditchfield, P., Pestrea, S., Combourieu-Nebout, N., Santisteban, C. y Grimalt, J. O. (1998). Sedimentary and diagenetic

markers of the restriction in a marine basin: the Lorca basin (SE Spain) during the Messinian. *Sedimentary Geology* 121, 23-55.

Rouchy, J. M., Taberner, C., Blanc-Valleron, M. M., Sprovieri, R., Russell, M., Pierre, C., Di Stefano, E., Pueyo, J. J., Caruso, A., Dinarès-Turell, J., Wolf, G., Grimalt, J., Marshall, J., Santisteban, C. y Pestrea, S. (1997). Depositional changes during the Messinian salinity crisis in a marginal basin (Lorca basin, SE Spain): Records of local and global forcings. *Neogene Mediterranean Paleoceanography. Abstract Volume*, 62-63.

Ryan, C. G. (1995). Development of PIXE and proton microprobe and the non-destructive analysis of fluid inclusions in minerals. *Boletín de la Sociedad Española de Mineralogía* 18(1), 214-215.

Ryan, C. G., Cousens, D. R., Heinrich, C. A., Griffin, W. L., Sie, S. H. y Mernagh, T. P. (1991). Quantitative PIXE microanalysis of fluid inclusions based on a layered yield model. *Nuclear Instruments and Methods in Physics Research B*54, 292-297.

Ryan, C. G., Heinrich, C. A. y Mernagh, T. P. (1993). PIXE microanalysis of fluid inclusions and its application to study ore metal segregation between magmatic brine and vapor. *Nuclear Instruments and Methods in Physics Research B*77, 463-471.

Rye, D. M. y Bradbury, H. J. (1988). Fluid flow in the crust: an example from a Pyrenean thrust ramp. *American Journal of Sciences* 288, 197-235.

Sáez, A. y Salvany, J. M. (1990). Las formaciones evaporíticas de Barbastro y Puente La Reina (Eoceno Superior-Oligoceno basal de la Cuenca Surpirenaica). En: *Formaciones evaporíticas de la cuenca del Ebro y cadenas periféricas, y de la zona de Levante* (ed. F. Ortí y J. M. Salvany). ENRESA.

Sáez, A., Vergés, J., Taberner, C., Pueyo, J. J., Muñoz, J. A. y Busquets, P. (1991). Eventos evaporíticos paleógenos en la cuenca antepaís surpirenaica: ¿causas climáticas-causas tectónicas? En: Vol. 5 (ed. F. Colombo), pp. 85. I Congreso del Grupo Español del Terciario.

Sanford, W. E. y Wood, W. W. (1991). Brine evolution and mineral deposition in hydrologically open evaporate basins. *American Journal of Science* 291, 687-710.

Sans, M., Muñoz, J. A. y Vergés, J. (1996a). Triangle zone and thrust wedge geometries related to evaporitic horizons (southern Pyrenees). *Bulletin of Canadian Petroleum Geology* 44(2), 375-384.

Sans, M., Sánchez, A. L. y Santanach, P. (1996b). Internal structure of a detachment horizon in the most external part of the Pyrenean fold and thrust belt (northern Spain). En: *Salt Tectonics*, Vol. 100 (ed. D. J. B. y I. D. G. I. Alsop), pp. 65-76. Geological Society Special Publication.

Sans, M. y Vergés, J. (1995). Fold development related to contractional salt tectonics: southeastern pyrenean thrust front, Spain. En: *Salt Tectonics: a Global Perspective*, Vol. 65 (ed. M. P. A. Jackson, D. G. Roberts y S. Snelson), pp. 369-378. AAPG Memoir.

- Sanz de Galdeano, C. (1990). Geologic evolution of the Betic cordilleras in the western Mediterranean, Miocene to the present. *Tectonophysics* 172, 107-119.
- Sasaki, A. (1972). Variation in sulphur isotopic composition of oceanic sulphate. En: *24th International Geology Congress*, Vol. 10, pp. 342-345.
- Sato, T., Ohmori, Y. y Nambu, M. (1984). Quantitative analysis of metallic ions associated with fluid inclusions. *Springer Ser. Chem. Physics*, 36, *Second Ion Mass Spectrom, SIMS 4*, 471-474.
- Schmidt di Frieberg, G. (1962). Introduction à la géologie pétrolière de la Sicile. *Rev. Inst. Fr. Pétrole* 27, 635-667.
- Shan, L., Liu, X. D. y Wu, W. J. (1985). Uranium exploration by radon measurement in inclusions (abstr.). *Geological Society of America. Abstracts with Programs* 17, 713.
- Shatkay, M. y Magaritz, M. (1987). Dolomitization and sulfate reduction in the mixing zone between brine and meteoric water in the newly exposed shores of the Dead Sea. *Geochimica et Cosmochimica Acta* 51, 1135-1141.
- Shearman, D. J. (1966). Origin of marine evaporites by diagenesis. *Institution of Mining and Metallurgy Transactions* 75, 208-215.
- Shepherd, T. J., Ayora, C., Cendón, D.I., Chenery, S. R., Moissette, A. (1998). Quantitative solute analysis of single fluid inclusions in halite by LA-ICP-MS and cryo-SEM-EDS: complementary techniques. *European Journal of Mineralogy* 10, 1097-1108.
- Shepherd, T. J. y Chenery, S. R. (1995). Laser ablation ICP-MS elemental analysis of individual fluid inclusions: An evaluation study. *Geochimica et Cosmochimica Acta* 59(19), 3997-4007.
- Slaczka, A. y Kolasa, K. (1997). Resedimented salt in the Northern Carpathians Foredeep (Wieliczka, Poland). *Slovak Geological Magazine* 3(2), 135-155.
- Sonnenfeld, P. y Finetti, I. (1985). Messinian evaporites in the Mediterranean: A model of continuous inflow and outflow. En: *Geological Evolution of the Mediterranean Basin* (ed. D. J. S. y F.-C. Wezel), pp. 347-353. Springer-Verlag.
- Spencer, R. J. y Hardie, L. A. (1990). Control of seawater by mixing of rivers waters and mid-ocean ridge hydrothermal brines. *Fluid-Mineral Interactions: A Tribute to H.P. Eugster. The Geochemical Society, Special Publication* 2, 409-419.
- Stein, C. L. y Krumhansl, J. L. (1988). A model for the evolution of brines in salt from the lower Salado Formation, southeastern New Mexico. *Geochimica et Cosmochimica Acta* 52, 1037-1046.
- Steinhorn, I., Assaf, G., Gat, J. R., Nishry, A., Nissenbaum, A., Stiller, M., Beyth, M., Neev, D., Garber, R., Friedman, G. M. y Weiss, W. (1979). The Dead Sea: Deepening of the Mixolimnion signifies the overture to overturn of the water column. *Science* 206, 55-57.

- Sweeney, R. E. y Kaplan, I. R. (1980). Diagenetic sulfate reduction in marine sediments. *Marine Chemistry* 9, 165-174.
- Taberner, C. (1982). Evolución ambiental y diagenética de los depósitos del Terciario inferior (Paleógeno y Eoceno de la cuenca de Vic). Tesis doctoral, Universitat de Barcelona.
- Thode, H. G. y Monster, J. (1965). Sulfur-isotope geochemistry of petroleum, evaporites, and ancient seas. *AAPG Bulletin*, 363-373.
- Thode, H. G., Monster, J. y Dunford, H. B. (1961). Sulphur isotope geochemistry. *Geochimica et Cosmochimica Acta*, 25(3), 159-174.
- Travé, A., Labaume, P., Calvet, F. y Soler, A. (1997). Sediment dewatering and pore fluid migration along thrust faults in a foreland basin inferred from isotopic and elemental geochemical analyses (Eocene southern Pyrenees, Spain). *Tectonophysics* 282, 375-398.
- Travé, A., Labaume, P., Calvet, F., Soler, A., Tritlla, J., Buatier, M., Potdevin, J. L., Séguret, M., Raynaud, S. y Briquieu, L. (1998). Fluid migration during Eocene thrust emplacement in the south Pyrenean foreland basin (Spain): an integrated structural, mineralogical and geochemical approach. En: *Cenozoic Foreland Basins of Western Europe*, Vol. 134 (ed. C. P. A. Mascle, H. P. Luterbacher, y M. Fernández), pp. 163-188. Geological society special publications.
- Urquiola, M. M. (1994). Aplicación de indicadores geoquímicos al estudio de las facies anóxicas del Eoceno superior de la cuenca de antepaís Surpirenaica (Sector Oriental). Tesis doctoral, Universitat de Barcelona.
- Usiglio, J. (1849). Analyse de l'eau de la Méditerranée sur les côtes de France. *Annalen der Chemie* 27, 92-107, 172-191.
- Utrilla, R. (1985). Estudi sedimentològic i geoquímic de les Salines de la Trinitat (Delta de l'Ebre) i San Pedro del Pinatar (Mar Menor). Tesis de licenciatura, Universitat de Barcelona.
- Utrilla, R. (1989). Les composicions isotòpiques ($\delta^{18}\text{O}$, $\delta^{34}\text{S}$) del sulfat, com a indicadors de l'origen de les evaporites del Mesozoic i del Cenozoic de la península Ibèrica i les Illes Balears. Tesis doctoral, Universitat de Barcelona.
- Utrilla, R., Ortí, F., Pierre, C. y Pueyo, J. J. (1991). Composición isotópica de las evaporitas Terciarias continentales de la cuenca del Ebro: relación con los ambientes deposicionales. *Revista de la Sociedad Geológica de España* 4(3-4), 353-360.
- Utrilla, R., Pierre, C., Orti, F. y Pueyo, J. J. (1992). Oxygen and sulphur isotope compositions as indicators of the origin of Mesozoic and Cenozoic evaporites from Spain. *Chemical Geology (Isotope Geoscience Section)* 102, 229-244.
- Valyashko, M. G. (1956). Geochemistry of bromine in the processes of salt deposition and the use of the bromine content as a genetic and prospecting criterion. *Geochemistry* 6, 570-589.

- Vanko, D. A., Sutton, S. R., Rivers, M. L. y Bodnar, R. J. (1993). Major-element ratios in synthetic fluid inclusions by synchrotron X-ray fluorescence microprobe. *Chemical Geology* 109, 125-134.
- Van't Hoff, J. H. (1912). *Untersuchungen über die Bildungsverhältnisse der ozeanischen Salzablagerungen, insbesondere das Stassfurter Salzlagere*. Akad. Verlags. ges.
- Von Matter, A., Peters, T. y Ramseyer, K. (1987). $^{87}\text{Sr}/^{86}\text{Sr}$ -Verhältnisse und Sr-gehalte von tiefengrundwässern, mineralien sowie gesteinen aus dem kristallin und der trias der Nordschweiz. *Eclogae Geologicae Helvetiae* 80(2), 579-592.
- Wardlaw, N. C. (1972). Unusual marine evaporites with salts of calcium and magnesium chloride in Cretaceous basins of Sergipe, Brazil. *Economic Geology* 67, 156-168.
- Wardlaw, N. C. y Schwerdtner, W. M. (1966). Halite-Anhydrite seasonal layers in the Middle Devonian Prairie Evaporite formation, Saskatchewan, Canada. *Geological Society of America Bulletin* 77, 331-342.
- Wendt, R. H. y Fassel, V. A. (1965). Induction-coupled plasma spectrometric excitation source. *Anal. Chem.* 37, 920-922.
- Wild, H. (1982). Das salzlager des mitterleren muschelkalkes in süddeutschen Raum, Schichtenfolge und Ausbildung-Rückschlüsse auf die Genese. En: *5th International Symposium on Salt*, Vol. 1 (ed. A. H. Coogan y L. Hauber), pp. 185-192. The Northern Ohio Geological Society.
- Wilkinson, J. J., Rankin, A. H., Mulshaw, S. C., Nolan, J. y Ramsey, M. H. (1994). Laser ablation-ICP-AES for determination of metals in fluid inclusions: An application to the study of magmatic ore fluids. *Geochimica et Cosmochimica Acta* 58(3), 1133-1146.
- Williams-Stroud, S. C. (1994). Solution to the paradox? results of some chemical equilibrium and mass balance calculations applied to the Paradox basin evaporite deposit. *American Journal of Science* 294, 1189-1228.
- Wopenka, B., Pasteris, J. D. y Freeman, J. J. (1990). Analysis of individual fluid inclusions by Fourier transform infrared and Raman microspectroscopy. *Geochimica et Cosmochimica Acta* 54, 519-533.
- Worden, R. H., Smalley, P. C. y Fallick, A. E. (1997). Sulfur cycle in buried evaporites. *Geology* 25(7), 643-646.
- Zak, I., Sakai, H. y Kaplan, I.R. (1980). Factor controlling the $^{18}\text{O}/^{16}\text{O}$ and $^{34}\text{S}/^{32}\text{S}$ isotope ratios of ocean sulfates, evaporites and interstitial sulfates from modern deep sea sediments. En: *Isope Marine Chemistry* (ed. E.D. Goldberg, Y. Horibe y K. Suruhashi), pp. 339-373. Rokakuho, Tokyo.
- Zs-Nagy, I., Lustyik, G. y Bertoni-Freddari, C. (1982). Intracellular water and dry mass content as measured in bulk specimens by energy-dispersive X-ray microanalysis. *Tissue Cel* 14, 47-60.

Zs-Nagy, I., Pieri, C., Giuli, C., Bertoni-Freddari, C. y Zs-Nagy, V. (1977). Energy-dispersive X-ray microanalysis of the electrolytes in biological bulk specimen, I. Specimen preparation, beam penetration and quantitative analysis. *Ultrastruct. Res.* 58, 22-33.