

VII. BIBLIOGRAFÍA**A**

- Abellán MT, Adell A, Honrubia MA, Mengod G, Artigas F (2000a) GABA(B)-RI receptors in serotonergic neurons: effects of baclofen on 5-HT output in rat brain. *Neuroreport* 11:941-945.
- Abellán MT, Jolas T, Aghajanian GK, Artigas F (2000b) Dual control of dorsal raphe serotonergic neurons by GABA(B) receptors. *Electrophysiological and microdialysis studies. Synapse* 36:21-34.
- Abi-Dargham A, Mawlawi O, Lombardo I, Gil R, Martinez D, Huang Y, Hwang DR, Keilp J, Kochan L, Van Heertum R, Gorman JM, Laruelle M (2002) Prefrontal dopamine D1 receptors and working memory in schizophrenia. *J.Neurosci.* 22:3708-3719.
- Adell A y Artigas F (1998) A microdialysis study of the *in vivo* release of 5-HT in the median raphe nucleus of the rat. *Br.J.Pharmacol.* 125:1361-137.
- Adell A, Sarna GS, Hutson PH, Curzon G (1989) An *in vivo* dialysis and behavioural study of the release of 5-HT by p-chloroamphetamine in reserpine-treated rats. *Br.J.Pharmacol.* 97:206-212.
- Adell A, Carceller A, Artigas F (1993) *In vivo* brain dialysis study of the somatodendritic release of serotonin in the raphe nuclei of the rat. Effects of 8-hydroxy-2-(di-n-propylamino)tetralin. *J.Neurochem.* 60:1673-1681.
- Adell A, Celada P, Artigas F (2001) The role of 5-HT_{1B} receptors in the regulation of serotonin cell firing and release in the rat brain. *J.Neurochem.* 79:172-182.
- Adell A, Celada P, Abellán MT, Artigas F (2002) Origin and functional role of the extracellular serotonin in the midbrain raphe nuclei. *Brain Res.Brain Res.Rev.* 39:154-180.
- Adham N, Borden LA, Schechter LE, Gustafson EL, Cochran TL, Vaysse PJJ, Weinshank RL, Branchek TA (1993) Cell-Specific Coupling of the Cloned Human 5-HT(1F) Receptor to Multiple Signal Transduction Pathways. *Naunyn-Schmied.Arch.Pharmacol.* 348:566-575.
- Aghajanian GK (1978) Feedback regulation of central monoaminergic neurons: evidence from single cell recording studies. *Essays Neurochem.Neuropharmacol.* 3:1-32.
- Aghajanian GK (1985) Modulation of a transient outward current in serotonergic neurones by alpha 1-adrenoceptors. *Nature* 315:501-503.
- Aghajanian GK y Lakoski JM (1984) Hyperpolarization of serotonergic neurons by serotonin and LSD: studies in brain slices showing increased K⁺ conductance. *Brain Res.* 305:181-185.

- Aghajanian GK y Marek GJ (1997) Serotonin induces excitatory postsynaptic potentials in apical dendrites of neocortical pyramidal cells. *Neuropharmacology* 36:589-599.
- Aghajanian GK y Marek GJ (1999a) Serotonin, via 5-HT_{2A} receptors, increases EPSCs in layer V pyramidal cells of prefrontal cortex by an asynchronous mode of glutamate release. *Brain Res.* 825:161-171.
- Aghajanian GK y Marek GJ (1999b) Serotonin-glutamate interactions: A new target for antipsychotic drugs. *Neuropsychopharmacology* 21:S122-S133.
- Aghajanian GK y Marek GJ (2000) Serotonin model of schizophrenia: emerging role of glutamate mechanisms. *Brain Res. Rev.* 31:302-312.
- Aghajanian GK y VanderMaelen CP (1982) Intracellular recording *in vivo* from serotonergic neurons in rat dorsal raphe nucleus: methodological considerations. *J Histochem Cytochem* 30:813-814.
- Aghajanian GK y Wang RY (1977) Habenular and other midbrain raphe afferents demonstrated by a modified retrograde tracing technique. *Brain Res.* 122:229-242.
- Aguilar J, Soto C, Rivadulla C, Canedo A (2002) The lemniscal-cuneate recurrent excitation is suppressed by strychnine and enhanced by GABA_A antagonists in the anaesthetized cat. *Eur J Neurosci* 16:697-704.
- Andreasen NC, O'Leary DS, Flaum M, Nopoulos P, Watkins GL, Boles Ponto LL, Hichwa RD (1997) Hypofrontality in schizophrenia: distributed dysfunctional circuits in neuroleptic-naïve patients. *Lancet* 349:1730-1734.
- Araneda R y Andrade R (1991) 5-Hydroxytryptamine₂ and 5-hydroxytryptamine_{1A} receptors mediate opposing responses on membrane excitability in rat association cortex. *Neuroscience* 40:399-412.
- Arango V, Underwood MD, Mann JJ (1997) Postmortem findings in suicide victims. Implications for *in vivo* imaging studies. *Ann. N. Y. Acad. Sci.* 836:269-287.
- Arango V, Underwood MD, Mann JJ (2002) Serotonin brain circuits involved in major depression and suicide. *Prog. Brain Res.* 136:443-453.
- Artigas F, Celada P, Laruelle M, Adell A (2001) How does pindolol improve antidepressant action? *Trends Pharmacol. Sci.* 22:224-228.
- Artigas F, Romero L, de Montigny C, Blier P (1996) Acceleration of the effect of selected antidepressant drugs in major depression by 5-HT_{1A} antagonists. *Trends Neurosci.* 19:378-383.

- Arvanov VL, Liang XF, Magro P, Roberts R, Wang RY (1999) A pre- and postsynaptic modulatory action of 5-HT and the 5-HT_{2A}, (2C) receptor agonist DOB on NMDA-evoked responses in the rat medial prefrontal cortex. *Eur. J. Neurosci.* 11:2917-2934.
- Ashby CR, Edwards E, Harkins K, Wang RY (1989) Characterization of 5-hydroxytryptamine3 receptors in the medial prefrontal cortex: A microiontophoretic study. *Eur. J. Pharmacol.* 173:193-196.
- Ashby CR, Minabe Y, Edwards E, Wang RY (1991) 5-HT₃-like receptors in the rat medial prefrontal cortex: an electrophysiological study. *Brain Res.* 550:181-191.
- Ashby CR, Edwards E, Wang RY (1992) Action of serotonin in the medial prefrontal cortex: mediation by serotonin3-like receptors. *Synapse* 10:7-15.
- Ashby CR, Edwards E, Wang RY (1994) Electrophysiological evidence for a functional interaction between 5-HT(1A) and 5-HT(2A) receptors in the rat medial prefrontal cortex: An iontophoretic study. *Synapse* 17:173-181.
- Audet MA, Descarries L, Doucet G (1989) Quantified regional and laminar distribution of the serotonin innervation in the anterior half of adult rat cerebral cortex. *J. Chem. Neuroanat.* 2:29-44.
- Auerbach SB, Minzenberg MJ, Wilkinson LO (1989) Extracellular serotonin and 5-hydroxyindoleacetic acid in hypothalamus of the unanesthetized rat measured by in vivo dialysis coupled to high-performance liquidchromatography with electrochemical detection: dialysate serotonin reflects neuronal release. *Brain Res.* 499:281-290.
- Azmitia EC (1981) The visualization and characterization of 5-HT reuptake sites in the rodent and primate hippocampus. A preliminary study. *J. Physiol.* 77:175-182.
- Azmitia EC y Henriksen SJ (1976) A modification of the Falck-Hillarp technique for 5-HT fluorescence employing hypertonic formaldehyde perfusion. *J. Histochem. Cytochem.* 24:1286-1288.
- Azmitia EC y Segal M (1978) An autoradiographic analysis of the differential ascending projections of the dorsal and median raphe nuclei in the rat. *J. Comp. Neurol.* 179:641-668.
- Azmitia EC, Gannon PJ, Kheck NM, Whitakerazmitia PM (1996) Cellular localization of the 5-HT1A receptor in primate brain neurons and glial cells. *Neuropsychopharmacology* 14:35-46.

B

- Baker DA y Kalivas PW (2000) Cocaine-induced alterations in extracellular glutamate levels: role of cystine/glutamate exchange. *Soc. Neurosci. Abstr.* 26:792-792.

- Baraban JM y Aghajanian GK (1980a) Suppression of firing activity of 5-HT neurons in the dorsal raphe by alpha-adrenoceptor antagonists. *Neuropharmacology* 19:355-363.
- Baraban JM y Aghajanian GK (1980b) Suppression of serotonergic neuronal firing by alpha-adrenoceptor antagonists: evidence against GABA mediation. *Eur. J. Pharmacol.* 66:287-294.
- Baraban JM y Aghajanian GK (1981) Noradrenergic innervation of serotonergic neurons in the dorsal raphe: demonstration by electronic microscopic autoradiography. *Brain Res.* 24:1-11.
- Barbas H y Blatt GJ (1995) Topographically specific hippocampal projections target functionally distinct prefrontal areas in the rhesus monkey. *Hippocampus* 5:511-533.
- Beaulieu C (1993) Numerical data on neocortical neurons in adult rat, with special reference to the GABA population. *Brain Res.* 609:284-292.
- Belin MF, Nanopoulos D, Didier M, Aguera M, Steinbusch H, Verhofstad A, Maitre M, Pujol JF (1983) Immunohistochemical evidence for the presence of gamma-aminobutyric acid and serotonin in one nerve cell. A study on the raphe nuclei of the rat using antibodies to glutamate decarboxylase and serotonin. *Brain Res.* 275:329-339.
- Berendse HW y Groenewegen HJ (1991) Restricted cortical termination fields of the midline and intralaminar thalamic nuclei in the rat. *Neuroscience* 42:73-102.
- Bertolino A, Esposito G, Callicott JH, Mattay VS, Van Horn JD, Frank JA, Berman KF, Weinberger DR (2000) Specific relationship between prefrontal neuronal N-acetylaspartate and activation of the working memory cortical network in schizophrenia. *Am. J. Psychiatry* 157:26-33.
- Blier P y de Montigny C (1994) Current advances and trends in the treatment of depression. *Trends Pharmacol. Sci.* 15:220-226.
- Blier P, de Montigny C, Tardif DE (1987) Short-term lithium treatment enhances responsiveness of postsynaptic 5-HT_{1A} receptors without altering 5-HT_{1A} autoreceptor sensitivity: an electrophysiological study in the rat brain. *Synapse* 1:225-232.
- Blier P, Serrano A, Scatton B (1990) Differential responsiveness of the rat dorsal and median raphe 5-HT systems to 5-HT₁ receptor agonists and p-chloroamphetamine. *Synapse* 5:120-133.
- Blue ME, Yagaloff KA, Mamounas LA, Hartig PR, Molliver ME (1988) Correspondence between 5-HT₂ receptors and serotonergic axons in rat neocortex. *Brain Res.* 453:315-328.
- Bonvento G, Scatton B, Claustre Y, Rouquier L (1992) Effect of local injection of 8-OH-DPAT into the dorsal or median raphe nuclei on extracellular levels

- of serotonin in serotonergic projection areas in the rat brain. *Neurosci. Lett.* 137:101-104.
- Bosker F, Klompmakers A, Westenberg H (1994) Extracellular 5-hydroxytryptamine in median raphe nucleus of the conscious rat is decreased by nanomolar concentrations of 8-hydroxy-2-(di-n-propylamino) tetralin and is sensitive to tetrodotoxin. *J. Neurochem.* 63:2165-2171.
- Bradley PB, Handley SL, Cooper SJ, Key BJ, Barnes NM, Coole JH (editores) (1992) Serotonin, CNS receptors and brain function. Oxford: Pergamon Press.
- Branchereau P, Van Bockstaele EJ, Chan J, Pickel VM (1996) Pyramidal neurons in rat prefrontal cortex show a complex synaptic response to single electrical stimulation of the locus coeruleus region: evidence for antidromic activation and GABAergic inhibition using *in vivo* intracellular recording and electron microscopy. *Synapse* 22:313-331.
- Brodmann K (1909) Vergleichende Lokalisationslehre der Grosshirnhinde. Leipzig.
-
- C**
- Carr DB y Sesack SR (1999) Terminals from the rat prefrontal cortex synapse on mesoaccumbens VTA neurons. *Ann. N. Y. Acad. Sci.* 877:676-678.
- Carr DB y Sesack SR (2000a) GABA-containing neurons in the rat ventral tegmental area project to the prefrontal cortex. *Synapse* 38:114-123.
- Carr DB y Sesack SR (2000b) Projections from the rat prefrontal cortex to the ventral tegmental area: target specificity in the synaptic associations with mesoaccumbens and mesocortical neurons. *J. Neurosci.* 20:3864-3873.
- Carr DB, O'Donnell P, Card JP, Sesack SR (1999) Dopamine terminals in the rat prefrontal cortex synapse on pyramidal cells that project to the nucleus accumbens. *J. Neurosci.* 19:11049-11060.
- Casanovas JM, Berton O, Celada P, Artigas F (2000) *In vivo* actions of the selective 5-HT_{1A} receptor agonist BAY x 3702 on serotonergic cell firing and release. *Naunyn Schmiedebergs Arch. Pharmacol.* 362:248-254.
- Casanovas JM, Hervás I, Artigas F (1999) Postsynaptic 5-HT_{1A} receptors control 5-HT release in the rat medial prefrontal cortex. *Neuroreport* 10:1441-1445.
- Cauli B, Audinat E, Lambolez B, Angulo MC, Ropert N, Tsuzuki K, Hestrin S, Rossier J (1997) Molecular and physiological diversity of cortical nonpyramidal cells. *J. Neurosci.* 17:3894-3906.

- Ceci A, Baschirotto A, Borsini F (1994) The inhibitor effect of 8-OH-DPAT on the firing activity of dorsal raphe neurons in rats is attenuated by lesion of the frontal cortex. *Neuropharmacology* 33:709-713.
- Celada P, Siuciak JA, Tran TM, Altar CA, Tepper JM (1996) Local infusion of brain-derived neurotrophic factor modifies the firing pattern of dorsal raphe serotonergic neurons. *Brain Res.* 712:293-298.
- Chinaglia G, Landwehrmeyer B, Probst A, Palacios JM (1993) Serotonergic terminal transporters are differentially affected in Parkinson's disease and progressive supranuclear palsy: an autoradiographic study with [³H]citalopram. *Neuroscience* 54:691-699.
- Clements JR, Madt JE, Johnson RL, Larson AA, Beitz AJ (1987) Localization of glutamate, glutaminase, aspartate and aspartate aminotransferase in the rat midbrain periaqueductal gray. *Exp. Brain Res.* 67:593-602.
- Clemett DA, Punhani T, Duxon MS, Blackburn TP, Fone KC (2000) Immunohistochemical localisation of the 5-HT_{2C} receptor protein in the rat CNS. *Neuropharmacology* 39:123-132.
- Coccaro EF, Murphy DL (editores) (1990) Serotonin in major psychiatric disorders. Washington: American Psychiatric Press.
- Colbert CM y Johnston D (1996) Axonal action-potential initiation and Na⁺ channel densities in the soma and axon initial segment of subiculum pyramidal neurons. *J. Neurosci.* 16:6676-6686.
- Compan V, Segu L, Buhot MC, Daszuta A (1998) Selective increases in serotonin 5-HT_{1B/1D} and 5-HT_{2A/2C} binding sites in adult rat basal ganglia following lesions of serotonergic neurons. *Brain Res.* 793:103-111.
- Connors BW y Gutnick MJ (1990) Intrinsic firing patterns of diverse neocortical neurons. *Trends Neurosci.* 13:365-366.
- Cortés R, Mengod G, Celada P, Artigas F (1993) p-Chlorophenylalanine increases tryptophan-5-hydroxylase mRNA levels in the rat dorsal raphe: A time course study using *in situ* hybridization. *J. Neurochem.* 60:761-764.
- Cross AJ (1990) Serotonin in Alzheimer-type dementia and other dementing illnesses. *Ann. N. Y. Acad. Sci.* 600:405-415.

D

- Dahlström A y Fuxe K (1964) Evidence for the existence of monoamine-containing neurons in the central nervous system. I Demonstration of monoamines in cell bodies of brain neurons. *Acta Physiol. Scand.* 62, suppl. 232:1-55.

- Danish University Antidepressant Group (1986) Citalopram: clinical effect profile in comparison with clomipramine. A controlled multicenter study. *Psychopharmacology* 90:131-138.
- Danish University Antidepressant Group (1990) Paroxetine: a selective serotonin reuptake inhibitor showing better tolerance, but weaker antidepressant effect than clomipramine in a controlled multicenter study. *J. Affect. Disord.* 18:289-299.
- De Chaffoy DE, Courcelles D, Leysen JE, De Clerck F, Van Belle H, Janssen PA (1985) Phospholipid turnover is the signal transducing system coupled to serotonin-S2 receptor sites. *J. Biol. Chem.* 260:7603-7608.
- De Felipe J, Arellano JI, Gomez A, Azmitia EC, Munoz A (2001) Pyramidal cell axons show a local specialization for GABA and 5-HT inputs in monkey and human cerebral cortex. *J. Comp. Neurol.* 433:148-155.
- De Quervain DJ, Henke K, Aerni A, Coluccia D, Wollmer MA, Hock C, Nitsch RM, Papassotiropoulos A (2003) A functional genetic variation of the 5-HT_{2A} receptor affects human memory. *Nat. Neurosci.* 6:1141-1142.
- DeFelipe J (2002) Cortical interneurons: from Cajal to 2001. *Prog. Brain Res.* 136:215-238.
- DeFelipe J, Hendry SH, Hashikawa T, Jones EG (1991) Synaptic relationships of serotonin-immunoreactive terminal baskets on GABA neurons in the cat auditory cortex. *Cereb. Cortex* 1:117-133.
- DeFelipe J, Alonso-Nanclares L, Arellano JI (2002) Microstructure of the neocortex: comparative aspects. *J. Neurocytol.* 31:299-316.
- Dégenètais E, Thierry AM, Glowinski J, Gioanni Y (2002) Electrophysiological properties of pyramidal neurons in the rat prefrontal cortex: an *in vivo* intracellular recording study. *Cereb. Cortex* 12:1-16.
- Delfs JM, Kong H, Mestek A, Chen Y, Yu L, Reisine T, Chesselet MF (1994) Expression of μ -opioid receptor mRNA in rat brain: an *in situ* hybridization study at the single cell level. *J. Comp. Neurol.* 345:46-68.
- Derkach V, Surprenant A, North RA (1989) 5-HT₃ receptors are membrane ion channels. *Nature* 339:706-709.
- Descarries L, Watkins KC, Garcia S, Beaudet A (1982) The serotonin neurons in nucleus raphe dorsalis of adult rat: a light and electron microscope radioautographic study. *J. Comp. Neurol.* 207:239-254.
- Doyle VM, Creba JA, Ruegg UT, Hoyer D (1986) Serotonin increases the production of inositol phosphates and mobilises calcium via de 5-HT₂ receptor in A7r5 smooth muscle cells. *Naunyn-Schmied. Arch. Pharmacol.* 333:98-103.

Drevets WC (2001) Functional anatomical abnormalities in limbic and prefrontal cortical structures in major depression. *Prog. Brain Res.* 126:413-31.

E

Elvevag B y Goldberg TE (2000) Cognitive impairment in schizophrenia is the core of the disorder. *Crit. Rev. Neurobiol.* 14:1-21.

Erickson JD, Eiden LE, Hoffman BJ (1992) Expression cloning of a reserpine-sensitive vesicular transporter. *Proc. Natl. Acad. Sci. USA* 89:10993-10997.

Erlander MG, Lovenberg TW, Baron BM, de Lecea L, Danielson PE, Racke M, Sloane AL, Siegel BW, Foye PE, Cannon K (1993) Two members of a distinct subfamily of 5-hydroxytryptamine receptors differentially expressed in rat brain. *Proc. Natl. Acad. Sci. USA* 90:3452-3456.

F

Fay R y Kubin L (2000) Pontomedullary distribution of 5-HT_{2A} receptor-like protein in the rat. *J. Comp. Neurol.* 418:323-345.

Férezou I, Cauli B, Hill EL, Rossier J, Hamel E, Lambolez B (2002) 5-HT₃ receptors mediate serotonergic fast synaptic excitation of neocortical vasoactive intestinal peptide/cholecystokinin interneurons. *J. Neurosci.* 22:7389-7397.

Ferré S, Cortés R, Artigas F (1994) Dopaminergic regulation of the serotonergic raphe-striatal pathway: microdialysis studies in freely moving rats. *J. Neurosci.* 14:4839-4846.

Foguet M, Nguyen H, Le H, Lubbert H (1992) Structure of the mouse 5-HT_{1C}, 5-HT₂ and stomach fundus serotonin receptor genes. *Neuroreport* 3:345-348.

Fozard JR (1989) The peripheral actions of 5-hydroxytryptamine. Oxford medical publications, Oxford.

Fuller JH y Schlag JD (1976) Determination of antidromic excitation by the collision test: problems of interpretation. *Brain Res.* 112:283-98.

Fuster JM (1997) The Prefrontal Cortex. Anatomy, Physiology and Neuropsychology of the Frontal Lobe. Philadelphia-New York: Lipincott-Raven.

- Fuster JM (2001) The prefrontal cortex--an update: time is of the essence. *Neuron* 30:319-333.
- Fuxe K (1965) Evidence for the existence of monoamine neurons in the central nervous system. 3. The monoamine nerve terminal. *Z. Zellforsch. Mikrosk. Anat.* 65:573-596.
- Fuxe K, Calza L, Benfenati F, Zini I, Agnati LF (1983) Quantitative autoradiographic localization of [³H]imipramine binding sites in the brain of the rat: relationship to ascending 5-hydroxytryptamine neuron systems. *Proc. Natl. Acad. Sci USA* 80:3836-3840.

G

- Gallager DW y Aghajanian GK (1976) Effect of antipsychotic drugs on the firing of dorsal raphe cells. II. Reversal by picrotoxin. *Eur. J. Pharmacol.* 39:357-364.
- Garratt JC, Kidd EJ, Wright IK, Marsden CA (1991) Inhibition of 5-hydroxytryptamine neuronal activity by the 5-HT agonist, DOI. *Eur. J. Pharmacol.* 199:349-355.
- Gartside SE, Hajos-Korcsok E, Bagdy E, Harsing LG, Sharp T, Hajós M (2000) Neurochemical and electrophysiological studies on the functional significance of burst firing in serotonergic neurons. *Neuroscience* 98:295-300.
- Geffard M, Tuffet S, Mons N, Chagnaud J-L (1987) Simultaneous detection of indoleamines and dopamine in rat dorsal raphe nuclei using specific antibodies. *Histochemistry* 88:61-64.
- Gershon MD, Liu KP, Karpiak SE, Tamir H (1983) Storage of serotonin *in vivo* as a complex with serotonin-binding protein in central and peripheral serotonergic neurons. *J. Neurosci.* 3:1901-1911.
- Glowinski J, Tassin JP, Thierry AM (1984) The mesocortico-prefrontal dopaminergic neurons. *Trends Neurosci.* 84:415-418.
- Goldman-Rakic PS (1995) Cellular basis of working memory. *Neuron* 14:477-485.
- Goldman-Rakic PS (1996) Regional and cellular fractionation of working memory. *Proc. Natl. Acad. Sci. USA* 93:13473-13480.
- Gras C, Herzog E, Bellenchi GC, Bernard V, Ravassard P, Pohl M, Gasnier B, Giros B, El Mestikawy S (2002) A third vesicular glutamate transporter expressed by cholinergic and serotonergic neurons. *J. Neurosci.* 22:5442-5451.

Groenewegen HJ y Uylings HB (2000) The prefrontal cortex and the integration of sensory, limbic and autonomic information. *Prog. Brain Res.* 126:3-28.:3-28.

Groenewegen HJ, Wright CI, Uylings HB (1997) The anatomical relationships of the prefrontal cortex with limbic structures and the basal ganglia. *J Psychopharmacol.* 11:99-106.

Grossman CJ, Kilpatrick GJ, Bunce KT (1993) Development of a radioligand binding assay for the 5-HT₄ receptors in guinea-pig and rat brain. *Br. J. Pharmacol.* 109:618-624.

Gurevich I, Tamir H, Arango V, Dwork AJ, Mann JJ, Schmauss C (2002) Altered editing of serotonin2C receptor pre-mRNA in the prefrontal cortex of depressed suicide victims. *Neuron* 34:349-356.

H

Hajós M y Sharp T (1996) Burst-firing activity of presumed 5-HT neurones of the rat dorsal raphe nucleus: Electrophysiological analysis by antidromic stimulation. *Brain Res.* 740:162-168.

Hajós M y Sharp T (1997) 5-HT neurones - Bursting with information? *Trends Neurosci.* 20:244-244.

Hajós M, Gartside SE, Villa AEP, Sharp T (1995) Evidence for a repetitive (burst) firing pattern in a sub-population of 5-hydroxytryptamine neurons in the dorsal and median raphe nuclei of the rat. *Neuroscience* 69:189-197.

Hajós M, Sharp T, Newberry NR (1996) Intracellular recordings from burst-firing presumed serotonergic neurones in the rat dorsal raphe nucleus *in vivo*. *Brain Res.* 737:308-312.

Hajós M, Richards CD, Szekely AD, Sharp T (1998) An electrophysiological and neuroanatomical study of the medial prefrontal cortical projection to the midbrain raphe nuclei in the rat. *Neuroscience* 87:95-108.

Hajós M, Hajos-Korcsok E, Sharp T (1999) Role of the medial prefrontal cortex in 5-HT_{1A} receptor-induced inhibition of 5-HT neuronal activity in the rat. *Br. J. Pharmacol.* 126:1741-1750.

Hall H, Farde L, Halldin C, Lundkvist C, Sedvall G (2000) Autoradiographic localization of 5-HT(2A) receptors in the human brain using [(3)H]M100907 and [(11)C]M100907. *Synapse* 38:421-431.

Hamada S, Senzaki K, Hamaguchi-Hamada K, Tabuchi K, Yamamoto H, Yamamoto T, Yoshikawa S, Okano H, Okado N (1998) Localization of 5-HT_{2A} receptor in rat cerebral cortex and olfactory system revealed by

- immunohistochemistry using two antibodies raised in rabbit and chicken. Mol. Brain Res. 54:199-211.
- Hamon M (1992) Central and peripheral 5-HT₃ receptors. London: Academic Press.
- Hamon M y Glowinski J (1974) Regulation of serotonin synthesis. Life Sci. 15:1533-1548.
- Hamon M, Bourgoin S, Glowinski J (1973) Feedback regulation of 5-HT synthesis in rat striatal slices. J. Neurochem. 20:1727-1745.
- Harder JA y Ridley RM (2000) The 5-HT_{1A} antagonist, WAY 100635, alleviates cognitive impairments induced by dizocilpine (MK-801) in monkeys. Neuropharmacology 39:547-52.
- Harvey JA, McMaster SE, Romano AG (1993) Methylenedioxymphetamine: neurotoxic effects on serotonergic projections to brainstem nuclei in the rat. Brain Res. 619:1-14.
- Heisler LK, Chu HM, Brennan TJ, Danao JA, Bajwa P, Parsons LH, Tecott LH (1998) Elevated anxiety and antidepressant-like responses in serotonin 5-HT_{1A} receptor mutant mice. Proc. Natl. Acad. Sci. USA 95:15049-15054.
- Higgins GA y Kilpatrick GJ (1999) 5-HT(3) receptor antagonists. Expert Opin. Investig. Drugs 8:2183-2188.
- Hjorth S y Magnusson T (1988) The 5-HT_{1A} receptor agonist 8-OH-DPAT preferentially activates cell body 5-HT autoreceptors in rat brain *in vivo*. Naunyn-Schmied. Arch. Pharmacol. 338:463-471.
- Hjorth S y Sharp T (1991) Effect of the 5-HT_{1A} receptor agonist 8-OH-DPAT on the release of 5-HT in dorsal and median raphe-innervated rat brain regions as measured by *in vivo* microdialysis. Life Sci. 48:1779-1786.
- Hjorth S y Tao R (1991) The putative 5-HT_{1B} receptor agonist CP-93,129 suppresses rat hippocampal 5-HT release *in vivo*: comparison with RU 24969. Eur. J. Pharmacol. 209:249-252.
- Hökfelt T, Johansson O, Fuxe K, Goldstein M, Park D (1976) Immunohistochemical studies on the localization and distribution of monoamine neuron systems in the rat brain. I. Tyrosine hydroxylase in the mesencephalon and diencephalon. Med. Biol. 54:427-453.
- Hoyer D y Schoeffter P (1988) 5-HT_{1D} Receptor-mediated inhibition of forskolin-stimulated adenylate cyclase activity in calf substantia nigra. Eur. J. Pharmacol. 147:145-147.
- Hoyer D, Engel G, Kalkman HO (1985) Molecular pharmacology of 5-HT₁ and 5-HT₂ recognition sites in rat and pig brain membranes: radioligand binding studies with (3H)5-HT, (3H)8-OH-DPAT, (-)

-)(125I)iodocyanopindolol, (3H)mesulergine and (3H)ketanserin. Eur. J. Pharmacol. 118:13-23.
- Hoyer D, Clarke DE, Fozard JR, Hartig PR, Martin GR, Mylecharane EJ, Saxena PR, Humphrey PPA (1994) International union of pharmacology classification of receptors for 5-Hydroxytryptamine (Serotonin). Pharmacol. Rev. 46:157-203.
- Hrdina PD, Foy B, Hepner A, Summers RJ (1990) Antidepressant binding sites in brain: autoradiographic comparison of [3H]paroxetine and [3H]imipramine, localization and relationship to serotonin transporter. J. Pharmacol. Exp. Ther. 252:410-418.
- Humphrey PP, Feniuk W, Perren MJ (1990a) Anti-migraine drugs in development: advances in serotonin receptor pharmacology. Headache 30:12-16.
- Humphrey PP, Feniuk W, Perren MJ, Beresford IJ, Skingle M, Whalley ET (1990b) Serotonin and migraine. Ann. N. Y. Acad. Sci. USA 600:587-598.
- Hutson PH, Sarna GS, O'Connell MT, Curzon G (1989) Hippocampal 5-HT synthesis and release *in vivo* is decreased by infusion of 8-OH-DPAT into the nucleus raphe dorsalis. Neurosci. Lett. 100:276-280.
-
- I**
- Ichikawa J, Ishii H, Bonaccorso S, Fowler WL, OLaughlin IA, Meltzer HY (2001) 5-HT_{2A} and D-2 receptor blockade increases cortical DA release via 5-HT_{1A} receptor activation: a possible mechanism of atypical antipsychotic-induced cortical dopamine release. J. Neurochem. 76:1521-1531.
- Innis RB, Nestler EJ, Aghajanian GK (1988) Evidence for G protein mediation of serotonin- and GABA_B-induced hyperpolarization of rat dorsal raphe neurons. Brain Res. 459:27-36.
- Invernizzi R, Carli M, Di Clemente A, Samanin R (1991) Administration of 8-Hydroxy-2-(Di-n-Propylamino)tetralin in raphe nuclei dorsalis and medianus reduces serotonin synthesis in the rat brain: differences in potency and regional sensitivity. J. Neurochem. 56:243-247.
- Iversen SD (1984) 5-HT and anxiety. Neuropharmacology 23:1553-1560.
-
- J**
- Jackson D y Abercrombie ED (1992) *In vivo* neurochemical evaluation of striatal serotonergic hyperinnervation in rats depleted of dopamine at infancy. J. Neurochem. 58:890-897.

- Jacobs BL y Azmitia EC (1992) Structure and function of the brain serotonin system. *Physiol. Rev.* 72:165-229.
- Jakab RL y Goldman-Rakic PS (1998) 5-Hydroxytryptamine(2A) serotonin receptors in the primate cerebral cortex: Possible site of action of hallucinogenic and antipsychotic drugs in pyramidal cell apical dendrites. *Proc. Natl. Acad. Sci. USA* 95:735-740.
- Jakab RL y Goldman-Rakic PS (2000) Segregation of serotonin 5-HT_{2A} and 5-HT₃ receptors in inhibitory circuits of the primate cerebral cortex. *J. Comp. Neurol.* 417:337-348.
- Jankowski MP y Sesack SR (2004) Prefrontal cortical proyections to the rat dorsal raphe nucleus: ultrastructural features and associations with serotonin and γ -aminobutyric acid neurons. *J. Comp. Neurol.* 468:518-529.
- Jansson A, Tinner B, Bancila M, Verge D, Steinbusch HW, Agnati LF, Fuxe K (2001) Relationships of 5-hydroxytryptamine immunoreactive terminal-like varicosities to 5-hydroxytryptamine-2A receptor-immunoreactive neuronal processes in the rat forebrain. *J. Chem. Neuroanat.* 22:185-203.
- Jodo E, Chiang C, Aston-Jones G (1998) Potent excitatory influence of prefrontal cortex activity on noradrenergic *locus coeruleus* neurons. *Neuroscience* 83:63-79.
- Johnson MD (1994) Synaptic glutamate release by postnatal rat serotonergic neurons in microculture. *Neuron* 12:433-442.
- Jolas T y Aghajanian GK (1997) Opioids suppress spontaneous and NMDA-induced inhibitory postsynaptic currents in the dorsal raphe nucleus of the rat *in vitro*. *Brain Res.* 755:229-245.
- Jolas T, Haj-Dahmane S, Lanfumey L, Fatacini CM, Kidd EJ, Adrien J, Gozlan H, Guardiolalemaître B, Hamon M (1993) (-)Tertatolol is a potent antagonist at presynaptic and postsynaptic serotonin 5-HT(1A) receptors in the rat brain. *Naunyn-Schmied. Arch. Pharmacol.* 347:453-463.

K

- Kalén P, Karlson M, Wiklund L (1985) Possible excitatory amino acid afferents to nucleus raphe dorsalis of the rat investigated with retrograde wheat germ agglutinin and D-[3H]aspartate tracing. *Brain Res.* 360:285-297.
- Kalén P, Skagerber G, Lindvall O (1988a) Projections from the ventral tegmental area and mesencephalic raphe to the dorsal raphe nucleus in the rat. Evidence for a minor dopaminergic component. *Exp. Brain Res.* 73:69-77.

- Kalén P, Strecker RE, Rosengren E, Björklund A (1988b) Endogenous release of neuronal serotonin and 5-hydroxyindoleacetic acid in the caudate-putamen of the rat as revealed by intracerebral dialysis coupled to high-performance liquid chromatography with fluorimetric detection. *J. Neurochem.* 51:1422-1435.
- Kalén P, Strecker RE, Rosengren E, Björklund A (1989) Regulation of striatal serotonin release by the lateral habenula-dorsal raphe pathway in the rat as demonstrated by *in vivo* microdialysis: role of excitatory amino acids and GABA. *Brain Res.* 492:187-202.
- Katz DM y Kimelberg HK (1985) Kinetics and autoradiography of high affinity uptake of serotonin by primary astrocyte cultures. *J. Neurosci.* 5:1901-1908.
- Kawa K (1994) Distribution and functional properties of 5-HT₃ receptors in the rat hippocampal dentate gyrus: a patch-clamp study. *J. Neurophysiol.* 71:1935-1947.
- Kawaguchi Y y Kondo S (2002) Parvalbumin, somatostatin and cholecystokinin as chemical markers for specific GABAergic interneuron types in the rat frontal cortex. *J. Neurocytol.* 31:277-287.
- Kawasaki T y Sato Y (1981) Afferent projections to the caudal part of the dorsal nucleus of the raphe in cats. *Brain Res.* 211:439-444.
- Kia HK, Brisorgueil MJ, Hamon M, Calas A, Verge D (1996a) Ultrastructural localization of 5-hydroxytryptamine(1A) receptors in the rat brain. *J. Neurosci. Res.* 46:697-708.
- Kia HK, Miquel MC, Brisorgueil MJ, Daval G, Riad M, Elmestikawy S, Hamon M, Vergé D (1996b) Immunocytochemical localization of serotonin(1A) receptors in the rat central nervous system. *J. Comp. Neurol.* 365:289-305.
- Kilpatrick GJ, Jones BJ, Tyers MB (1987) Identification and distribution of 5-HT₃ receptors in rat brain using radioligand binding. *Nature* 330:746-305.
- Kimelberg HK y Katz DM (1985) High-affinity uptake of serotonin into immunocytochemically identified astrocytes. *Science* 228:889-891.
- Kondo S y Kawaguchi Y (2001) Slow synchronized bursts of inhibitory postsynaptic currents (0.1-0.3 Hz) by cholinergic stimulation in the rat frontal cortex *in vitro*. *Neuroscience* 107:551-560.
- Kosofsky BE y Molliver ME (1987) The serotonergic innervation of cerebral cortex: Different classes of axon terminals arise from dorsal and median raphe nuclei. *Synapse* 1:153-168.
- Kreiss DS y Lucki I (1994) Differential regulation of 5-HT release in the striatum and hippocampus by 5-HT_{1A} autoreceptors of the dorsal and median raphe nuclei. *J. Pharmacol. Exp. Ther.* 269:1268-1279.

- Krimer LS y Goldman-Rakic PS (2001) Prefrontal microcircuits: membrane properties and excitatory input of local, medium, and wide arbor interneurons. *J. Neurosci.* 21:3788-3796.
- Kroeze WK y Roth BL (1998) The molecular biology of serotonin receptors: therapeutic implications for the interface of mood and psychosis. *Biol. Psychiatry* 44:1128-1142.
- Kühn DM, Wolf WA, Youdim MBH (1985) 5-Hydroxytryptamine release *in vivo* from cytoplasmic pool: studies on the 5-HT behavioural syndrome in reserpinized rats. *Br. J. Pharmacol.* 84:121-129.
- Kuroda M, Yokofujita J, Murakami K (1998) An ultrastructural study of the neural circuit between the prefrontal cortex and the mediodorsal nucleus of the thalamus. *Prog. Neurobiol.* 54:417-458.
- Kursar JD, Nelson DL, Wainscott DB, Cohen ML, Baez M (1992) Molecular cloning, functional expression, and pharmacological characterization of a novel serotonin receptor (5-hydroxytryptamine 2F) from rat stomach fundus. *Mol. Pharmacol.* 42:549-557.

L

- Lakoski JM y Aghajanian GK (1985) Effects of ketanserin on neuronal responses to serotonin in the prefrontal cortex, lateral geniculate and dorsal raphe nucleus. *Neuropharmacology* 24:265-273.
- Lambe EK, Goldman-Rakic PS, Aghajanian GK (2000) Serotonin induces EPSCs preferentially in layer V pyramidal neurons of the frontal cortex in the rat. *Cereb. Cortex*. 10:974-980.
- Lance JW (1993) Current concepts of migraine pathogenesis. *Neurology* 43:S11-S15.
- Laruelle M, Abi-Dargham A, van Dyck CH, Gil R, D'Souza CD, Erdos J, McCance E, Rosenblatt W, Fingado C, Zoghbi SS, Baldwin RM, Seibyl JP, Krystal JH, Charney DS, Innis RB (1996) Single photon emission computerized tomography imaging of amphetamine-induced dopamine release in drug-free schizophrenic subjects. *Proc. Natl. Acad. Sci. USA* 93:9235-9240.
- Lee EHY y Geyer MA (1984) Dopamine autoreceptor mediation of the effects of apomorphine on serotonin neurons. *Pharmacol. Biochem. Behav.* 21:301-311.
- Lee HS, Kim MA, Valentino RJ, Waterhouse BD (2003) Glutamatergic afferent projections to the dorsal raphe nucleus of the rat. *Brain Res.* 963:57-71.
- Leibowitz SF (1990) The role of serotonin in eating disorders. *Drugs* 39, suppl. 3:33-48.

Leonhardt S, Herrick-Davis K, Titeler M (1989) Detection of a novel serotonin receptor subtype (5-HT_{1E}) in human brain: interaction with a GTP-binding protein. *J. Neurochem.* 53:465-471.

Levine ES y Jacobs BL (1992) Neurochemical afferents controlling the activity of serotonergic neurons in the dorsal raphe nucleus: microiontophoretic studies in the awake cat. *J. Neurosci.* 12:4037-4044.

Li YQ, Li H, Kaneko T, Mizuno N (2001) Morphological features and electrophysiological properties of serotonergic and non-serotonergic projection neurons in the dorsal raphe nucleus. An intracellular recording and labeling study in rat brain slices. *Brain Res.* 900:110-118.

Liu R, Jolas T, Aghajanian G (2000) Serotonin 5-HT₂ receptors activate local GABA inhibitory inputs to serotonergic neurons of the dorsal raphe nucleus. *Brain Res.* 873:34-45.

López-Giménez JF, Mengod G, Palacios JM, Vilàro MT (1997) Selective visualization of rat brain 5-HT_{2A} receptors by autoradiography with [³H]MDL 100,907. *Naunyn Schmiedebergs Arch. Pharmacol.* 356:446-454.

Lovenberg TW, Erlander MG, Baron BM, Racke M, Slone AL, Siegel BW, Craft CM, Burns JE, Danielson PE, Sutcliffe JG (1993) Molecular cloning and functional expression of 5-HT_{1E}-like rat and human 5-Hydroxytryptamine receptor genes. *Proc. Natl. Acad. Sci. USA* 90:2184-2188.

M

Malenka RC y Nicoll JA (1999) Long-term potentiation--a decade of progress? *Science* 285:1870-184.

Mamounas LA y Molliver ME (1988) Evidence for dual serotonergic projections to neocortex: Axons from the dorsal and median raphe nuclei are differentially vulnerable to the neurotoxin p-chloroamphetamine (PCA). *Exp. Neurol.* 102:23-36.

Manrique C, Francoisbellan AM, Segu L, Becquet D, Hery M, Faudon M, Hery F (1994) Impairment of serotonergic transmission is followed by adaptive changes in 5HT(1B) binding sites in the rat suprachiasmatic nucleus. *Brain Res.* 663:93-100.

Mantz J, Godbour R, Tassin J-P, Glowinski J, Thierry AM (1990) Inhibition of spontaneous and evoked unit activity in the rat medial prefrontal cortex by mesencephalic raphe nuclei. *Brain Res.* 524:22-30.

Marek GJ y Aghajanian GK (1998) 5-hydroxytryptamine-induced excitatory postsynaptic currents in neocortical layer V pyramidal cells: suppression by mu-opiate receptor activation. *Neuroscience* 86:485-497.

- Marek GJ, Wright RA, Schoepp DD, Monn JA, Aghajanian GK (2000) Physiological antagonism between 5-hydroxytryptamine(2A) and group II metabotropic glutamate receptors in prefrontal cortex. *J. Pharmacol. Exp. Ther.* 292:76-87.
- Marek GJ, Wright RA, Gewirtz JC, Schoepp DD (2001) A major role for thalamocortical afferents in serotonergic hallucinogen receptor function in the rat neocortex. *Neuroscience* 105:379-392.
- Maricq AV, Peterson AS, Brake AJ, Myers RM, Julius D (1991) Primary structure and functional expression of the 5HT₃ receptor, a serotonin-gated ion channel. *Science* 254:432-437.
- Martín-Ruiz R y Ugedo L (2001) Electrophysiological evidence for postsynaptic 5-HT_{1A} receptor control of dorsal raphe 5-HT neurones. *Neuropharmacology* 41:72-78.
- Martín-Ruiz R, Ugedo L, Honrubia MA, Mengod G, Artigas F (2001) Control of serotonergic neurons in rat brain by dopaminergic receptors outside the dorsal raphe nucleus. *J. Neurochem.* 77:762-775.
- Matos FF, Urban C, Yocca FD (1996) Serotonin (5-HT) release in the dorsal raphe and ventral hippocampus: raphe control of somatodendritic and terminal 5-HT release. *J. Neural Transm.* 103:173-190.
- Maura G, Gemignani A, Versace P, Martire M, Raiteri M (1982) Carrier-mediated and carrier-independent release of serotonin from isolated central nerve endings. *Neurochem. Int.* 4:219-224.
- Maura G, Roccatagliata E, Raiteri M (1986) Serotonin autoreceptor in rat hippocampus: pharmacological characterization as a subtype of the 5-HT₁ receptor. *Naunyn-Schmied. Arch. Pharmacol.* 334:323-326.
- McCormick DA, Connors BW, Lighthall JW, Prince DA (1985) Comparative electrophysiology of pyramidal and sparsely spiny stellate neurons of the neocortex. *J. Neurophysiol.* 54:782-806.
- McMahon LL y Kauer JA (1997) Hippocampal interneurons are excited via serotonin-gated ion channels. *J. Neurophysiol.* 78:2493-2502.
- McQuade R y Sharp T (1995) Release of cerebral 5-hydroxytryptamine evoked by electrical stimulation of the dorsal and median raphe nuclei: effect of a neurotoxic amphetamine. *Neuroscience* 68:1079-1088.
- McQuade R y Sharp T (1997) Functional mapping of dorsal and median raphe 5-hydroxytryptamine pathways in forebrain of the rat using microdialysis. *J. Neurochem.* 69:791-796.
- Mello NK y Negus SS (2001) Effects of indatraline and buprenorphine on self-administration of speedball combinations of cocaine and heroin by rhesus monkeys. *Neuropsychopharmacology* 25:104-117.

- Meltzer HY (1999) The role of serotonin in antipsychotic drug action. *Neuropsychopharmacology* 21:S106-S115.
- Millan MJ (2000) Improving the treatment of schizophrenia: focus on serotonin (5-HT)(1A) receptors. *J. Pharmacol. Exp. Ther.* 295:853-861.
- Miller EK y Cohen JD (2001) An integrative theory of prefrontal cortex function. *Ann. Rev. Neurosci.* 24:167-202.
- Miller KJ y Teitler M (1992) Quantitative autoradiography of 5-CT-sensitive (5-HT_{1D}) and 5-CT-insensitive (5-HT_{1E}) serotonin receptors in human brain. *Neurosci. Lett.* 136:223-226.
- Miner LA, Backstrom JR, Sanders-Bush E, Sesack SR (2003) Ultrastructural localization of serotonin(2A) receptors in the middle layers of the rat prelimbic prefrontal cortex. *Neuroscience* 116:107-117.
- Miquel MC, Doucet E, Boni C, Elmestikawy S, Matthiessen L, Daval G, Verge D, Hamon M (1991) Central serotonin 1A receptors respective distributions of encoding messenger RNA, receptor protein and binding sites by *in situ* hybridization histochemistry, radioimmunohistochemistry and autoradiographic mapping in the rat brain. *Neurochem. Int.* 19:453-465.
- Misane I y Ogren SO (2003) Selective 5-HT_{1A} antagonists WAY 100635 and NAD-299 attenuate the impairment of passive avoidance caused by scopolamine in the rat. *Neuropsychopharmacology* 28:253-264.
- Molliver DC y Molliver ME (1990) Anatomic evidence for a neurotoxic effect of (+/-)-fenfluramine upon serotonergic projections in the rat. *Brain Res.* 511:165-168.
- Morales M y Bloom FE (1997) The 5-HT₃ receptor is present in different subpopulations of GABAergic neurons in the rat telencephalon. *J. Neurosci.* 17:3157-3167.
- Morales M, Battenberg E, Delecea L, Bloom FE (1996) The type 3 serotonin receptor is expressed in a subpopulation of GABAergic neurons in the rat neocortex and hippocampus. *Brain Res.* 731:199-202.
- Murphy DL, Zohar J, Benkelfat C, Pato MT, Pigott TA, Nsel TR (1989) Obsessive-compulsive disorder as a 5-HT subsystem-related behavioural disorder. *Br. J. Psychiatry* 155:15-24.

N

- Nanopoulos D, Belin MF, Maitre M, Vincendon G, Pujol JF (1982) Immunocytochemical evidence for the existence of GABAergic neurones in the nucleus raphe dorsalis. Possible existence of neurones containing serotonin and GABA. *Brain Res.* 232:375-389.

Nishikawa T y Scatton B (1985) Inhibitory influence of GABA on central serotonergic transmission. Raphe nuclei as the neuroanatomical site of the GABAergic inhibition of cerebral serotonergic neurons. Brain Res. 331:91-103.

O

O'Donnell P (2003) Dopamine gating of forebrain neural ensembles. Eur. J. Neurosci. 17:429-435.

O'Hearn E, Battaglia G, De Souza EB, Kuhar MJ, Molliver ME (1988) Methylenedioxymphetamine (MDA) and methylenedioxymethamphetamine (MDMA) cause selective ablation of serotonergic axon terminals in forebrain: immunocytochemical evidence for neurotoxicity. J. Neurosci. 8:2788-2803.

O'Hearn E y Molliver ME (1984) Organization of raphe-cortical projections in rat: A quantitative retrograde study. Brain Res. Bull. 13:709-726.

Oleskevich S y Descarries L (1990) Quantified distribution of the serotonin innervation in adult rat hippocampus. Neuroscience 34:19-33.

Olsten L y Seiger A (1972) Early prenatal ontogeny of central monoamine neurons in the rat: fluorescence histochemical observation. Anat. Entwicklungsgesch 137:301-316.

P

Palacios JM, Markstein R, Pazos A (1986) Serotonin-1C sites in the choroid plexus are not linked in a stimulatory or inhibitory way to adenylate cyclase. Brain Res. 380:151-154.

Pan ZZ y Williams JT (1989) GABA- and glutamate-mediated synaptic potentials in rat dorsal raphe neurons *in vitro*. J. Neurophysiol. 61:719-726.

Parks CL, Robinson PS, Sibille E, Shenk T, Toth M (1998) Increased anxiety of mice lacking the serotonin(1A) receptor. Proc. Natl. Acad. Sci. USA 95:10734-10739.

Paxinos G y Watson C (1986) The Rat Brain in Stereotaxic Coordinates. Sydney: Academic Press.

Paxinos G y Watson C (1998) The rat brain in stereotaxic coordinates. Sydney: Academic Press.

- Pazos A y Palacios JM (1985) Quantitative autoradiographic mapping of serotonin receptors in the rat brain. I. Serotonin-1 receptors. *Brain Res.* 346:205-230.
- Pazos A, Cortés R, Palacios JM (1985) Quantitative autoradiographic mapping of serotonin receptors in the rat brain. II. Serotonin-2 receptors. *Brain Res.* 346:231-249.
- Peroutka SJ y Snyder SH (1979) Multiple serotonin receptors: differential binding of [³H]5-hydroxytryptamine, [³H]lysergic acid, diethylamide and [³H]spiroperidol. *Mol. Pharmacol.* 16:687-699.
- Peters JA, Malone HM, Lambert JJ (1992) Recent advances in the electrophysiological characterization of 5-HT₃ receptors. *Trends Pharmacol. Sci.* 13:391-397.
- Peyron C, Luppi PH, Kitahama K, Fort P, Hermann DM, Jouvet M (1995) Origin of the dopaminergic innervation of the rat dorsal raphe nucleus. *Neuroreport* 6:2527-2531.
- Peyron C, Luppi PH, Fort P, Rampon C, Jouvet M (1996) Lower brainstem catecholamine afferents to the rat dorsal raphe nucleus. *J. Comp. Neurol.* 364:402-413.
- Peyron C, Petit JM, Rampon C, Jouvet M, Luppi PH (1998) Forebrain afferents to the rat dorsal raphe nucleus demonstrated by retrograde and anterograde tracing methods. *Neuroscience* 82:443-468.
- Pirot S, Jay TM, Glowinski J, Thierry AM (1994) Anatomical and electrophysiological evidence for an excitatory amino acid pathway from the thalamic mediodorsal nucleus to the prefrontal cortex in the rat. *Eur. J. Neurosci.* 6:1225-1234.
- Pompeiano M, Palacios JM, Mengod G (1992) Distribution and cellular localization of mRNA coding for 5-HT_{1A} receptor in the rat brain: correlation with receptor binding. *J. Neurosci.* 12:440-453.
- Pompeiano M, Palacios JM, Mengod G (1994) Distribution of the serotonin 5-HT₂ receptor family mRNAs: comparison between 5-HT_{2A} and 5-HT_{2C} receptors. *Mol. Brain Res.* 23:163-178.
- Portas CM y McCarley RW (1994) Behavioral state-related changes of extracellular serotonin concentration in the dorsal raphe nucleus: a microdialysis study in the freely moving cat. *Brain Res.* 648:306-312.
- Pritchett DB, Bach AW, Wozny M, Taleb O, Dal Toso R, Shih JC, Seuberg PH (1988) Structure and functional expression of cloned rat serotonin 5HT-2 receptor. *EMBO J.* 20:4135-4140.

Q

Qian Y, Melikian HE, Rye BR, Levey AI, Blakely RD (1995) Identification and characterization of antidepressant-sensitive serotonin transporter proteins using site-specific antibodies. *J. Neurosci.* 15:1261-1274.

R

Ramón y Cajal S (1911) *Histologie du système nerveux de l'homme et des vertébrés II*. Paris: Malone.

Ray JP y Price JL (1993) The organization of projections from the mediodorsal nucleus of the thalamus to orbital and medial prefrontal cortex in macaque monkeys. *J. Comp. Neurol.* 337:1-31.

Riad M, Garcia S, Watkins KC, Jodoin N, Doucet E, Langlois X, El Mestikawy S, Hamon M, Descarries L (2000) Somatodendritic localization of 5-HT_{1A} and preterminal axonal localization of 5-HT_{1B} serotonin receptors in adult rat brain. *J. Comp. Neurol.* 417:181-94.

Robbins TW (2000) From arousal to cognition: the integrative position of the prefrontal cortex. *Prog. Brain Res.* 126:469-83.:469-483.

Romero L, Celada P, Artigas F (1994) Reduction of *in vivo* striatal 5-hydroxytryptamine release by 8-OH-DPAT after inactivation of Gi/Go proteins in dorsal raphe nucleus. *Eur. J. Pharmacol.* 265:103-106.

Romero L, Jernej B, Bel N, Cicin-Sain L, Cortés R, Artigas F (1998) Basal and stimulated extracellular serotonin concentration in the brain of rats with altered serotonin uptake. *Synapse* 28:313-321.

Ropert N y Guy N (1991) Serotonin facilitates GABAergic transmission in the CA1 region of rat hippocampus *in vitro*. *J. Physiol* 441:121-136.

Rose JE y Woolsey CN (1948) The orbitofrontal cortex and its connections with the mediodorsal nucleus in rabbit, sheep and cat. *Res. Pub. Ass. Res. Nerv. Ment. Dis.* 27:210-232.

Roth BL, Craig SC, Choudhary MS, Uluer A, Monsma FJ, Jr., Shen Y, Meltzer HY, Sibley DR (1994) Binding of typical and atypical antipsychotic agents to 5-hydroxytryptamine-6 and 5-hydroxytryptamine-7 receptors. *J. Pharmacol. Exp. Ther.* 268:1403-1410.

Rudnick G y Wall SC (1992) p-Chloroamphetamine induces serotonin release through serotonin transporters. *Biochemistry* 31:6710-6718.

Rueter LE y Jacobs BL (1996) Changes in forebrain serotonin at the light-dark transition: Correlation with behaviour. *Neuroreport* 7:1107-1111.

S

- Sakai K, Salvert D, Touret M, Jouvet M (1977) Afferent connections of the nucleus raphe dorsalis in the cat as visualized by the horseradish peroxidase technique. *Brain Res.* 137:11-35.
- Santana N, Bortolozzi A, Serrats J, Mengod G, Artigas F (2004) Expression of serotonin_{1A} and serotonin_{2A} receptors in pyramidal and GABAergic neurons of the rat prefrontal cortex. *Cereb. Cortex* x:xx-xx.
- Sara SJ y Herve-Minvielle A (1995) Inhibitory influence of frontal cortex on *locus coeruleus* neurons. *Proc. Natl. Acad. Sci. USA.* 92:6032-6036.
- Sargent PA, Kjaer KH, Bench CJ, Rabiner EA, Messa C, Meyer J, Gunn RN, Grasby PM, Cowen PJ (2000) Brain serotonin_{1A} receptor binding measured by positron emission tomography with [11C]WAY-100635: effects of depression and antidepressant treatment. *Arch. Gen. Psychiatry* 57:174-180.
- Sawyer SF, Tepper JM, Young SJ, Groves PM (1985) Antidromic activation of dorsal raphe neurons from neostriatum: physiological characterization and effects of terminal autoreceptor activation. *Brain Res.* 332:15-28.
- Schoeffter P y Waeber C (1994) 5-Hydroxytryptamine receptors with a 5-HT₆ receptor-like profile stimulating adenylyl cyclase activity in pig caudate membranes. *Naunyn-Schmied. Arch .Pharmacol.* 350:356-360.
- Scruggs JL, Patel S, Bubser M, Deutch AY (2000) DOI-induced activation of the cortex: dependence on 5-HT_{2A} heteroreceptors on thalamocortical glutamatergic neurons. *J. Neurosci.* 20:8846-8852.
- Serrats J, Artigas F, Mengod G, Cortés R (2003) GABA_B receptor mRNA in the raphe nuclei: co-expression with serotonin transporter and glutamic acid decarboxylase. *J. Neurochem.* 84:743-752.
- Serrats J, Raurich A, Vilardo MT, Mengod G, Cortés R (2004) 5-HT_{5B} receptor mRNA in the raphe nuclei: coexpression with serotonin transporter. *Synapse* 51:102-111.
- Sesack SR y Carr DB (2002) Selective prefrontal cortex inputs to dopamine cells: implications for schizophrenia. *Physiol. Behav.* 77:513-517.
- Sesack SR, Deutch AY, Roth RH, Bunney BS (1989) Topographical organization of the efferent projections of the medial prefrontal cortex in the rat: an anterograde tract-tracing study with *Phaseolus vulgaris* leucoagglutinin. *J. Comp. Neurol.* 290:213-242.
- Sharp T, Bramwell SR, Clark D, Grahame-Smith DG (1989a) *In vivo* measurement of extracellular 5-hydroxytryptamine in hippocampus of the anaesthetized rat using microdialysis: changes in relation to 5-hydroxytryptaminergic neuronal activity. *J. Neurochem.* 53:234-240.

- Sharp T, Bramwell SR, Grahame-Smith DG (1989b) 5-HT₁ agonists reduce 5-hydroxytryptamine release in rat hippocampus *in vivo* as determined by brain microdialysis. Br. J. Pharmacol. 96:283-290.
- Sharp T, Bramwell SR, Grahame-Smith DG (1990) Release of endogenous 5-hydroxytryptamine in rat ventral hippocampus evoked by electrical simulation of the dorsal raphe nucleus as detected by microdialysis: sensitivity to tetrodotoxin, calcium and calcium antagonists. Neuroscience 39:629-637.
- Sheline YI, Sanghavi M, Mintun MA, Gado MH (1999) Depression duration but not age predicts hippocampal volume loss in medically healthy women with recurrent major depression. J. Neurosci. 19:5034-5043.
- Sinton CM y Fallon SL (1988) Electrophysiological evidence for a functional differentiation between subtypes of the 5-HT₁ receptor. Eur. J. Pharmacol. 157:173-181.
- Sirota P, Mosheva T, Shabtay H, Giladi N, Korczyn AD (2000) Use of the selective serotonin 3 receptor antagonist ondansetron in the treatment of neuroleptic-induced tardive dyskinesia. Am. J. Psychiatry 157:287-289.
- Sleight AJ, Carolo C, Petit N, Zwingelstein C, Bourson A (1995) Identification of 5-hydroxytryptamine(7) receptor binding sites in rat hypothalamus: sensitivity to chronic antidepressant treatment. Mol. Pharmacol. 47:99-103.
- Smiley JF y Goldman-Rakic PS (1996) Serotonergic axons in monkey prefrontal cerebral cortex synapse predominantly on interneurons as demonstrated by serial section electron microscopy. J. Comp. Neurol. 367:431-43.
- Smith GS, Savery D, Marden C, Lopez Costa JJ, Averill S, Priestley JV, Rattray M (1994) Distribution of messenger RNAs encoding enkephalin, substance P, somatostatin, galanin, vasoactive intestinal polypeptide, neuropeptide Y, and calcitonin gene-related peptide in the midbrain periaqueductal grey in the rat. J. Comp. Neurol. 350:23-40.
- Somogyi P, Tamas G, Lujan R, Buhl EH (1998) Salient features of synaptic organisation in the cerebral cortex. Brain Res. Rev. 26:113-135.
- Sotelo C, Cholley B, El Mestikawy S, Gozlan H, Hamon M (1990) Direct immunohistochemical evidence of the existence of 5-HT autoreceptors on serotonergic neurons in the midbrain raphe nuclei. Eur. J. Neurosci. 2:1144-1154.
- Sprouse JS y Aghajanian GK (1986) (-)-Propanolol blocks the inhibition of serotonergic dorsal raphe cell firing by 5-HT_{1A} selective agonists. Eur. J. Pharmacol. 128:295-298.
- Sprouse JS y Aghajanian GK (1987) Electrophysiological responses of serotonergic dorsal raphe neurons to 5-HT_{1A} and 5-HT_{1B} agonists. Synapse 1:3-9.

- Stamp JA y Semba K (1995) Extent of colocalization of serotonin and GABA in the neurons of the rat raphe nuclei. *Brain Res.* 677:39-49.
- Staubli U y Xu FB (1995) Effects of 5-HT₃ receptor antagonism on hippocampal theta rhythm, memory, and LTP induction in the freely moving rat. *J. Neurosci.* 15:2445-2452.
- Stern WC, Johnson A, Bronzino JD, Morgane PJ (1981) Neurpharmacology of the afferent projections from the lateral habenula and substantia nigra to the anterior raphe in the rat. *Neuropharmacology* 20:979-989.
- Strazielle C, Lalonde R, Hebert C, Reader TA (1999) Regional brain distribution of noradrenaline uptake sites, and of alpha1-alpha2- and beta-adrenergic receptors in PCD mutant mice: a quantitative autoradiographic study. *Neuroscience* 94:287-304.
- Svensson TH, Bunney BS, Aghajanian GK (1975) Inhibition of both noradrenergic and serotonergic neurons in brain by the alpha-adrenergic agonist clonidine. *Brain Res.* 92:291-306.
- Swanson LW (1998) Brain maps: structure of the rat brain. Amsterdam: Elsevier.

T

- Takagishi M y Chiba T (1991) Efferent projections of the infralimbic (area 25) region of the medial prefrontal cortex in the rat: an anterograde tracer PHA-L study. *Brain Res.* 566:26-39.
- Talvik-Lotfi M, Nyberg S, Nordstrom AL, Ito H, Halldin C, Brunner F, Farde L (2000) High 5HT_{2A} receptor occupancy in M100907-treated schizophrenic patients. *Psychopharmacology* 148:400-43.
- Tanaka E y North RA (1993) Actions of 5-hydroxytryptamine on neurons of the rat cingulate cortex. *J. Neurophysiol.* 69:1749-1757.
- Tao R y Auerbach SB (1996) Differential effect of NMDA on extracellular serotonin in rat midbrain raphe and forebrain sites. *J. Neurochem.* 66:1067-1075.
- Tao R y Auerbach SB (2000) Regulation of serotonin release by GABA and excitatory amino acids. *J. Psychopharmacol.* 14:100-113.
- Tao R y Auerbach SB (2003) Influence of inhibitory and excitatory inputs on serotonin efflux differs in the dorsal and median raphe nuclei. *Brain Res.* 961:109-120.
- Tao R, Ma Z, Auerbach SB (1996) Differential regulation of 5-hydroxytryptamine release by GABA_A and GABA_B receptors in midbrain raphe nuclei and forebrain of rats. *Br. J. Pharmacol.* 119:1375-1384.

- Tao R, Ma Z, Auerbach SB (1997) Influence of AMPA/kainate receptors on extracellular 5-hydroxytryptamine in rat midbrain raphe and forebrain. Br. J. Pharmacol. 121:1707-1715.
- Thiébot MH (1986) Are serotonergic neurons involved in the control of anxiety and in the anxiolytic activity of benzodiazepines? Pharmacol. Biochem. Behav. 24:1471-1477.
- Timmerman W y Westerink BHC (1997) Brain microdialysis of GABA and glutamate: What does it signify? Synapse 27:242-261.
- Toghi H, Abe T, Takahashi S, Kimura M, Takahashi J, Kikuchi T (1992) Concentrations of serotonin and its related substances in the cerebrospinal fluid in patients with Alzheimer type dementia. Neurosci. Lett. 141:9-12.
- Törk I (1990) Anatomy of the serotonergic system. Ann. N. Y. Acad. Sci. 600:9-34.

V

- Van der Kooy D, Hunt SP, Steinbusch HW, Verhofstad AA (1981) Separate populations of cholecystokinin and 5-hydroxytryptamine-containing neuronal cells in the rat dorsal raphe, and their contribution to the ascending raphe projections. Neurosci. Lett. 26:25-30.
- Van der Werf YD, Witter MP, Groenewegen HJ (2002) The intralaminar and midline nuclei of the thalamus. Anatomical and functional evidence for participation in processes of arousal and awareness. Brain Res. Rev. 39:107-140.
- VanderMaelen CP y Aghajanian GK (1983) Electrophysiological and pharmacological characterization of serotonergic dorsal raphe neurons recorded extracellular and intracellular in rat brain slices. Brain Res. 289:109-119.
- VanderMaelen CP, Matheson GK, Wilderman RC, Patterson LA (1986) Inhibition of serotonergic dorsal raphe neurons by systemic and iontophoretic administration of buspirone, a non-benzodiazepine anxiolytic drug. Eur. J. Pharmacol. 129:123-130.
- Varga V, Szekely AD, Csillag A, Sharp T, Hajós M (2001) Evidence for a role of GABA interneurons in the cortical modulation of midbrain 5-hydroxytryptamine neurones. Neuroscience 106:783-792.
- Vergé C, Daval G, Patey A, Gozlan H, El Mestikawy S, Hamon M (1985) Presynaptic 5-HT autoreceptors on serotonergic cell bodies and/or dendrites but not terminals are of the 5-HT_{1A} subtype. Eur. J. Pharmacol. 113:463-464.

Vertes RP (2004) Differential projections of the infralimbic and prelimbic cortex in the rat. *Synapse* 51:32-58.

W

Waeber C, Hoyer D, Palacios JM (1989) 5-Hydroxytryptamine3 receptors in the human brain: autoradiographic visualization using (3H)ICS 205-930. *Neuroscience* 31:393-400.

Waldmeier PC, Wicki P, Feldtrauer JJ, Baumann PA (1988) Potential involvement of a baclofen-sensitive autoreceptor in the modulation of the release of endogenous GABA from rat brain slices *in vitro*. *Naunyn-Schmied. Arch. Pharmacol.* 337:289-295.

Wang QP, Ochiai H, Nakai Y (1992) GABAergic innervation of serotonergic neurons in the dorsal raphe nucleus of the rat studied by electron microscopy double immunostaining. *Brain Res. Bull.* 29:943-948.

Wang RY y Aghajanian GK (1977) Antidromically identified serotonergic neurons in the rat midbrain raphe: evidence for collateral inhibition. *Brain Res.* 132:186-193.

Wang RY y Aghajanian GK (1977) Physiological evidence for habenula as major link between forebrain and midbrain raphe. *Science* 197:89-91.

Wang RY, Gallager DW, Aghajanian GK (1976) Stimulation of pontine reticular formation suppresses firing of serotonergic neurones in the dorsal raphe. *Nature* 264:365-368.

Weinberger DR, Aloia MS, Goldberg TE, Berman KF (1994) The frontal lobes and schizophrenia. *J. Neuropsychiatry Clin. Neurosci.* 6:419-427.

Wilkinson LO, Auerbach SB, Jacobs BL (1991) Extracellular serotonin levels change with behavioral state but not with pyrogen-induced hyperthermia. *J. Neurosci.* 11:2732-2741.

Williams GV y Goldman-Rakic PS (1995) Modulation of memory fields by dopamine D1 receptors in prefrontal cortex. *Nature* 376:572-575.

Williams GV, Rao SG, Goldman-Rakic PS (2002) The physiological role of 5-HT_{2A} receptors in working memory. *J. Neurosci.* 22:2843-2854.

Willins DL, Deutch AY, Roth BL (1997) Serotonin 5-HT_{2A} receptors are expressed on pyramidal cells and interneurons in the rat cortex. *Synapse* 27:79-82.

Wilson MA y Molliver ME (1991) The organization of serotonergic projections to cerebral cortex in primates retrograde transport studies. *Neuroscience* 44:555-570.

Wright IK, Garratt JC, Marsden CA (1990) Effects of a selective 5-HT₂ agonist, DOI, on 5-HT neuronal firing in the dorsal raphe nucleus and 5-HT release and metabolism in the frontal cortex. Br. J. Pharmacol. 99:221-222.

X

Xu T y Pandey SC (2000) Cellular localization of serotonin(2A) (5HT_(2A)) receptors in the rat brain. Brain Res. Bull. 51:499-505.

Xu ZQD y Hökfelt T (1997) Expression of galanin and nitric oxide synthase in subpopulations serotonin neurons of the rat dorsal raphe nucleus. J. Chem. Neuroanat. 13:169-187.

Y

Yang CR, Seamans JK, Gorelova N (1996) Electrophysiological and morphological properties of layers V-VI principal pyramidal cells in rat prefrontal cortex *in vitro*. J. Neurosci. 16:1904-1921.

Z

Zastawny RL, George SR, Nguyen T, Chen R, Tsatsos J, Briones-Urbina R, O'Dowd BF (1994) Cloning, characterization, and distribution of a μ-opiod receptor in rat brain. J. Neurochem. 62:2099-2105.

Zhou FM y Hablitz JJ (1999) Activation of serotonin receptors modulates synaptic transmission in rat cerebral cortex. J. Neurophysiol. 82:2989-2999.